

You are taking "Final Exam (TIMED)" as a timed exam. The timer on the right shows the time remaining in the exam. To receive credit for problems, you must select "Submit" for each problem before you select "End My Exam". **Show Less** 

**End My Exam** 

45:43:50 Ø

<u>Course</u> > <u>Final exam</u> > <u>Final Exam (TIMED)</u> > 4. F4.

## 4. F4.

4(a)

2/2 points (graded)

Find all negative real numbers  $\lambda_k$  for  $k=1,2,3,\ldots$  , for which the boundary value problem

$$egin{array}{lll} rac{d^2}{dx^2} v_k(x) &=& \lambda_k v_k(x) \ v_k(0) &=& 0 \ v_k(\pi/6) &=& 0 \end{array}$$

has a solution  $v_k(x)$  that is not the zero function.

(Note that  $\lambda_k$  is negative, k is nonnegative, and  $|\lambda_1|<|\lambda_2|<|\lambda_3|<\cdots$ .)

$$\lambda_k = oxed{-36 ^* ext{k^2}}$$

Find the k=1 eigenfunction  $v_{1}\left( x
ight)$  with amplitude 1.

$$v_1\left(x
ight) = egin{bmatrix} \sin\left(6^*\mathbf{x}
ight) \ & \sin\left(6\cdot x
ight) \end{bmatrix}$$

FORMULA INPUT HELP

Submit

You have used 1 of 3 attempts

✓ Correct (2/2 points)

## 4(b)

2/2 points (graded)

Find all negative real numbers  $\lambda_k$  for  $k=0,1,2,3,\ldots$  , for which the boundary value problem

$$egin{array}{lll} rac{d^2}{dx^2} v_k\left(x
ight) &=& \lambda_k v_k\left(x
ight) \ &v_k'\left(0
ight) &=& 0 \ &v_k\left(\pi
ight) &=& 0 \end{array}$$

has a solution  $v_k\left(x
ight)$  that is not the zero function.

(Note that  $\lambda_k$  is negative, k is nonnegative, and  $|\lambda_0|<|\lambda_1|<|\lambda_2|<\cdots$ .)

Find the k=0 eigenfunction  $v_{0}\left( x\right)$  with amplitude 1.



© All Rights Reserved