

MITx: 6.041x Introduction to Probability - The Science of Uncertainty



▶ Unit 0: Overview

- ▶ Entrance Survey
- ▶ Unit 1: Probability models and axioms
- ▶ Unit 2: Conditioning and independence
- ▶ Unit 3: Counting
- Unit 4: Discrete random variables
- Exam 1
- ▶ Unit 5: Continuous random variables
- Unit 6: Further topics on random variables
- **▼** Unit 7: **Bayesian** inference

Unit 7: Bayesian inference > Lec. 17: Linear least mean squares (LLMS) estimation > Lec 17 Linear least mean squares LLMS estimation vertical6

■ Bookmark

Exercise: Choice of representations

(1/1 point)

We wish to estimate an unknown quantity Θ . Our measuring equipment produces an observation of the form $X=\Theta^3+W$, where W is a noise term which is small relative to the range of Θ . Which type of linear estimator is preferable in such a situation?

$$\widehat{\Theta} = aX + b$$

$$\widehat{\Theta}=aX^3+b$$

$$oldsymbol{\widehat{\Theta}} = aX^{1/3} + b$$

Answer:

If the noise W were completely absent, we would estimate Θ by letting $\widehat{\Theta} = X^{1/3}$. In the presence of small noise, our estimator should again have a similar form, which argues in favor of the third option.

You have used 1 of 1 submissions

Unit overview

Lec. 14: Introduction to **Bayesian inference** Exercises 14 due Apr 06, 2016 at 23:59 UT 🗗

Lec. 15: Linear models with normal noise Exercises 15 due Apr 06, 2016 at 23:59 UT 4

Problem Set 7a Problem Set 7a due Apr 06, 2016 at 23:59 UTC

Lec. 16: Least mean squares (LMS) estimation Exercises 16 due Apr 13, 2016 at 23:59 UT 🗗

Lec. 17: Linear least mean squares (LLMS) estimation

Exercises 17 due Apr 13, 2016 at 23:59 UT (2)

Problem Set 7b Problem Set 7b due Apr 13, 2016 at 23:59 UTC

Solved problems

Additional theoretical material

Unit summary

© All Rights Reserved



© edX Inc. All rights reserved except where noted. EdX, Open edX and the edX and Open EdX logos are registered trademarks or trademarks of edX Inc.

















