

Ţ <u>Help</u>

sandipan\_dey >

**Discussion** <u>Syllabus</u> laff routines **Community** <u>Progress</u> <u>Outline</u> <u>Course</u> <u>Dates</u>

☆ Course / Week 9: Vector Spaces / 9.4 Vector Spaces

**(** 

< Previous 

Next >

### 9.4.4 The Null Space

☐ Bookmark this page

**■** Calculator

Week 9 due Dec 9, 2023 18:12 IST Completed

# 9.4.4 The Null Space

#### **Video**



Start of transcript. Skip to the end.

Robert van de Geijn: The second really important subspace of Rn

is known as the null space.

Recall.

We're interested in the solution of Ax equals b.

You've seen that if we have a specific solution

**▶** 0:00 / 0:00

▶ 2.0x









▲ Download video file

#### **Transcripts**

### Reading Assignment

0 points possible (ungraded) Read Unit 9.4.4 of the notes. [LINK]



Done



Submit

✓ Correct

#### Discussion

**Topic:** Week 9 / 9.4.4

**Hide Discussion** 

by recent activity >

Add a Post

Show all posts

There are no posts in this topic yet.

×

**⊞** Calculator

#### Homework 9.4.4.1

1/1 point (graded)

Let  $A \in \mathbb{R}^{m imes n}$  . The null space of A ,  $\mathcal{N}\left(A
ight)$  , is a subspace

TRUE ✓ ✓ Answer: TRUE

- $0 \in \mathcal{N}(A)$ : A0 = 0.
- If  $x,y\in\mathcal{N}\left(A\right)$  then  $x+y\in\mathcal{N}\left(A\right)$ : Let  $x,y\in\mathcal{N}\left(A\right)$  so that Ax=0 and Ay=0. Then  $A\left(x+y\right)=Ax+Ay=0+0=0$  which means means that  $x+y\in\mathcal{N}\left(A\right)$ .
- If  $lpha\in\mathbb{R}$  and  $x\in\mathcal{N}\left(A\right)$  then  $lpha x\in\mathcal{N}\left(A\right)$ : Let  $lpha\in\mathbb{R}$  and  $x\in\mathcal{N}\left(A\right)$  so that Ax=0. Then  $A\left(lpha x
  ight)=Alpha x=lpha Ax=lpha 0=0$  which means means that  $lpha x\in\mathcal{N}\left(A\right)$ .

Hence  $\mathcal{N}\left(A\right)$  is a subspace.

**Submit** 


Answers are displayed within the problem

#### Homework 9.4.4.2

8/8 points (graded)

Note: This exercise (and some in future units) does not seem to render right in Chrome. You may want to try another browser.

For each of the matrices on the left match the set of vectors on the right that describes its null space. (You should be able to do this "by examination.")



8. 
$$\begin{pmatrix} 2 & -4 \end{pmatrix}$$

✓ Answer: i

$$\text{(h)}\left\{\alpha\left(\frac{1}{2}\right)\middle|\alpha\in\mathbb{R}\right\}$$

$$\text{(i)}\left\{\alpha\left(\frac{2}{1}\right)\middle|\alpha\in\mathbb{R}\right\}$$

(Recall that V is the logical "or" operator.)

1. 
$$\begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$$
 **Answer:** (a) Any vector in  $\mathbb{R}^2$  maps to the zero vector.

2. 
$$\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$$
 Answer: (c)  $\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} \chi_0 \\ \chi_1 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$  means  $\chi_1 = 0$  with no restriction on  $\chi_0$ .

3. 
$$\begin{pmatrix} 0 & -2 \\ 0 & 0 \end{pmatrix}$$
 Answer: (c)  $\begin{pmatrix} 0 & -2 \\ 0 & 0 \end{pmatrix}$   $\begin{pmatrix} \chi_0 \\ \chi_0 \end{pmatrix}$  =  $\begin{pmatrix} 0 \\ 0 \end{pmatrix}$  means  $-2\chi_1 = 0$  with no restriction on  $\chi_0$ .

Previous Next >

© All Rights Reserved



### edX

**About** 

<u>Affiliates</u>

edX for Business

Open edX

**Careers** 

<u>News</u>

## Legal

Terms of Service & Honor Code

**Privacy Policy** 

**Accessibility Policy** 

**Trademark Policy** 

<u>Sitemap</u>

Cookie Policy

Your Privacy Choices

## Connect

<u>Idea Hub</u>

Contact Us

**Help Center** 

**Security** 

Media Kit

















© 2023 edX LLC. All rights reserved.

深圳市恒宇博科技有限公司 <u>粤ICP备17044299号-2</u>