

< Previous







Next >

# 12.4.1 Predicting the Weather, One Last Time

 Bookmark this page

Week 12 due Dec 29, 2023 10:42 IST

# 12.4.1 Predicting the Weather, One Last Time

## Video 12.4.1 Part 1

Homework (difficult)

If  $\lambda \in \Lambda(A)$  then  $\lambda \in \Lambda(A^T)$ .

True/False

11 / 44

important in a minute.

It sure seemed like there was a vector such that P times x was equal to x.

And therefore 1 would have to be an eigenvalue of the transition matrix P.

Why is this?

So here is a homework that actually is quite challenging.

So don't fret about this one too long.

**If you get frustrated, just go to the next video.**

### Video

Download video file

### Transcripts

Download SubRip (.srt) file

Download Text (.txt) file

## Reading Assignment

0 points possible (ungraded)

Read Unit 12.4.1 of the notes. [\[LINK\]](#)

☒ Done



Submit

☒ Correct

## Discussion

Topic: Week 12 / 12.4.1

Hide Discussion

Add a Post

Show all posts

by recent activity

HW 12.4.1.1: Explanation of the leap in logic

1

Type in PDF notes?

2

Calculator

Video 12.4.1 Part 2

Summary

► An understanding of eigenvalues and eigenvectors helped us understand why  $x^{(k)}$  eventually satisfies

$x^{(k)} \approx Px^{(k)}$

44 / 44

and you end up with a vector that points in the direction of the eigenvector  $v_0$ .

And bingo!

That explains the magic of predicting the weather many days from now.

So in summary, it's understanding eigenvalues and eigenvectors that allowed us to explain that when we keep hitting this vector  $x$  with  $P$ , eventually we end up with a vector that doesn't change anymore, **at least not noticeably.**

► 8:22 / 8:22

► 2.0x

🔊

⌵

CC

“

Video

📄 Download video file

Transcripts

📄 Download SubRip (.srt) file

📄 Download Text (.txt) file

Homework 12.4.1.1

1/1 point (graded)  
If  $\lambda \in \Lambda(A)$  then  $\lambda \in \Lambda(A^T)$ .

True ▼


✔ Answer: True

ANSWER: True

$\lambda \in \Lambda(A) \Rightarrow (A - \lambda I)$  is singular  
 $\Rightarrow$  < equivalent conditions >  
 $\dim(\mathcal{N}(A - \lambda I)) = k > 0$   
 $\Rightarrow$  <  $A$  is square, fundamental space picture >  
 $\dim(\mathcal{N}((A - \lambda I)^T)) = k > 0$   
 $\Rightarrow$  <  $(A - \lambda I)^T = A^T - \lambda I$  >  
 $\dim(\mathcal{N}(A^T - \lambda I)) = k > 0$   
 $\Rightarrow$  < equivalent conditions >  
 $(A^T - \lambda I)$  is singular  
 $\Rightarrow$  < property of eigenvalue >  
 $\lambda \in \Lambda(A^T)$

Submit

🧮 Calculator

 Answers are displayed within the problem

Homework 12.4.1.2

1/1 point (graded)

$\lambda \in \Lambda(A)$  if and only if  $\lambda \in \Lambda(A^T)$ .

TRUE 

 Answer: TRUE

This follows immediately from the last homework and the fact that  $(A^T)^T = A$ .

Submit

< Previous

Next >

© All Rights Reserved



edX

- [About](#)
- [Affiliates](#)
- [edX for Business](#)
- [Open edX](#)
- [Careers](#)
- [News](#)

Legal

- [Terms of Service & Honor Code](#)
- [Privacy Policy](#)
- [Accessibility Policy](#)
- [Trademark Policy](#)
- [Sitemap](#)
- [Cookie Policy](#)
- [Your Privacy Choices](#)

Connect

- [Idea Hub](#)
- [Contact Us](#)
- [Help Center](#)
- [Security](#)
- [Media Kit](#)



 Calculator

© 2023 edX LLC. All rights reserved.

深圳市恒宇博科技有限公司 [粤ICP备17044299号-2](#)