



You are taking "[Exam \(Timed, No Correctness Feedback\)](#)," as a timed exam. [Show more](#)

End My Exam

44:08:03

< Previous

✓

✓

✓

✓

✓

✓

✓

✓

✓

✓

✓

✓

✓

Next >

11. Applications to physics

Bookmark this page

Lecture due Aug 18, 2021 20:30 IST Completed




Apply

Find the vector decomposition

3.0/3 points (graded)
Consider a block of mass **1**kg sitting on a plane inclined to an angle of $\theta = \pi/6$. Approximate the force due to gravity to be **10**N pointing straight down. Find the vector decomposition into tangent and perpendicular vector components by following the method on the previous page.

(Decompose vectors with respect to the coordinate system where \hat{i} points horizontally to the right, and \hat{j} points vertically upwards.)



Find a vector \vec{F} representing the force due to gravity. (Enter vectors surrounded by square brackets: e.g. `[1,0]` .)

$\vec{F} =$ ✓ Answer: [0,-10]

Find a vector \vec{u} tangent to the inclined plane. Use the vector \vec{u} and the vector \vec{F} to find the vector \vec{a} that is the component of the vector \vec{F} tangent to the inclined plane.

(Enter vectors surrounded by square brackets: e.g. `[1,0]` .)

$\vec{a} =$ ✓ Answer: [-5*cos(pi/6),-5*sin(pi/6)]

Find the vector \vec{b} normal to the inclined plane so that $\vec{F} = \vec{a} + \vec{b}$. (Enter vectors surrounded by square brackets: e.g. `[1,0]` .)

$\vec{b} =$ ✓ Answer: [5*cos(pi/6),-15/2]

Solution:

The vector \vec{F} in the standard coordinate system is

$$\vec{F} = \langle 0, -10 \rangle.$$

A vector \vec{u} tangent to the inclined plane is

Calculator

Hide Notes

$$\vec{u} = \langle \cos(\pi/6), \sin(\pi/6) \rangle (= \langle \sqrt{3}/2, 1/2 \rangle).$$

Note that any vector that points in the same (or opposite) direction as this vector works!

Now to find the vector \vec{a} we use the dot product as in the worked example on the previous page.


$$\begin{aligned} \vec{a} &= \frac{\vec{F} \cdot \vec{u}}{\vec{u} \cdot \vec{u}} \vec{u} \\ &= \frac{\langle 0, -10 \rangle \cdot \langle \cos(\pi/6), \sin(\pi/6) \rangle}{\langle \cos(\pi/6), \sin(\pi/6) \rangle \cdot \langle \cos(\pi/6), \sin(\pi/6) \rangle} \langle \cos(\pi/6), \sin(\pi/6) \rangle \\ &= \frac{-5}{1} \langle \cos(\pi/6), \sin(\pi/6) \rangle = \langle -5\sqrt{3}/2, -5/2 \rangle. \end{aligned}$$

The vector \vec{b} is found using vector subtraction

$$\vec{b} = \vec{F} - \vec{a} = \langle 0, -10 \rangle - \langle -5\sqrt{3}/2, -5/2 \rangle = \langle 5\sqrt{3}/2, -15/2 \rangle.$$

Submit

You have used 2 of 10 attempts







 Answers are displayed within the problem

11. Applications to physics

Hide Discussion

Topic: Unit 2: Geometry of Derivatives / 11. Applications to physics

Add a Post

Show all posts	by recent activity
 <u>This one was poorly worded and confusing.</u> <u>I kept getting the wrong answer due to confusion with coordinates. Did anyone else have that issue? :(</u>	5
 <u>Where is the origin?</u> <u>Where is the origin? Any hints about this problem?</u>	2
 <u>units</u> <u>E, Ftan (= a) and Fnorm (= b) are forces, so they have to have a unit (Newton). But the system does not allow put a unit ("N", for exa...</u>	2
 <u>Strange wording and not enough tries</u> <u>I had a similar problem to a few of the others. It was strangely worded. It took me a while to finally figure out what you really wanted...</u>	5
 <u>[Staff] Just Say No to Default Coordinate Systems</u>	3
 <u>STAFF: The last question should be asking for b not a.</u> <u>The last question should be asking for b not a.</u>	6 new_ 11



edX

- [About](#)
- [Affiliates](#)
- [edX for Business](#)
- [Open edX](#)
- [Careers](#)
- [News](#)

Legal

- [Terms of Service & Honor Code](#)
- [Privacy Policy](#)
- [Accessibility Policy](#)
- [Trademark Policy](#)
- [Sitemap](#)

Connect

- [Blog](#)
- [Contact Us](#)
- [Help Center](#)
- [Media Kit](#)
- [Donate](#)



© 2021 edX Inc. All rights reserved.
深圳市恒宇博科技有限公司 [粤ICP备17044299号-2](#)



Calculator



Hide Notes