



[Lecture 14: Wald's Test, Likelihood Ratio Test, and Implicit Hypothesis](#)

4. Interlude: Square Roots of Matrices

[Course](#) > [Unit 4 Hypothesis testing](#) > [Test](#)

Audit Access Expires Dec 24, 2019

You lose all access to this course, including your progress, on Dec 24, 2019.

Upgrade by Nov 4, 2019 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)

4. Interlude: Square Roots of Matrices

Interlude: Square root of a positive semi-definite matrix

Recall that a matrix \mathbf{A} of size $d \times d$ is **positive semi-definite** if $\mathbf{x}^T \mathbf{A} \mathbf{x} \geq 0$ for all $\mathbf{x} \in \mathbb{R}^d$. Two example classes of positive semi-definite matrices are:

- Diagonal matrices with non-negative entries: $\mathbf{D} = \begin{pmatrix} c_1 & 0 & \dots & 0 \\ 0 & c_2 & & 0 \\ \vdots & & \ddots & \vdots \\ 0 & & \dots & c_d \end{pmatrix}$ where $c_i \geq 0$ for all i . (You have shown in exercise in a previous

lecture that indeed $\mathbf{x}^T \mathbf{D} \mathbf{x} \geq 0$ for all \mathbf{x} .

- Matrix products $\mathbf{P}^T \mathbf{D} \mathbf{P}$ where \mathbf{P} is an invertible (square) matrix and \mathbf{D} is a diagonal matrix with non-negative entries (as above). **Proof:**
 $\mathbf{x}^T (\mathbf{P}^T \mathbf{D} \mathbf{P}) \mathbf{x} = (\mathbf{P} \mathbf{x})^T \mathbf{D} (\mathbf{P} \mathbf{x}) = \mathbf{y}^T \mathbf{D} \mathbf{y} \geq 0$ for all vectors \mathbf{x} .

Generating Speech Output

The **positive semi-definite square root** (or simply the square root) of a positive semi-definite matrix \mathbf{A} is another positive semi-definite matrix, denoted by $\mathbf{A}^{1/2}$, satisfying $\mathbf{A}^{1/2} \mathbf{A}^{1/2} = \mathbf{A}$. It is the case that for any positive semi-definite matrix (positive definite matrix, respectively), the positive semi-definite square root (positive definite square root, respectively) is unique.

Square Root of a Matrix

1/1 point (graded)

Using the definition above of the square root of a matrix, find the square root $\mathbf{D}^{1/2}$ of $\mathbf{D} = \begin{pmatrix} 2 & 0 \\ 0 & 0 \end{pmatrix}$.

(Enter your answer as a matrix, e.g. by typing `[[1,2],[5,1]]` for the matrix $\begin{pmatrix} 1 & 2 \\ 5 & 1 \end{pmatrix}$. Note the square brackets, and the commas as separators.)

$\mathbf{D}^{1/2} =$

`[[sqrt(2),0],[0,0]]`

✓ Answer: `[[sqrt(2),0],[0,0]]`

STANDARD NOTATION

Solution:

Since

$$\begin{pmatrix} \sqrt{2} & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} \sqrt{2} & 0 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} 2 & 0 \\ 0 & 0 \end{pmatrix},$$

We have $\mathbf{D}^{1/2} = \begin{pmatrix} \sqrt{2} & 0 \\ 0 & 0 \end{pmatrix}$.

Submit

You have used 2 of 3 attempts

❗ Answers are displayed within the problem

Generating Speech Output

(Optional): Square Root of a Matrix

0 points possible (ungraded)

Let

$$\mathbf{A} = \mathbf{P}^T \mathbf{D} \mathbf{P} \quad \text{where} \quad \mathbf{D} = \begin{pmatrix} 3 & 0 \\ 0 & 0 \end{pmatrix}$$
$$\mathbf{P} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix}.$$

Note that $\mathbf{P}^T = \mathbf{P}^{-1}$.

Find the square root $\mathbf{A}^{1/2}$ of the matrix \mathbf{A} .

Hint: $\mathbf{P}^T \mathbf{B}^2 \mathbf{P} = \mathbf{P}^T \mathbf{B} (\mathbf{P} \mathbf{P}^T) \mathbf{B} \mathbf{P}$.

(Enter your answer as a matrix, e.g. by typing `[[1,2],[5,-1]]` for the matrix $\begin{pmatrix} 1 & 2 \\ 5 & -1 \end{pmatrix}$. Note the square brackets, and commas as separators.)

$\mathbf{A}^{1/2} =$

[[sqrt(3)/2,-sqrt(3)/2],[-sqrt(3)/2,sqrt(3)/2]]



Answer: sqrt(3)/2*[[1,-1],[-1,1]]

STANDARD NOTATION

Solution:

$$\begin{aligned} (\mathbf{P}^T \mathbf{D}^{1/2} \mathbf{P}) (\mathbf{P}^T \mathbf{D}^{1/2} \mathbf{P}) &= \mathbf{P}^T \mathbf{D}^{1/2} (\mathbf{P} \mathbf{P}^T) \mathbf{D}^{1/2} \mathbf{P} \\ &= \mathbf{P}^T \mathbf{D}^{1/2} (\mathbf{P} \mathbf{P}^{-1}) \mathbf{D}^{1/2} \mathbf{P} \quad \text{since } \mathbf{P}^T = \mathbf{P}^{-1} \\ &= \mathbf{P}^T \mathbf{D}^{1/2} \mathbf{D}^{1/2} \mathbf{P} \\ &= \mathbf{P}^T \mathbf{D} \mathbf{P} \end{aligned}$$

Hence $\mathbf{A}^{1/2} = \mathbf{P}^T \mathbf{D}^{1/2} \mathbf{P}$. Plugging in the values of \mathbf{D} and \mathbf{P} , we get

Generating Speech Output

$$\begin{aligned}
 \mathbf{A}^{1/2} &= \mathbf{P}^T \mathbf{D}^{1/2} \mathbf{P} = \left(\frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix} \right) \begin{pmatrix} \sqrt{3} & 0 \\ 0 & 0 \end{pmatrix} \left(\frac{1}{\sqrt{2}} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix} \right) \\
 &= \frac{1}{2} \begin{pmatrix} \sqrt{3} & 0 \\ -\sqrt{3} & 0 \end{pmatrix} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix} \\
 &= \frac{\sqrt{3}}{2} \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix}
 \end{aligned}$$

Submit

You have used 1 of 3 attempts

i Answers are displayed within the problem

Discussion

Hide Discussion

Topic: Unit 4 Hypothesis testing: Lecture 14: Wald's Test, Likelihood Ratio Test, and Implicit Hypothesis Test / 4.
Interlude: Square Roots of Matrices

Add a Post

Show all posts ▼

by recent activity ▼

There are no posts in this topic yet.

✕

Learn About Verified Certificates

© All Rights Reserved

Generating Speech Output