

edX and its Members use cookies and other tracking technologies for performance, analytics, and marketing purposes. By using this website, you accept this use. Learn more about these technologies in the [Privacy Policy](#).



[Homework 6 Maximum Likelihood](#)

[Course](#) > [Unit 3 Methods of Estimation](#) > [Estimation and Method of Moments](#) > 5. Censored data

Audit Access Expires Dec 24, 2019

You lose all access to this course, including your progress, on Dec 24, 2019.

Upgrade by Nov 4, 2019 to get unlimited access to the course as long as it exists on the site. [Upgrade now](#)

5. Censored data

In a given population, n individuals are sampled randomly, with replacement, and each sampled individual is asked whether his/her salary is greater than some fixed threshold z . Assume that the salary of a randomly chosen individual has the exponential distribution with unknown parameter λ . Asking whether the salary overcomes a given threshold rather than directly asking for the salary increases the number people that are willing to answer and decreases the number of mistakes in the collected answers.

Denote by X_1, \dots, X_n the binary responses of the n sampled individuals, so that $X_i \in \{0, 1\}$. We call the X_i **censored data**.

(a)

2/2 points (graded)

What kind of distribution do the X_i s follow?



Exponential distribution with parameter $\mu(\lambda)$

Generating Speech Output

☒ Bernoulli with parameter $\mu(\lambda)$

☐ Poisson with parameter $\mu(\lambda)$



Give the parameter of this distribution in terms of λ and z :

Parameter $\mu(\lambda) =$

$e^{-(\lambda \cdot z)}$



$e^{-\lambda \cdot z}$

Submit

You have used 1 of 2 attempts

✓ Correct (2/2 points)

(b)

1/1 point (graded)

Let \bar{X}_n be the proportion of sampled individuals whose response was 1 (corresponding to Yes). Convince yourself that \bar{X}_n is asymptotically normal.

What is its asymptotic variance?

$V(\bar{X}_n) =$

$e^{-(\lambda \cdot z)} \cdot (1 - e^{-(\lambda \cdot z)})$



$e^{-\lambda \cdot z} \cdot (1 - e^{-\lambda \cdot z})$

Submit

You have used 1 of 3 attempts

Generating Speech Output

✓ Correct (1/1 point)

(c)

1/1 point (graded)

Find a function f such that $f(\bar{X}_n)$ is a consistent estimator of λ .

Write **barX_n** for the sample average \bar{X}_n .

$$f(\bar{X}_n) = -\ln(\text{barX}_n)/z$$

✓

$-\frac{\ln(\text{barX}_n)}{z}$

Submit

You have used 1 of 3 attempts

✓ Correct (1/1 point)

(d)

1/1 point (graded)

Convince yourself that $f(\bar{X}_n)$ is asymptotically normal and compute its asymptotic variance.

$$V(f(\bar{X}_n)) = (e^{(\lambda z)} - 1)/z^2$$

✓

$\frac{e^{\lambda z} - 1}{z^2}$

Submit

You have used 1 of 3 attempts

Generating Speech Output

✓ Correct (1/1 point)

(e)

1/1 point (graded)

What equation must z satisfy in order to minimize the asymptotic variance computed in part (d)? Write this equation in the form $g_\lambda(z) = z$, where g_λ is a function that depends on the unknown parameter λ .

$$g_\lambda(z) = \frac{2 \cdot (1 - e^{-\lambda z})}{\lambda}$$

Submit

You have used 2 of 3 attempts

✓ Correct (1/1 point)

(f)

1/1 point (graded)

Let Y_1, \dots, Y_n be the salaries of the n sampled people. If one could actually observe Y_1, \dots, Y_n , what would be the Fisher information of Y , $I_Y(\lambda)$, depending on λ ?

$$I_Y(\lambda) = \frac{1}{\lambda^2}$$

Submit

You have used 1 of 3 attempts

Generating Speech Output

✓ Correct (1/1 point)

(g)

1/1 point (graded)

In the model where only the X_i 's are observed (with fixed threshold z), what is the Fisher information? Denote it by $I_X(\lambda)$.

$I_X(\lambda)$ ✓

$$\frac{z^2}{e^{\lambda \cdot z} - 1}$$

Submit

You have used 2 of 3 attempts

✓ Correct (1/1 point)

(h)

2/2 points (graded)

Compare $I_Y(\lambda)$ and $I_X(\lambda)$:

☒ $I_Y(\lambda) \geq I_X(\lambda)$ for all λ

☐ $I_Y(\lambda) \leq I_X(\lambda)$ for all λ

☐ $I_Y(\lambda) \geq I_X(\lambda)$ for some λ , $I_Y(\lambda) < I_X(\lambda)$ for others.

✓

Generating Speech Output interpret this in this model?

☐ It depends on the parameter λ whether the censored data or the actual data provides a better estimate.

☒ The actual data always provides a better estimate

☐ The censored data always provides a better estimate.



Submit

You have used 1 of 2 attempts

✓ Correct (2/2 points)

Discussion

Hide Discussion

Topic: Unit 3 Methods of Estimation: Homework 6 Maximum Likelihood Estimation and Method of Moments / 5.
Censored data

Add a Post

Show all posts ▼

by recent activity ▼

? [Parts e,g](#)

4

[For part e, not getting a neat answer from a solver. Maybe am way off in the method. For part g, compared to similar version of same problem with the parameter from this o...](#)

💬 [e\) trouble with Lambert W function](#)

1

[Hey, I got a green mark on d\). So e\) seemed quite trivial. But my function involves the Lambert W function \(that surprised me\) or to be precise an approximation of the Output v...](#)

Learn About Verified Certificates

© All Rights Reserved

Generating Speech Output