



Bookmarks

- [Module 1: The Basics of R and Introduction to the Course](#)
- [Entrance Survey](#)
- [Module 2: Fundamentals of Probability, Random Variables, Distributions, and Joint Distributions](#)
- [Module 3: Gathering and Collecting Data, Ethics, and Kernel Density Estimates](#)
- [Module 4: Joint, Marginal, and Conditional Distributions & Functions of Random Variable](#)

Module 6: Special Distributions, the Sample Mean, the Central Limit Theorem, and Estimation > Human Subjects and Special Distributions > The Negative Binomial Distribution - Quiz

## The Negative Binomial Distribution - Quiz

🔖 Bookmark this page

### Question 1

1.0 point possible (graded)

Which of the following examples would be useful to model using the negative binomial distribution? (Select all that apply.)


- ☐ a. The number of shots that a basketball player makes out of the next 40 attempts
- ☒ b. The number of times that you can operate a machine before there is an issue with the machine and the machine breaks down
- ☒ c. The number of people that you would need to survey to reach a certain number of respondents that are male and above the age of 65.
- ☒ d. The number of cards you sample from a deck of 52 before drawing an ace




► [Module 5: Moments of a Random Variable, Applications to Auctions, & Intro to Regression](#)

▼ [Module 6: Special Distributions, the Sample Mean, the Central Limit Theorem, and Estimation](#)


### **Human Subjects and Special Distributions**

Finger Exercises due Nov 07, 2016 at 05:00 IST 

### **The Sample Mean, Central Limit Theorem, and Estimation**

Finger Exercises due Nov 07, 2016 at 05:00 IST 

### **Module 6: Homework**

Homework due Oct 31, 2016 at 05:00 IST 

► [Module 7: Assessing and Deriving Estimators - Confidence Intervals, and Hypothesis Testing](#)

Submit

You have used 1 of 2 attempts

✓ Correct (1/1 point)

## Question 2

0 points possible (ungraded)

Let's expand on the example described above of surveying homes to reach 100 males over the age of 65. Suppose that you know that in the population of interest, the probability that a home will have one male resident above the age of 65 is 15% or .15. (Assume that all homes either have 0 or 1 men over 65.) What is the expected number of homes you will need to visit before reaching 100 males above 65?

*Please round your answer to the nearest whole number (no decimals).*

567

✓ Answer: 567

567

## Explanation

The average number of homes that you will need to visit can be calculated as:

$$E[X] = 100(1 - p)/p + 100 = 100/0.15 = 667.$$

► [Exit Survey](#)

Submit

You have used 1 of 2 attempts

✓ Correct

### Discussion

**Topic:** Module 6 / The Negative Binomial Distribution - Quiz

Show Discussion

© All Rights Reserved



© 2016 edX Inc. All rights reserved except where noted. EdX, Open edX and the edX and Open EdX logos are registered trademarks or trademarks of edX Inc.

POWERED BY  
**OPENedX**



