








< Previous



Next >

12.2.1 Explicit and Implicit Methods

 Bookmark this page

MO2.7

The stability issue observed with the Forward Euler method turns out to be a problem for the other methods we have also learned about (i.e. Runge-Kutta methods and leap frog). The common feature all of these methods is that they all extrapolate from the current state \underline{v}^n and the current forcing $\underline{f}(\underline{v}^n, t^n)$ to estimate \underline{v}^{n+1} . This is clear for Forward Euler,

$$\underline{v}^{n+1} = \underline{v}^n + \Delta t \underline{f}(\underline{v}^n, t^n)$$

(12.21)

where \underline{v}^{n+1} is explicitly calculated from \underline{v}^n and $\underline{f}(\underline{v}^n, t^n)$ and hence Forward Euler (and our other methods so far) are known as *explicit* methods.

A different approach is to use an *implicit* method, the simplest implicit method the Backward Euler method given by:

$$\underline{v}^{n+1} = \underline{v}^n + \Delta t \underline{f}(\underline{v}^{n+1}, t^{n+1})$$

(12.22)

© All Rights Reserved



[About](#)
[Affiliates](#)
[edX for Business](#)
[Open edX](#)
[Careers](#)
[News](#)



[Terms of Service & Honor Code](#)
[Privacy Policy](#)
[Accessibility Policy](#)
[Trademark Policy](#)
[Sitemap](#)
[Cookie Policy](#)
[Your Privacy Choices](#)



[Blog](#)
[Contact Us](#)
[Help Center](#)
[Security](#)
[Media Kit](#)

Discussions

All posts sorted by recent activity



Next >



© 2023 edX LLC. All rights reserved.

深圳市恒宇博科技有限公司 [粤ICP备17044299号-2](#)

