<u>Help</u>

sandipan\_dey >

<u>Calendar</u> **Discussion** <u>Notes</u> <u>Course</u> <u>Progress</u> <u>Dates</u>

☆ Course / Unit 2: Geometry of Derivat... / Lecture 4: Introduction to vectors and dot pro...



You are taking "Exam (Timed, No Correctness Feedback)" as a timed exam. Show more

**End My Exam** 

44:21:47





Lecture due Aug 18, 2021 20:30 IST Completed



**Practice** 

### **Vector magnitude**



PROFESSOR: Speaking of magnitude,

Start of transcript. Skip to the end.

we write it this way.

or length,

The norm of the vector, the absolute value of a vector,

this means the length of the vector, also called the magnitude.

And in terms of v1 and v2, we can find it from the picture

0:00 / 0:00

▶ 2.0x

X


CC 66

**Video** Download video file **Transcripts** Download SubRip (.srt) file Download Text (.txt) file

**Definition 4.1** The **magnitude** of a vector  $\vec{v}$  is equal to its length and is denoted by  $|\vec{v}|$ .

We can find the magnitude of the vector  $ec{v}=\langle v_1,v_2
angle$  by using the Pythagorean theorem. We draw a triangle with base  $v_1$  , height  $v_2$  , and hypotenuse |v| . Then the magnitude of the vector  $ec{v}=\langle v_1,v_2
angle$  is given by

$$|\vec{v}| = \sqrt{v_1^2 + v_2^2}. (3.3)$$



#### **→** Spoiler: Magnitude in higher dimensions

Consider a vector with n components given by  $ec{v}=\langle v_1,v_2,\ldots,v_n
angle$ . The magnitude of  $ec{v}$  is given by

$$|\vec{v}| = \sqrt{v_1^2 + v_2^2 + \dots + v_n^2}. (3.4)$$

<u>Hide</u>

**A note about terminology:** In some videos, the professor uses the word "norm" when referring to the magnitude. This is because the magnitude of a vector is given by what is called the Euclidean norm. There are many different kinds of norms used to measure different mathematical objects such as numbers, vectors, and functions.

### Vector magnitude practice 1

3/3 points (graded)

Find the magnitude of the following vectors.

(Note you may enter math expressions or numbers. You need only enter decimals to 2 decimal places.)

### Solution:

From the definition of magnitude, we have

$$ert ec v ert \ = \ ert \langle 3,4 
angle ert = \sqrt{3^2 + 4^2} = \sqrt{25} = 5$$
  $ert ec w ert \ = \ ert \langle -2,0 
angle ert = \sqrt{(-2)^2 + 0^2} = \sqrt{4} = 2$   $ert ec u ert \ = \ ert \langle -1,1,-10 
angle ert = \sqrt{(-1)^2 + (1)^2 + (-10)^2} = \sqrt{102}.$ 

Submit


You have used 1 of 5 attempts

**1** Answers are displayed within the problem

## Vector magnitude practice 2

1/1 point (graded)

Find the magnitude of the vector drawn below.



(Note you may enter math expressions or numbers. You need only enter decimals to 2 decimal places.)

 $|\vec{v}| =$ **✓ Answer:** sqrt(1.25) 1.118034

### **Solution:**

The vector plotted is given by  $ec{v}=\langle 1,-0.5 
angle$  . The magnitude is therefore

$$|\vec{v}| = |\langle 1, -0.5 \rangle| = \sqrt{1^2 + 0.5^2} = \sqrt{1.25}.$$
 (3.5)

Submit

You have used 1 of 3 attempts

**1** Answers are displayed within the problem

### 4. Magnitude

Topic: Unit 2: Geometry of Derivatives / 4. Magnitude

**Hide Discussion** 

**Add a Post** 

Show all posts $\checkmark$ by recent activity		ity 🗸
•	tick marks on line above not working  Although I correctly answered the questions in this particular section, the tick marks for these exercises are not showing on the top I	7
<b>Q</b>	Why does the pythagorean theorem work beyond two dimensions?  Could someone provide some insights on this matter? Thanks in advance!	4


Previous	Next >
----------	--------

© All Rights Reserved



edX

**About Affiliates** edX for Business



Open edX

Careers

<u>News</u>

# Legal

Terms of Service & Honor Code

**Privacy Policy** 

**Accessibility Policy** 

**Trademark Policy** 

<u>Sitemap</u>

## **Connect**

<u>Blog</u>

**Contact Us** 

Help Center

Media Kit

**Donate** 















© 2021 edX Inc. All rights reserved.

深圳市恒宇博科技有限公司 <u>粤ICP备17044299号-2</u>