



You are taking "[Exam \(Timed, No Correctness Feedback\)](#)," as a timed exam. [Show more](#)

End My Exam

44:32:45



< Previous



Next >

10. Variables and units

Bookmark this page



Calculator



Hide Notes

Lecture due Aug 4, 2021 20:30 IST Completed



Reflect

In the first lecture, we mentioned that the world around us is multivariable, and that multivariable functions can be used to represent the world around us. So far, we've been thinking about x , y , and z as spatial variables, which we often expect to have units like meters, or inches. Let's think about what this means for how we represent functions first.

Suppose that x and y represent position in units of meters. And $z = f(x, y)$ is an altitude measured again in meters. Let's look at our first example function $z = x^2 + y^2$.

Both x^2 and y^2 have units of square meters. How can z have units of meters as we expect?

Secretly, there must be constant multiples that also have units embedded within this function that help all of the units to make sense. Really, what we have is

$$z = 1x^2 + 1y^2$$

where both of the constants 1 have units of 1/meters. In this way, every term in this function has the same units.

General rules for units of multivariable functions

- 1. All terms that are added together or equated in a multivariable function must have the same units. That is, we cannot add a term with units of meters to a term with units of square meters. It doesn't have a physical meaning that we can interpret.
- 2. Functions like \sin , \cos , \ln , $e^{f(x,y)}$ must have input arguments that are unit-less.
- 3. Taking derivatives changes units. (See exercise below.)

▼ (Optional) Remark about unit-less input arguments

If you think about expressing a function like $\sin(f(x))$ in one variable as their Taylor series, it makes sense that an infinite sum of terms must have every term unit-less. Otherwise, we would break the first rule above. The same holds for multivariable functions with multivariable Taylor series!

Hide

Example 10.1

Consider the ideal gas law

$$PV = nRT$$

This relationship involves 5 variables:

Variable	Quantity	Units
P	pressure	Newtons per square meter (N/m^2)
V	volume	cubic meters (m^3)

Calculator

Hide Notes

n	amount	number of moles (mol)
R	ideal gas constant	(Exercise below)
T	temperature	Kelvin (K)

Exercise

1.0/1 point (graded)
Find the units of the ideal gas constant R so that the equation makes physical sense.

(Type for Newtons, type for meters, type for moles, type for Kelvins.)

✓ **Answer:** N*m/(mol*K)

? INPUT HELP

Solution:

The units of both sides of the equation must be equal for this to make sense.

$$\frac{P}{\text{N/m}^2} V = n \frac{RT}{\text{mol K}}$$

Therefore the units of R must be $\frac{\text{N} \cdot \text{m}}{\text{mol} \cdot \text{K}}$ for the units of both sides of this equation to balance.

Submit

You have used 1 of 3 attempts

i Answers are displayed within the problem

Units and partial derivatives

2.0/2 points (graded)
Suppose that a quantity $z = f(x, t)$ has units of temperature measured in Kelvins, the variable x has units of length measured in meters, and the variable t has units of time measured in seconds.

What are the units of the partial derivative f_x ?

(Type for Newtons, type for meters, type for moles, type for Kelvins, type for seconds.)

✓ **Answer:** K/m

What are the units of the partial derivative f_t ?

(Type for Newtons, type for meters, type for moles, type for Kelvins, type for seconds.)

✓ **Answer:** K/s

? INPUT HELP

Solution:


Note that the units of the partial derivative f_x are the same as the units of its approxi

Note that the units of the partial derivative f_x are the same as the units of its approximation $\frac{\Delta f}{\Delta x}$, which is Kelvin per meter.

Similarly, the units of the partial derivative f_t are Kelvin per second.

Submit

You have used 1 of 3 attempts



 Answers are displayed within the problem

10. Variables and units

Hide Discussion

Topic: Unit 1: Functions of two variables / 10. Variables and units

Add a Post

Show all posts	▼	by recent activity	▼
	<u>Secret Constants</u>	8	▼
<u>I would like to politely say I am having some difficulty with the propositions advanced in this section. One of the early examples of a...</u>			
	<u>What is the function f that we have to find fx and fy derivatives?</u>	4	▼
<u>What is the function f that we have to find fx and fy derivatives?</u>			

< Previous

Next >



edX

- About
- Affiliates
- edX for Business
- Open edX
- Careers
- News

Legal

 Calculator

 Hide Notes

- [Terms of Service & Honor Code](#)
- [Privacy Policy](#)
- [Accessibility Policy](#)
- [Trademark Policy](#)
- [Sitemap](#)

Connect

- [Blog](#)
- [Contact Us](#)
- [Help Center](#)
- [Media Kit](#)
- [Donate](#)



© 2021 edX Inc. All rights reserved.
深圳市恒宇博科技有限公司 [粤ICP备17044299号-2](#)