





< Previous









Next >

12.3.1 Eigenvalues and Eigenvectors of $n \times n$ Matrices: Special Cases

 Bookmark this page

Week 12 due Dec 29, 2023 10:42 IST Completed

12.3.1 Eigenvalues and Eigenvectors of n x n Matrices: Special Cases

Video 12.3.1 Part 1

Start of transcript. Skip to the end.

0:00 / 0:00

▶ 2.0x

🔊

🔍

CC

🗣️

Dr. Robert van de Geijn: So now that we've examined 2 by 2 matrices, and 3 by 3 matrices, and how to find the eigenvalues and eigenvectors of those, let's move on to the general case where we have n by n matrices. We've seen the general case a little bit

Video

📄 [Download video file](#)

Transcripts

📄 [Download SubRip \(.srt\) file](#)

📄 [Download Text \(.txt\) file](#)

Reading Assignment

0 points possible (ungraded)
Read Unit 12.3.1 of the notes. [\[LINK\]](#)

☒ Done



Submit

✔ Correct

Discussion

Topic: Week 12 / 12.3.1

Hide Discussion

Add a Post

Show all posts ▼

by recent activity ▼

There are no posts in this topic yet.

✕

Calculator

Homework 12.3.1.1

1/1 point (graded)

Let $A \in \mathbb{R}^{n \times n}$ be a diagonal matrix: $A = \begin{pmatrix} \alpha_{0,0} & 0 & 0 & \cdots & 0 \\ 0 & \alpha_{1,1} & 0 & \cdots & 0 \\ 0 & 0 & \alpha_{2,2} & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & \alpha_{n-1,n-1} \end{pmatrix}.$

Then e_i is an eigenvector associated with eigenvalue $\alpha_{i,i}$.

TRUE

✓ Answer: TRUE

Just multiply it out. Without loss of generality (which means: take as a typical case), let $i = 1$. Then

$$\begin{pmatrix} \alpha_{0,0} & 0 & 0 & \cdots & 0 \\ 0 & \alpha_{1,1} & 0 & \cdots & 0 \\ 0 & 0 & \alpha_{2,2} & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & \alpha_{n-1,n-1} \end{pmatrix} \begin{pmatrix} 0 \\ 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix} = \alpha_{1,1} \begin{pmatrix} 0 \\ 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix}$$

Here is another way of showing this, leveraging our notation: Partition

$$A = \left(\begin{array}{c|c|c} A_{00} & 0 & 0 \\ \hline 0 & \alpha_{11} & 0 \\ \hline 0 & 0 & A_{22} \end{array} \right) \quad \text{and} \quad e_j = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix},$$

where α_{11} denotes diagonal element $\alpha_{j,j}$. Then

$$\left(\begin{array}{c|c|c} A_{00} & 0 & 0 \\ \hline 0 & \alpha_{11} & 0 \\ \hline 0 & 0 & A_{22} \end{array} \right) \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 0 \\ \alpha_{11} \\ 0 \end{pmatrix} = \alpha_{11} \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}$$

Submit


i Answers are displayed within the problem

Video 12.3.1 Part 2

[Start of transcript. Skip to the end.](#)



Dr. Robert van de Geijn: So here

 Calculator

▶ 0:00 / 0:00

▶ 2.0x

🔊

🔍

📄

🗣️

answer.

You partition the matrix where you identify

the i-th entry on the diagonal.

And we'll call that alpha 1 1.

And if you then take the unit basis vector

Video

📄 Download video file

Transcripts

📄 Download SubRip (.srt) file

📄 Download Text (.txt) file

Homework 12.3.1.2

1/1 point (graded)

Let $A = \left(\begin{array}{c|c|c} A_{00} & a_{01} & A_{02} \\ \hline 0 & \alpha_{11} & a_{12}^T \\ \hline 0 & 0 & A_{22} \end{array} \right)$, where A_{00} is square. Then α_{11} is an eigenvalue of A and $\begin{pmatrix} -(A_{00} - \alpha_{11}I)^{-1}a_{01} \\ 1 \\ 0 \end{pmatrix}$ is a corresponding eigenvector (provided $A_{00} - \alpha_{11}I$ is nonsingular).

TRUE ▼

✔ Answer: TRUE

What we are going to show is that $(A - \alpha_{11}I)x = 0$ for the given vector.

$$\begin{pmatrix} (A_{00} - \alpha_{11}I) & a_{01} & A_{02} \\ \hline 0 & 0 & a_{12}^T \\ \hline 0 & 0 & (A_{22} - \alpha_{11}I) \end{pmatrix} \begin{pmatrix} -(A_{00} - \alpha_{11}I)^{-1}a_{01} \\ 1 \\ 0 \end{pmatrix} = \begin{pmatrix} (A_{00} - \alpha_{11}I)[-(A_{00} - \alpha_{11}I)^{-1}a_{01}] + a_{01} + 0 \\ 0 + 0 + 0 \\ 0 \end{pmatrix} = \begin{pmatrix} -a_{01} + a_{01} + 0 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}$$

A more constructive way of verifying the result is to notice that clearly

$$\left(\begin{array}{c|c|c} (A_{00} - \alpha_{11}I) & a_{01} & A_{02} \\ \hline 0 & 0 & a_{12}^T \end{array} \right)$$

🧮 Calculator


$$\left(\begin{array}{c|c|c} & & \tilde{12} \\ \hline & 0 & \\ \hline & 0 & (A_{22} - \alpha_{11}I) \\ \hline \end{array}\right)$$

is singular since if one did Gaussian elimination with it, a zero pivot would be encountered exactly where the **0** in the middle appears. Now, consider a vector of form $\begin{pmatrix} \frac{x_0}{1} \\ 0 \end{pmatrix}$. Then

$$\left(\begin{array}{c|c|c} (A_{00} - \alpha_{11}I) & a_{01} & A_{02} \\ \hline 0 & 0 & a_{12}^T \\ \hline 0 & 0 & (A_{22} - \alpha_{11}I) \\ \hline \end{array}\right)\begin{pmatrix} \frac{x_0}{1} \\ 0 \end{pmatrix} = \begin{pmatrix} \frac{0}{0} \\ 0 \\ 0 \end{pmatrix}$$

means that $(A_{00} - \alpha_{11}I)x_0 + a_{01} = 0$. (First component on both sides of the equation.) Solve this to find that $x_0 = -(A_{00} - \alpha_{11}I)^{-1}a_{01}$.

Submit

 Answers are displayed within the problem

Video 12.3.1 Part 3

[Start of transcript. Skip to the end.](#)



 0:00 / 0:00

 **2.0x**











Video
 [Download video file](#)

Transcripts
 [Download SubRip \(.srt\) file](#)
 [Download Text \(.txt\) file](#)

Homework 12.3.1.3

1/1 point (graded)
The eigenvalues of a triangular matrix can be found on its diagonal.

TRUE 



< Previous

Next >



edX

- [About](#)
- [Affiliates](#)
- [edX for Business](#)
- [Open edX](#)
- [Careers](#)
- [News](#)

Legal

- [Terms of Service & Honor Code](#)
- [Privacy Policy](#)
- [Accessibility Policy](#)
- [Trademark Policy](#)
- [Sitemap](#)
- [Cookie Policy](#)
- [Your Privacy Choices](#)

Connect

- [Idea Hub](#)
- [Contact Us](#)
- [Help Center](#)
- [Security](#)
- [Media Kit](#)



© 2023 edX LLC. All rights reserved.
深圳市恒宇博科技有限公司 [粤ICP备17044299号-2](#)