

All posts sorted by recent activity

15.3.7 Confidence Intervals for Mean Estimation

Bookmark this page

- MO2.3
- MO2.13
- MO2.14
- MO2.15

The following video discusses confidence intervals. As shown in that video, for large N , we can prove that for **95%** of all possible samples, the population mean will be found in the *confidence interval*:

$$\bar{x} - 1.96 \frac{\sigma_x}{\sqrt{N}} < \mu_x < \bar{x} + 1.96 \frac{\sigma_x}{\sqrt{N}}$$

(15.22)

A higher confidence can be achieved, with a wider confidence interval. For example, **99%** of all possible samples will satisfy,


$$\bar{x} - 2.576 \frac{\sigma_x}{\sqrt{N}} < \mu_x < \bar{x} + 2.576 \frac{\sigma_x}{\sqrt{N}}$$

(15.23)

Note that in practice: σ_x is not known and so is estimated from the sample, i.e. $\sigma_x \approx s_x$.

The Python script used in this video (and several others) is available [here](#).

Video on confidence intervals for means



0:00 / 0:00

▶ 2.0x









[Start of transcript. Skip to the end.](#)

PROFESSOR: Now we're going to discuss the implications of the central limit theorem, or CLT.

And this is one of the classic results and plays a key role in probability

Video

 [Download video file](#)

Transcripts



edX

- [About](#)
- [Affiliates](#)
- [edX for Business](#)
- [Open edX](#)
- [Careers](#)
- [News](#)

Legal

- [Terms of Service & Honor Code](#)
- [Privacy Policy](#)
- [Accessibility Policy](#)
- [Trademark Policy](#)
- [Sitemap](#)
- [Cookie Policy](#)
- [Your Privacy Choices](#)

Connect

- [Idea Hub](#)
- [Contact Us](#)
- [Help Center](#)
- [Security](#)
- [Media Kit](#)



© 2023 edX LLC. All rights reserved.
深圳市恒宇博科技有限公司 [粤ICP备17044299号-2](#)