EdX and its Members use cookies and other tracking technologies for performance, analytics, and marketing purposes. By using this website, you accept this use. Learn more about these technologies in the Privacy Policy.





Homework 6 Maximum Likelihood

2. Recap: Maximum Likelihood

Course > Unit 3 Methods of Estimation > Estimation and Method of Moments > Estimators and Fisher information

Audit Access Expires Dec 24, 2019

You lose all access to this course, including your progress, on Dec 24, 2019.

Upgrade by Nov 4, 2019 to get unlimited access to the course as long as it exists on the site. **Upgrade now**

2. Recap: Maximum Likelihood Estimators and Fisher information **Instructions:**

For each of the following distributions, compute the maximum likelihood estimator based on n i.i.d. observations X_1, \ldots, X_n and the Fisher information, if defined. If it is not, enter **DNE** in each applicable input box.

(a)

3/3 points (graded)

$$X_i \sim \mathsf{Ber}\left(p
ight), \quad p \in (0,1)$$

(Enter **barX_n** for the sample average \overline{X}_n

$$\overline{X}_n = rac{1}{n} \sum_{i=1}^n X_i.$$

Hint: Use the definition of Fisher information that leads to the shorter computation.

(If the Fisher information is not defined, enter **DNE**.)

Fisher information
$$I\left(p\right)= \fbox{ 1/p/(1-p)}$$

Use Fisher Information to find the asymptotic variance $\,V\left(\hat{p}\,
ight)\,$ of the MLE $\,\hat{p}\,$.

$$V\left(\hat{p}
ight) = egin{bmatrix} eg$$

STANDARD NOTATION

Submit

You have used 1 of 3 attempts

(b)

3/3 points (graded)

$$X_i \sim \mathsf{Poiss}\left(\lambda
ight), \quad \lambda > 0,$$

which means that each X_i has distribution

$$\mathbf{P}_{\lambda}\left(X=k
ight)=e^{-\lambda}rac{\lambda^{k}}{k!},\quad k\in\mathbb{N}.$$

(Enter **barX_n** for the sample average \overline{X}_n .)

Maximum likelihood estimator $\hat{\lambda} = oldsymbol{f barX_n}$

(If the Fisher information is not defined, enter **DNE**.)

Fisher information $I(\lambda) = \boxed{ \begin{tabular}{c} 1/lambda \end{tabular}}$

Use Fisher Information to find the asymptotic variance $\,V\left(\hat{\lambda}
ight)\,$ of the MLE $\,\hat{\lambda}$.

$$V\left(\hat{\lambda}
ight) = egin{bmatrix} \mathsf{lambda} & oldsymbol{\checkmark} \ \lambda & oldsymbol{} \end{pmatrix}$$

STANDARD NOTATION

Submit

You have used 1 of 3 attempts

(c)

3/3 points (graded)

$$X_{i} \sim \mathsf{Exp}\left(\lambda
ight), \quad \lambda > 0,$$

which means that each X_1 has density

$$f_{\lambda}\left(x
ight) =\lambda e^{-\lambda x},\quad x>0.$$

(Enter **barX_n** for \overline{X}_n the sample average.)

Maximum likelihood estimator $\hat{\lambda} = \boxed{1/\text{barX_n}}$ \checkmark (If the Fisher information is not defined, enter **DNE**.)

Fisher information
$$I\left(\lambda\right)=$$
 1/lambda^2
$$\frac{1}{\lambda^2}$$

Use Fisher Information to find the asymptotic variance $\,V\left(\hat{\lambda}
ight)\,$ of the MLE $\,\hat{\lambda}$.

$$V\left(\hat{\lambda}
ight) = egin{bmatrix} \mathsf{lambda^2} \ \lambda^2 \end{bmatrix}$$

STANDARD NOTATION

Submit

You have used 1 of 3 attempts

(d)

7.0/7 points (graded)

$$X_{i}\sim\mathcal{N}\left(\mu,\sigma^{2}
ight),\quad\mu\in\mathbb{R},\,\sigma^{2}>0,$$

which means that each $\,X_1\,$ has density

$$f_{\mu,\sigma^2}\left(x
ight) = rac{1}{\sqrt{2\pi\sigma^2}} \mathrm{exp}\left(-rac{(x-\mu)^2}{2\sigma^2}
ight).$$

Hint: Keep in mind that we consider σ^2 as the parameter, not σ . You may want to write $au=\sigma^2$ in your computation.

(Enter ${\bf barX_n}$ for the sample average \overline{X}_n and ${\bf bar(X_n^2)}$ for the sample average of second moments $\overline{X_n^2}$.)

(Enter ${\bf barX_n}$ for the sample average \overline{X}_n and ${\bf bar(X_n^2)}$ for the sample average of second moments $\overline{X_n^2}$.)

Maximum likelihood estimator $\widehat{\sigma^2} = \boxed{ \mathsf{bar}(\mathsf{X}_n^2)-\mathsf{barX}_n^2}$

Hint: One of the formulas for Fisher information will lead to a much shorter computation.

(If the Fisher information is not defined, enter **DNE** for all boxes below.)

$$[I\left(\mu,\sigma^2\right)]_{1,1} = \begin{bmatrix} 1/\mathrm{sigma^2} & & & \\ & \frac{1}{\sigma^2} & & & \\ & &$$

Using the Fisher Information you obtain above, what is the asymptotic variance $V(\widehat{\sigma^2})$ of the MLE $\widehat{\sigma^2}$?Compare this with your result from Homework 5 Problem 3.

$$V\left(\widehat{\sigma^2}
ight) = oxed{2^* ext{sigma}^4}$$

STANDARD NOTATION

Submit

You have used 2 of 3 attempts

(e)

6/6 points (graded)

 X_i follows a shifted exponential distribution with parameters $\,a\in\mathbb{R}\,$ and $\,\lambda>0$. That means each $\,X_i\,$ has density

$$f_{a,\lambda}\left(x
ight)=\lambda e^{-\lambda\left(x-a
ight)}\mathbf{1}\{x\geq a\},\quad x\in\mathbb{R}.$$

(Enter ${f barX_n}$ for the sample average \overline{X}_n , and if applicable, use ${f min_i(X_i)}$ for ${\min_{1 < i < n} X_i}$).

Maximum likelihood estimator $\hat{a} = \boxed{\min_{i}(X_i)}$

Maximum likelihood estimator $\hat{\lambda} = \boxed{1/(\text{barX_n-min_i(X_i)})}$

Hint: Think of the effect of the indicator function on the derivatives. (If the Fisher information is not defined, enter **DNE** in all boxes below.)

$$[I\left(a,\lambda
ight)]_{1,1} = egin{bmatrix} ext{DNE} \ ext{DNE} \end{bmatrix}$$
 , $[I\left(a,\lambda
ight)]_{1,2} = egin{bmatrix} ext{DNE} \ ext{DNE} \end{bmatrix}$

2. Recap: Maximum Likelihood Estimators and Fisher information | Homework 6 Maximum Likelihood Estimation and Method of Moments | 18.6501x Courseware | edX $[I(a,\lambda)]_{2,1} =$ $ightharpoonup , \left[I\left(a,\lambda
ight)
ight]_{2,2} =$ DNE DNEDNESTANDARD NOTATION You have used 3 of 3 attempts Submit ✓ Correct (6/6 points) Discussion **Hide Discussion** Topic: Unit 3 Methods of Estimation: Homework 6 Maximum Likelihood Estimation and Method of Moments / 2. Recap: Maximum Likelihood Estimators and Fisher information Add a Post **≺** All Posts part e) Matrix Order question posted about 16 hours ago by **nbourbon** Is it ok to assume that the first value I_1_1 it corresponds to the second derivative of "a"? or of "lambda"?. I'm assuming that if I get one of the values "0" the grader expects "0" while "not defined" is a different thing right? This post is visible to everyone. **Erocha** (Community TA) about 14 hours ago - marked as answer about 13 hours ago by **nbourbon** Check slide 24. 0 is not the same as DNE. But pay attention to the hint.

| thanks I got the green tick now I was struggling with underst | tanding that indicator effect and it's key was not intuit | tive at the beginning but now I | |
|---|---|---------------------------------|--------------|
| posted about 13 hours ago by <u>nbourbon</u> | | | |
| Add a comment | | | |
| | | | |
| | | | |
| | | // | |
| Preview | | | |
| Submit | La aura Ala auta Vanifia d'Gantifi | | |
| | Learn About Verified Certificates | © All Righ | nts Reserved |