



< Previous



Next >

2. Planes, normal vectors, and tangent planes

 Bookmark this page

Part A due Oct 5, 2021 20:30 IST



Practice

5A-2

1/1 point (graded)
Let \mathbf{P} be the plane defined by $x + 2y + 3z = 17$. Find a vector which is normal to this plane. (It doesn't have to be a unit vector.)

[1,2,3]

✓ Answer: [1,2,3]

? INPUT HELP

Solution:

A normal vector can be taken as

$$\vec{n} = \nabla (x + 2y + 3z) = \langle 1, 2, 3 \rangle.$$

Submit

You have used 1 of 3 attempts

Answers are displayed within the problem

5A-3

1/1 point (graded)
Find the equation of a plane which is perpendicular to the vector $\langle 3, 2, 1 \rangle$ and passes through the point $(1, 1, 1)$.

(Express the plane in the format $0 = ax + by + cz + d$. We provide the $0 =$ for you, so you do not need to type that part.)

0 =

3*x+2*y+z-6

✓ Answer: 3*x + 2*y + z -6

Solution:

To find the equation of the plane we can use the fact that for any other point in the plane (x, y, z) the vector from $(1, 1, 1)$ to (x, y, z) , which is $\langle x - 1, y - 1, z - 1 \rangle$ is normal to the vector $\langle 3, 2, 1 \rangle$. That is the dot product between these two vectors is zero.

Therefore equation of the plane is

$$\langle 3, 2, 1 \rangle \cdot \langle x - 1, y - 1, z - 1 \rangle = 0$$
$$3(x - 1) + 2(y - 1) + (z - 1) = 0$$

(6.261)

(6.262)

, or equivalently, $3x + 2y + z - 6 = 0$.

Alternatively, we only need that a normal vector is $\langle 3, 2, 1 \rangle$, so the plane takes the form $3x + 2y + z + d = 0$. We can plug in the point $(1, 1, 1)$ as it lies on the plane.

This gives

$$3(1) + 2(1) + (1) + d = 0 \quad (6.263)$$

$$6 + d = 0 \quad (6.264)$$

$$d = -6 \quad (6.265)$$

Submit

You have used 1 of 3 attempts

i Answers are displayed within the problem

5A-4

2/2 points (graded)

Suppose that $f(x, y, z) = xz - yz^2 + xy$. Suppose that we start at the point $(2, 1, 1)$ and increase z slightly. Does f increase, decrease, or stay the same?

☐ f increases

☒ f decreases

☐ f stays the same



Is $f_z(2, 1, 1)$ positive, negative, or zero?

☐ positive

☐ negative

☒ zero



Solution:

A direct calculation yields

$$f(2, 1, 1+t) - f(2, 1, 1) = -t^2 < 0.$$

Hence **f decreases.**

Computing the partial derivative, you find that

$$f_z(x, y, z) = x - 2yz \quad (6.266)$$

$$f_z(2, 1, 1) = 2 - 2(1)(1) = 0 \quad (6.267)$$

Can you see why the function can both decrease as z increases, but have a zero partial derivative? (Hint, the function is not equal to its linear approximation.)

Submit

You have used 2 of 2 attempts

i Answers are displayed within the problem

Calculator

Hide Notes

5A-5

1/1 point (graded)

Let \mathcal{S} be the surface defined by $x^2 + 4y^2 + 4z^2 = 3$. The point $(1, 0.5, 0.5)$ lies in \mathcal{S} . Find a normal vector to the surface \mathcal{S} at the point $(1, 0.5, 0.5)$. (It does not have to be a unit vector.)

[2,4,4]

✔ Answer: [2,4,4]

? INPUT HELP

Solution:

A normal vector is

$$\nabla (x^2 + 4y^2 + 4z^2) (1, 0.5, 0.5) = \langle 2x, 8y, 8z \rangle|_{(1,0.5,0.5)} \tag{6.268}$$
$$= \langle 2 (1) , 8 (0.5) , 8 (0.5) \rangle \tag{6.269}$$
$$= \boxed{\langle 2, 4, 4 \rangle}. \tag{6.270}$$

Note that any constant multiple is also correct.

Submit

You have used 1 of 3 attempts

ⓘ Answers are displayed within the problem

2. Planes, normal vectors, and tangent planes

Hide Discussion

Topic: Unit 5: Curves and Surfaces / 2. Planes, normal vectors, and tangent planes

Add a Post

◀ All Posts

[STAFF] Which is the preferred answer?

discussion posted 2 days ago by [valleymd](#)

5A-4 has different answers depending upon whether the linear approximation approach is used with the gradient or if an actual calculation with a test value is used. Which is the desired approach?

This post is visible to everyone.

Add a Response

1 response

yves-M (Community TA)

2 days ago

+

★

⋮

The first one is better. The second one has a flaw: think of $f(x) = x^3$ and what happens at $x = 0$.

⋮

I used the first approach (the linear approximation with the gradient) as you suggested and was marked incorrect. My second approach (the direct calculation which was actually my first impulse) was the desired approach. Can this problem be reset so that I might get credit for realizing the difference in answers depending upon the approach used?

posted 2 days ago by [valleymd](#)

⋮

I have also used the method of linear approximation and the one based on the partial derivative. They both return the same answer .The direct calculation shows a different answer . It is problematic if one cannot have absolute confidence in these tools even if the conditions to apply them are met. The origin of the error is not at all o

Calculator

Hide Notes

https://learning.edx.org/course/course-v1:MITx+18.02.1x+2T2021/block-v1:MITx+18.02.1x+2T2021+type@sequential+block@ps5A-sequential/block-v1:MITx+18.02.1x+2T2021+type@vertical+block@ps5A-tab2

4/6

to spot.

posted a day ago by [bdpedx](#)



I think the confusion arises from the fact that we often ask the opposite question: Is the sign of f_z positive, negative, or zero. But truly, despite the fact that the z direction is tangent to the level curve here, it is in fact increasing.

I guess this is a problem with how this question is posed! I will reword, reframe, and reset attempts.

posted a day ago by [jfrench](#) (Staff)



A variable change of 0.1% results in a total change of 0.0001%. As far as I'm concerned, for estimation purposes, that's staying the same.

posted about 7 hours ago by [mmark](#)

Add a comment

Showing all responses

Add a response:

Preview

Submit



edX

- About
- Affiliates
- edX for Business
- Open edX
- Careers

Calculator

Hide Notes

[News](#)

Legal

[Terms of Service & Honor Code](#)

[Privacy Policy](#)

[Accessibility Policy](#)

[Trademark Policy](#)

[Sitemap](#)

Connect

[Blog](#)

[Contact Us](#)

[Help Center](#)

[Media Kit](#)

[Donate](#)



© 2021 edX Inc. All rights reserved.
深圳市恒宇博科技有限公司 [粤ICP备17044299号-2](#)