



MITx: 6.041x Introduction to Probability - The Science of Uncertainty



Bookmarks

- ▶ Unit 0: Overview
- ▶ Entrance Survey
- ▶ Unit 1: Probability models and axioms
- ▶ Unit 2: Conditioning and independence

▼ Unit 3: Counting

Lec. 4: Counting

Exercises 4 due Feb 24, 2016 at 23:59 UTC

Solved problems**Problem Set 3**

Problem Set 3 due Feb 24, 2016 at 23:59 UTC

Unit 3: Counting > Lec. 4: Counting > Lec 4 Counting vertical3



Bookmark

Exercise: Binomial probabilities

(2/2 points)

Recall that the probability of obtaining k Heads in n independent coin tosses is $\binom{n}{k} p^k (1-p)^{n-k}$, where p is the probability of Heads for any given coin toss.

Find the value of $\sum_{k=0}^n \binom{n}{k} p^k (1-p)^{n-k}$. (Your answer should be a number.)

1

**Answer: 1****Answer:**

Note that the events "0 Heads", "1 Heads", ..., " n Heads" are disjoint, and their union is the entire sample space. The summation is adding up the probability of all of these events. Hence, the sum must be 1. In other words, each term in the summation gives the probability of obtaining k Heads out of n tosses. We then sum over all values of k from 0 to n . Since the number of Heads must be one of $0, 1, \dots, n$ these probabilities must sum up to 1.

You have used 1 of 2 submissions

© All Rights Reserved



© edX Inc. All rights reserved except where noted. EdX, Open edX and the edX and Open EdX logos are registered trademarks or trademarks of edX Inc.

POWERED BY
OPENedX



