EdX and its Members use cookies and other tracking technologies for performance, analytics, and marketing purposes. By using this website, you accept this use. Learn more about these technologies in the Privacy Policy.





Homework 3: Introduction to

2. Concept Check: Hypothesis Test

Course > Unit 2 Foundation of Inference > Hypothesis Testing

> Using a Single Observation

Currently enrolled in **Audit Track** (expires December 25, 2019) <u>Upgrade (\$300)</u>

2. Concept Check: Hypothesis Test Using a Single Observation Setup:

Let X be a **single** (i.e. n=1) Gaussian random variable with unknown mean μ and variance 1. Consider the following hypotheses:

$$H_0: \mu=0 \quad ext{vs} \quad H_1: \mu
eq 0.$$

(a)

1/1 point (graded)

Define a test $\,\psi_lpha:\mathbb{R} o\{0,1\}\,$ with level $\,lpha\,$ that is of the form

$$\psi_{\alpha} = \mathbf{1}\{f_{\alpha}(X) > 0\},\,$$

for some function $\,f_{lpha}:\mathbb{R}
ightarrow\mathbb{R}\,.$

Generating Speech Output \downarrow ir test ψ above to satisfy the following:

- ullet symmetric in the value of X
- its "acceptance region" is an interval. (The **acceptance region** of a test is the region in which the null hypothesis is **not rejected**, i.e. the complement of its rejection region.)

Specify the function $f_{lpha}\left(X
ight)$ in terms of lpha below.

(Type **alpha** for α . If applicable, enter **abs(x)** for |x|, **Phi(x)** for $\Phi(x)=\mathbf{P}(Z\leq x)$ where $Z\sim\mathcal{N}(0,1)$, and **q(alpha)** for q_{α} , the $1-\alpha$ -quantile of a standard normal distribution, e.g. enter **q(0.01)** for $q_{0.01}$.)

$$f(X) =$$
abs(X)-q(alpha/2)

STANDARD NOTATION

Submit

You have used 2 of 3 attempts

✓ Correct (1/1 point)

(b)

3/3 points (graded)

Assume you observe $\,X=1.32$, and What is the value of your test ψ_{lpha} with level lpha=0.05?

What is the p-value of your test (keeping in mind the symmetry and interval requirements)? (If applicable, enter **abs(x)** for |x|, **Phi(x)** for $\Phi(x) = \mathbf{P}(Z \le x)$ where $Z \sim \mathcal{N}(0,1)$, and **q(alpha)** for q_{α} , the $1-\alpha$ -quantile of a standard normal distribution, e.g. enter **q(0.01)** for $q_{0.01}$.)

What is the conclusion of the test? \bigcirc Accept H_0 lacksquare Do not reject H_0 \bigcirc Accept H_1 \bigcirc Do not reject H_1 STANDARD NOTATION You have used 1 of 3 attempts Submit ✓ Correct (3/3 points) Discussion **Hide Discussion Topic:** Unit 2 Foundation of Inference:Homework 3: Introduction to Hypothesis Testing / 2. Concept Check: Hypothesis Test Using a Single Observation Add a Post Show all posts by recent activity ▼ Not able to figure out (a) 3 Lhave tried a bunch of functions that similar to |A| - |X|, which is symmetric around 0. But the grader keeps telling that the answer is incorrect. Can someone please help m... 22 --- edited ---**Generating Speech Output**

Audit Access Expires Dec 24, 2019

You lose all access to this course, including your progress, on Dec 24, 2019.

Upgrade by Nov 4, 2019 to get unlimited access to the course as long as it exists on the site. **Upgrade now**

Learn About Verified Certificates

© All Rights Reserved

Generating Speech Output