

edX and its Members use cookies and other tracking technologies for performance, analytics, and marketing purposes. By using this website, you accept this use. Learn more about these technologies in the [Privacy Policy](#).



Lecture 4: Parametric Estimation

[Course](#) > [Unit 2 Foundation of Inference](#) > [and Confidence Intervals](#)

> 5. Quadratic Risk of Estimators

Audit Access Expires Dec 24, 2019

You lose all access to this course, including your progress, on Dec 24, 2019.

Upgrade by Nov 4, 2019 to get unlimited access to the course as long as it exists on the site. **[Upgrade now](#)**

5. Quadratic Risk of Estimators

Quadratic Risk of Estimators

So here's what an exercise would look like.

Take blah to be an estimator.

Compute its bias.

Compute its variance.

Compute its quadratic risk.

Well, hopefully, if you compute the bias

Quadratic risk

- ▶ We want estimators to have low bias and low variance at the same time.



- ▶ The *Risk* (or *quadratic risk*) of an estimator $\hat{\theta}_n \in \mathbb{R}$ is

$$R(\hat{\theta}_n) = \mathbb{E} [|\hat{\theta}_n - \theta|^2]$$

$$\mathbb{E}[(\hat{\theta}_n - \mathbb{E}[\hat{\theta}_n] + \mathbb{E}[\hat{\theta}_n] - \theta)^2] = \underbrace{\mathbb{E}[(\hat{\theta}_n - \mathbb{E}[\hat{\theta}_n])^2]}_{\text{var}(\hat{\theta}_n)} + \underbrace{\mathbb{E}[(\mathbb{E}[\hat{\theta}_n] - \theta)^2]}_{\text{bias}(\hat{\theta}_n)^2} + 2 \underbrace{\mathbb{E}[(\hat{\theta}_n - \mathbb{E}[\hat{\theta}_n])(\mathbb{E}[\hat{\theta}_n] - \theta)]}_{0}$$

- ▶ Low quadratic risk means that both bias and variance are small:

$$\text{quadratic risk} = \text{VARIANCE} + \text{BIAS}^2$$

and you compute the variance, it's

going to cost you much to compute the quadratic risk.

OK.

Are there any questions?



Video

[Download video file](#)

Transcripts

[Download SubRip \(.srt\) file](#)

[Download Text \(.txt\) file](#)

Find the Quadratic Risk

1/1 point (graded)

Let $\hat{\theta}_n$ denote an estimator for a true parameter θ . The **quadratic risk** of $\hat{\theta}_n$ is defined to be

$$\mathbb{E}[(\hat{\theta}_n - \theta)^2].$$

As in the previous problem on variance, let $X_1, \dots, X_n \stackrel{iid}{\sim} \mathcal{U}([a, a+1])$ where a is an unknown parameter. What is the quadratic risk of the estimator $\bar{X}_n - \frac{1}{2}$?

Quadratic risk :

✓ Answer: 1/(12*n)


Solution:

Recall that

$$\text{quadratic risk} = \text{variance} + \text{bias}^2.$$

We showed in a previous question that this estimator is unbiased. Also note that $\text{Var}(\bar{X}_n) = \text{Var}(\bar{X}_n - \frac{1}{2}) = \frac{1}{12n}$. Hence, the quadratic risk is also $\frac{1}{12n}$.

You have used 1 of 3 attempts

 Answers are displayed within the problem

Properties of Estimators

1/1 point (graded)

Let $\hat{\theta}_n$ denote an estimator for a true parameter θ . Here n denotes the sample size. Which of the following properties of $\hat{\theta}_n$ would ensure that $\hat{\theta}_n$ converges in probability to θ as $n \rightarrow \infty$? (Choose all that apply.)

☒ $\hat{\theta}_n$ is consistent.

☐ $\hat{\theta}_n$ is unbiased.

☒ The quadratic risk of $\hat{\theta}_n$ goes to 0 as $n \rightarrow \infty$.

☐ The variance of $\hat{\theta}_n$ goes to 0 as $n \rightarrow \infty$.



Solution:

The first choice is correct, because by definition, consistency implies that the estimator $\hat{\theta}_n \rightarrow \theta$ as $n \rightarrow \infty$. The third choice, "The quadratic risk of $\hat{\theta}_n$ goes to 0 as $n \rightarrow \infty$.", is correct because if the quadratic risk $\mathbb{E}[(\hat{\theta}_n - \theta)^2] \rightarrow 0$ then $\hat{\theta}_n \rightarrow \theta$ in L^2 . By the properties of convergence, this implies that $\hat{\theta}_n \rightarrow \theta$ in probability.

Recall: Refer to Chapter 1 to review the relationship between the different types of convergence.

The second choice, " $\hat{\theta}_n$ is unbiased.", is incorrect. We give an example that shows that this choice is incorrect. Note that if $X_1, \dots, X_n \stackrel{iid}{\sim} N(\mu, 1)$, then $\hat{\theta}_n := X_1$ is an unbiased estimator for μ because $\mathbb{E}[X_1] = \mu$. However, it is not consistent: $X_1 - \mu$ does not tend to 0 as $n \rightarrow \infty$.

Using this same example, we can also see that the fourth choice "The variance of $\hat{\theta}_n$ goes to 0 as $n \rightarrow \infty$." is incorrect. The estimator $\hat{\theta}_n := 0$ (as an estimator for μ) has variance 0 for all n . If $\mu \neq 0$, then $\hat{\theta}_n - \mu = -\mu$, which is constant for all n and does not converge to 0.

Submit

You have used 2 of 3 attempts

i Answers are displayed within the problem

Discussion


Hide Discussion

Topic: Unit 2 Foundation of Inference: Lecture 4: Parametric Estimation and Confidence Intervals
/ 5. Quadratic Risk of Estimators


Add a Post

Show all posts ▼

by recent activity ▼

 [Properties of Estimation and Chebyshev's Inequality.](#)

4

 [Understanding why the Quadratic Risk/Mean Squared Error Equals Variance\(estimator\) + \(Bias\(estimator\)\)^2](#)

1

☒ Quadratic risk and Mean Squared Error

3

Are they the same thing?☐ What's the use of the quadratic risk when its expression involves the unknown?

2

While the expression in the example is a happy one, in which I know the variance does not depend on the parameter... the example may be differe...☒ Properties of Estimators - problem #3

2

[Learn About Verified Certificates](#)

© All Rights Reserved