



Newton's method with Hessian Matrix

Asked 3 years, 5 months ago Modified 3 years, 5 months ago Viewed 546 times



Let's say we want to use Newton's method to find optimal x_* for $f : \mathbb{R}^n \rightarrow \mathbb{R}$, with $x_{t+1} = x_t + H_f(x_t)^{-1} \nabla f(x_t)$

2



Can $H_f(x_t)^{-1}$ be calculated only once and used in every iteration step, or do we need to calculate it every time?



Does equation $H_f(x_0)(x_* - x_0) = \nabla f(x_0)$ need to be calculated only once to receive optimal value?



optimization

hessian-matrix

Share Cite Edit Follow Flag

asked Feb 16, 2020 at 21:24



Corgam

99 1 8

1 Answer

Sorted by:

Highest score (default)



"Can $H_f(x_t)^{-1}$ be calculated only once and used in every iteration step, or do we need to calculate it every time?"

1



No, the equation

$$H_f(x_t)(x_{t+1} - x_t) = -\nabla f(x_t)$$



needs to be solved at every iteration. This is the reason every iteration in Newton's method is costly compared to other methods with slower convergence. The exception is the quadratic problem



$$\min f(x), \quad f(x) = \frac{1}{2}x^T A x + b^T x$$

with $\nabla f(x) = Ax + b$ and $H_f(x) = A$. This problem is exactly solved by one step of Newton's.

$$\begin{aligned} A(x_1 - x_0) &= -Ax_0 - b \\ x_1 &= -A^{-1}b \end{aligned}$$

for which we see that $\nabla f(x_1) = 0$. Provided that A is positive semidefinite this is a solution to $\min f(x)$.

Share Cite Edit Follow Flag

answered Feb 18, 2020 at 10:38



Mikal

1,381

9

14