

◀ Previous				Next ▶
----------------------------	---	---	---	------------------------

2. Rotations

 Bookmark this page

Recitation due Sep 15, 2021 20:30 IST



Practice

Rotate a Vector

1/1 point (graded)

Recall:The rotation matrix $R_\theta = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$.

Find a two-dimensional vector \boldsymbol{v} such that the angle between \boldsymbol{v} and the vector $\begin{pmatrix} 1 \\ 2 \end{pmatrix}$ is 30° . There is more than one possible answer.

(Enter a vector using notation such as `[a,b]`.)

$\boldsymbol{v} =$ ✓ **Answer:** [sqrt(3)/2 - 1, sqrt(3) + 1/2]

? INPUT HELP

Solution:

We have $\boldsymbol{v} = R_{\frac{\pi}{6}} \begin{pmatrix} 1 \\ 2 \end{pmatrix} = \begin{pmatrix} \sqrt{3}/2 - 1 \\ \sqrt{3} + 1/2 \end{pmatrix}$.

It would also be correct to answer $R_{-\frac{\pi}{6}} \boldsymbol{v} = \begin{pmatrix} \sqrt{3}/2 + 1 \\ \sqrt{3} - 1/2 \end{pmatrix}$, or any scalar multiple of these.

Submit

You have used 1 of 5 attempts

ⓘ Answers are displayed within the problem

Rotate twice

1/1 point (graded)

Let $M_1 = R_{\frac{\pi}{3}}$ and $M_2 = R_{\frac{\pi}{6}}$. Find the matrix product $M_1 M_2$.

(Enter a matrix using notation such as `[[a,b],[c,d]]`.)

$M_1 M_2 =$ ✓ **Answer:** [[0,-1],[1,0]]


? INPUT HELP

Solution:

$M_1 M_2 = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$. This is $R_{\frac{\pi}{2}}$.

Submit

You have used 1 of 5 attempts

 Answers are displayed within the problem

Rotating Complementary Angles

1/1 point (graded)
Suppose θ and ϕ are complementary angles, that is, two angles such that $\theta + \phi = \frac{\pi}{2}$. Compute the product $R_\theta R_\phi$. Your answer should not involve θ or ϕ .

(Enter a matrix using notation such as `[[a,b],[c,d]]` .)

$R_\theta R_\phi =$  **Answer:** `[[0,-1],[1,0]]`


 INPUT HELP

Solution:

The product $R_\theta R_\phi = R_{\theta+\phi} = R_{\frac{\pi}{2}} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$.

Submit

You have used 1 of 5 attempts

 Answers are displayed within the problem

Power of rotation

1/1 point (graded)
Let $M = R_{\frac{\pi}{4}}$. What is the smallest value of $k > 0$ such that $M^k = I$?


$k =$  **Answer:** 8

Solution:

The product $R_\theta R_\phi = R_{\theta+\phi}$. It takes 8 rotations by $\pi/4$ to come back to the start. Therefore $k = 8$, or any integer multiple of 8.

Submit

You have used 1 of 5 attempts


 Answers are displayed within the problem


2. Rotations

Hide Discussion


Topic: Unit 4: Matrices and Linearization / 2. Rotations

Add a Post

Show all posts 

by recent activity 

There are no posts in this topic yet.



[< Previous](#)

[Next >](#)

© All Rights Reserved



edX

- [About](#)
- [Affiliates](#)
- [edX for Business](#)
- [Open edX](#)
- [Careers](#)
- [News](#)

Legal

- [Terms of Service & Honor Code](#)
- [Privacy Policy](#)
- [Accessibility Policy](#)
- [Trademark Policy](#)
- [Sitemap](#)

Connect

- [Blog](#)
- [Contact Us](#)
- [Help Center](#)
- [Media Kit](#)
- [Donate](#)



© 2021 edX Inc. All rights reserved.
深圳市恒宇博科技有限公司 [粤ICP备17044299号-2](#)

Calculator

Hide Notes