



All posts sorted by recent activity

15.1.2 A New Model Problem: Projectile Motion with Uncertainty

Bookmark this page

MO2.13

As we explore the fundamentals of probability and statistics, we will use a simpler example than the Martian lander to reduce the computational expense required to demonstrate these concepts. Specifically, as shown in Figure 15.1, consider a projectile launched from the ground with an initial speed V_0 at an angle θ_0 . g is the gravitational acceleration. The projectile impacts the ground at the location x_f .




Figure 15.1: Projectile motion problem definition. When only gravitational force is included (i.e. neglected aerodynamic drag), then the impact location can be found to be:

$$x_f = \frac{V_0^2}{g} \sin 2\theta_0$$

(15.1)

We will consider the nominal case for the projectile motion to be $V_{0\text{nom}} = 30 \text{ m/s}$ and $\theta_{0\text{nom}} = 30^\circ$. Assuming gravity is $g = 9.81 \text{ m/s}^2$, then the impact location for these nominal conditions is $x_{f\text{nom}} = 79.45 \text{ m}$.



edX

- [About](#)
- [Affiliates](#)
- [edX for Business](#)
- [Open edX](#)
- [Careers](#)
- [News](#)

Legal

- [Terms of Service & Honor Code](#)
- [Privacy Policy](#)
- [Accessibility Policy](#)
- [Trademark Policy](#)
- [Sitemap](#)
- [Cookie Policy](#)
- [Your Privacy Choices](#)

Connect

- [Idea Hub](#)
- [Contact Us](#)
- [Help Center](#)
- [Security](#)
- [Media Kit](#)



© 2023 edX LLC. All rights reserved.
深圳市恒宇博科技有限公司 [粤ICP备17044299号-2](#)