

### BerkeleyX: CS190.1x Scalable Machine Learning

# PRINCIPAL COMPONENT VECTORS (1/1 point)

Principal component vectors have length

on the length of the dataset
o sqrt(n)
● d the number of features ✔
k the number of principal components selected

#### **EXPLANATION**

Principal component vectors have length d. The dot product between a principal component and an observation's features creates a score or transformed feature, so the number of features and the length of the principal component vector must match.

CHECK HIDE ANSWER

## **EIGENVALUES** (1/1 point)

The top k principal components correspond to the top k eigenvalues.



### **EXPLANATION**

Eigenvalues are proportional to the variance explained by the corresponding principal component. The top principal component has the largest eigenvalue, the second principal component has the next largest eigenvalue, etc.

CHECK

**HIDE ANSWER** 

# COMPUTATION AND STORAGE COMPLEXITY (1/1 point)

The distributed PCA algorithm we implemented in the lab used:

✓ O(d^2) local storage
✓ O(d^3) local computation
✓ O(d) local storage
O(d^2) local computation

Note: Make sure you select all of the correct options—there may be more than one!

### **EXPLANATION**

The outer product results in a d by d matrix which requires  $O(d^2)$  local storage. Eigendecomposition on the aggegated results of the outer products requires  $O(d^3)$  local computation.

CHECK

**HIDE ANSWER** 

# PLOTTING CORRELATED VARIABLES (1/1 point)

In Visualization 1, what would the data look like if covariance equaled -1.0?

A vertical line
Similar to covariance of 0
Similar to covariance of .9
A diagonal line

### **EXPLANATION**

A covariance of -1.0 would produce an exact relationship between x1 and x2, which would show up as a diagonal line of points from the top-left to the bottom-right of the graph.

CHECK HI

HIDE ANSWER

# HIGH POSITIVE COVARIANCE (1/1 point)

In Visualization 1 when the covariance is .9, where are most of the points on the plot?

- lower-left and lower-right
- upper-right and lower-left
- upper-left and lower-right
- upper-left and upper-right

### **EXPLANATION**

A positive covariance creates the tendency for the points to fall around a diagonal line with a positive slope, leading to most points falling in the lower-left and upper-right of the graph.

CHECK

**HIDE ANSWER** 

## PCA FUNCTION (1/1 point)

When running the pca function what is the largest k we should use?

- 0 1
- n -- length of dataset
- d -- number of features

#### **EXPLANATION**

We can generate up to d orthonormal eigenvectors from the covariance matrix, so k should be less than or equal to d.

CHECK

**HIDE ANSWER** 

## TIME BASED AGGREGATION (1/1 point)

In Visualization 9, does the resulting spatial map appear symmetric or asymmetric about the midline (horizontal line across the middle of the brain)?



#### **EXPLANATION**

The time aggregated information shows a symmetric response about the midline.

CHECK

**HIDE ANSWER** 

## DIRECTION BASED AGGREGATION (1/1 point)

In Visualization 10, does the resulting spatial map appear symmetric or asymmetric about the midline (horizontal line across the middle of the brain)?

<ul><li>Symmetric</li></ul>	
Asymmetric	✓

### **EXPLANATION**

The direction aggregated information shows an asymmetric response about the midline.

CHECK

**HIDE ANSWER** 

# SURVEY: LAB5 COMPLETION TIME (1/1 point)

How long did Lab FIVE take you to complete (in hours - decimals are OK)?





3

Please click "Check" to save your answers.

CHECK

**SHOW ANSWER** 

⊚ 🖲 🕲 🗑 Some Rights Reserved



About Blog News FAQs Contact Jobs Donate Sitemap

Terms of Service & Honor Code Privacy Policy Accessibility Policy

© edX Inc. All rights reserved except where noted. EdX, Open edX and the edX and Open EdX logos are registered trademarks or trademarks of edX Inc.

















