



< Previous



✓



✓



✓



✓



✓



✓



✓



✓



✓



✓



✓

















Next >

Notifications

×

11. Complex eigenvalues and eigenvectors

 Bookmark this page

Upgrade your course today

- ✓

You are taking the free version of this course.
- ✓

Upgrade today and, upon passing, receive your digital certificate signed by MIT faculty to highlight the knowledge and skills you’ve gained from this MITx course.

Upgrade

Recall that complex non-real eigenvalues come in pairs of complex conjugates. It turns out the eigenvectors of a complex eigenvalue are the complex conjugates of the eigenvectors of the conjugate eigenvalue.

Example 11.1 Find the eigenvalues and eigenvectors of the matrix $\mathbf{A} = \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$.

We have found previously that the eigenvalues are $1, \pm i$. Let us now find the eigenvectors. We will start with those of the complex eigenvalue i .

Eigenspace of i : $\text{NS}(i\mathbf{I} - \mathbf{A})$.

We reduce $i\mathbf{I} - \mathbf{A}$ to row-echelon form;

$$i\mathbf{I} - \mathbf{A} = \begin{pmatrix} i & 1 & 0 \\ -1 & i & 0 \\ 0 & 0 & i-1 \end{pmatrix} \rightarrow \begin{pmatrix} i & 1 & 0 \\ 0 & 0 & -1 \\ 0 & 0 & 0 \end{pmatrix}. \quad (3.14)$$

By back substitution (the first equation is $ix + y = 0$), the solutions are $\mathbf{c} \begin{pmatrix} i \\ 1 \\ 0 \end{pmatrix}$ for any scalar $\mathbf{c}$. That is,

$$\text{Eigenspace of } i = \text{NS}(i\mathbf{I} - \mathbf{A}) = \text{Span} \begin{pmatrix} i \\ 1 \\ 0 \end{pmatrix}. \quad (3.15)$$

Eigenspace of $-i$: $\text{NS}(-i\mathbf{I} - \mathbf{A})$.

Since $\mathbf{A}$ is a real matrix, $-i\mathbf{I} - \mathbf{A}$ is the complex conjugate of $i\mathbf{I} - \mathbf{A}$, and every step in the Gaussian elimination is the same as before except with i replaced by $-i$:

$$-i\mathbf{I} - \mathbf{A} = \begin{pmatrix} -i & 1 & 0 \\ -1 & -i & 0 \\ 0 & 0 & -i-1 \end{pmatrix} \rightarrow \begin{pmatrix} i & -1 & 0 \\ 0 & 0 & -1 \\ 0 & 0 & 0 \end{pmatrix}. \quad (3.16)$$

Therefore, the row-echelon form of $-i\mathbf{I} - \mathbf{A}$ is the complex conjugate of the row echelon form of $i\mathbf{I} - \mathbf{A}$ and the vectors in the nullspace of $-i\mathbf{I} - \mathbf{A}$ are the complex conjugates of the vectors in the nullspace of $i\mathbf{I} - \mathbf{A}$. Hence,

$$\text{Eigenspace of } -i = \text{NS}(-i\mathbf{I} - \mathbf{A}) = \text{Span} \begin{pmatrix} -i \\ 1 \\ 0 \end{pmatrix}. \quad (3.17)$$

Eigenspace of 1 : $\text{NS}(\mathbf{I} - \mathbf{A})$.

We start by reducing $\mathbf{I} - \mathbf{A}$:

$$\mathbf{I} - \mathbf{A} = \begin{pmatrix} 1 & 1 & 0 \\ -1 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}. \quad (3.18)$$

This gives the eigenspace

$$\text{Eigenspace of } 1 = \text{NS}(\mathbf{I} - \mathbf{A}) = \text{Span} \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}. \quad (3.19)$$

(You could also have guessed the solution $\mathbf{v} = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$ and plugged it into $\mathbf{A}\mathbf{v}$ to check that indeed it is an eigenvector of eigenvalue 1.)

Conclusion : The eigenvalues and corresponding eigenspaces of the matrix $\begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$ are:

$$\begin{array}{cc} \text{Eigenvalue} & \text{Corresponding eigenspace} \end{array} \quad (3.20)$$

$$\lambda = i; \text{Span} \begin{pmatrix} i \\ 1 \\ 0 \end{pmatrix} \quad (3.21)$$

$$(-i)$$

$$\lambda = -i; \text{Span} \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

(3.22)

$$\lambda = 1; \text{Span} \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}$$

(3.23)

(The eigenvectors for each eigenvalue are all nonzero vectors in the corresponding eigenspace.)

11. Complex eigenvalues and eigenvectors

Hide Discussion

Topic: Unit 2: Linear Algebra, Part 2 / 11. Complex eigenvalues and eigenvectors

Add a Post

Show all posts

by recent activity

There are no posts in this topic yet.

< Previous

Next >