

< Previous



Next >

### Question 13

 Bookmark this page

Final Exam due Jan 21, 2025 04:09 IST   Completed

### Question 13

1.0/1.0 point (graded)

The formulation of the question below is incomplete. It is included to avoid any ambiguity. The complete question and proposed solutions are available in the PDF file.

Suppose that we have three functions  $f(i)$ ,  $g(i)$  and  $h(i)$ ,  $i \in A \subset \mathbb{N}$  ( $A$  is an arbitrary subset of  $\mathbb{N}$ ), such that for all  $i$  we have  $0 \leq f(i) \leq M_f$ ,  $0 \leq g(i) \leq M_g$  and  $0 \leq h(i) \leq M_h$ . We would like to model that for each  $i$ , at least one of the three functions must be equal to 0. This can be modeled as (select the correct answer):

☐ Solution 1.

☒ Solution 2.

☐ Solution 3.

☐ Solution 4.



Submit

< Previous

Next >



edX

[About](#)

[Affiliates](#)

[edX for Business](#)

[Open edX](#)

[Careers](#)

XPERT

 Show Notes

# Legal

[Terms of Service & Honor Code](#)

[Privacy Policy](#)

[Accessibility Policy](#)

[Trademark Policy](#)

[Sitemap](#)

[Cookie Policy](#)

[Your Privacy Choices](#)

---

# Connect

[Idea Hub](#)

[Contact Us](#)

[Help Center](#)

[Security](#)

[Media Kit](#)

---

