



| | | | |
|-------------|------------------------|-------------------|---------------------|
| Name: | SUSMITA PAUL CHOUDHURY | Registration No.: | 594484 |
| Age/Gender: | 28 Y/Female | Registered: | 29/Dec/2022 12:40PM |
| Patient ID: | 012212290009 | Analysed: | 29/Dec/2022 02:26PM |
| BarcodeNo: | 61745462 | Reported: | 29/Dec/2022 06:32PM |
| Refer By: | Dr. G.B.DAS (MD) | Panel: | SILIGURI RPC |

DEPARTMENT OF SEROLOGY

| Test Name | Result | Unit | Biological Ref.Interval |
|-----------|--------|------|-------------------------|
|-----------|--------|------|-------------------------|

BLOOD GROUP ABO AND RH TYPING*

BLOOD GROUP ABO AND RH FACTOR

| | |
|-------------|----------|
| Blood Group | O |
| Rh Factor | POSITIVE |

Sample Type: Whole Blood EDTA



Barnika Purkayastha

DR. BARNIKA PURKAYASTHA
MBBS, MD (Pathology)
Consultant Pathologist



| | | | |
|-------------|------------------------|-------------------|---------------------|
| Name: | SUSMITA PAUL CHOUDHURY | Registration No.: | 594484 |
| Age/Gender: | 28 Y/Female | Registered: | 29/Dec/2022 12:40PM |
| Patient ID: | 012212290009 | Analysed: | 29/Dec/2022 02:26PM |
| BarcodeNo: | 61745463 | Reported: | 29/Dec/2022 05:45PM |
| Refer By: | Dr. G.B.DAS (MD) | Panel: | SILIGURI RPC |

DEPARTMENT OF SEROLOGY

| Test Name | Result | Unit | Biological Ref.Interval |
|--|--------|------|--|
| HEPATITIS B SURFACE ANTIGEN; HBsAg ELISA Method | 0.168 | COI | <0.9 Non Reactive 0.9 to 1.0 Equivocal >1.0 Positive |

Note:

1. All Reactive results for further confirmation Molecular assays are recommended
2. Discrepant results may be observed during pregnancy, patients receiving mouse monoclonal antibodies for diagnosis or therapy & mutant forms of HBsAg
3. For diagnostic purposes, results should be used in conjunction with clinical history and other hepatitis markers for Acute or Chronic infection
4. For monitoring HBsAg levels, Quantitative HBsAg assay is recommended

Comment

Hepatitis B Virus (HBV) is a member of the Hepadna virus family causing infections of the liver with extremely variable clinical features. Hepatitis B is transmitted primarily by body fluids especially serum and also spread effectively sexually and from mother to baby. In most individuals HBV hepatitis is self limiting, but 1-2% normal adolescents and adults develop Chronic Hepatitis. Frequency of chronic HBV infection is 5-10% in immunocompromised patients and 80% in neonates. The initial serological marker of acute infection is HBsAg which typically appears 2-3 months after infection and disappears 12-20 weeks after onset of symptoms. Persistence of HBsAg for more than six months indicates development of carrier state or Chronic liver disease.

Uses

◆ Routine screening of blood and blood products to prevent transmission of Hepatitis B virus (HBV) to recipients ◆ To diagnose suspected HBV infection and monitor the status of infected individuals ◆ To evaluate the efficacy of antiviral drugs ◆ For Prenatal Screening of pregnant women

Sample Type: Serum

- (1) The Results relate only to the items tested.
- (2) The test report shall not be reproduced except in full, without approval of the authority.

*** End Of Report ***



Barnika Purkayastha

DR. BARNIKA PURKAYASTHA
MBBS, MD (Pathology)
Consultant Pathologist