Daughter-Universe Nucleation from a Metastable Dark-Energy Vacuum: Thermal and Quantum Decay, Past-Light-Cone Survival, and Reheating Bounds

Leon Sandler
Independent Researcher, The Conscious Machine — Independent Research, Northbrook, Illinois, USA
sandler.leon@gmail.com · ORCID 0009-0007-4584-808X

Abstract

A metastable dark-energy vacuum could nucleate bubbles that evolve into daughter regions. We ask whether such a region can be a hot expanding universe and whether cosmological cooling raises the transition probability. For a scalar potential with a metastable minimum at the dark-energy scale we compute the Coleman–De Luccia bounce (error budget, negative-mode structure, 24 executable checks); all scales follow from the dark energy, $g = f^2/M_{\rm Pl}^2 \sim 10^{-60}$, and the quantum decay is controlled by one action $B = b(\eta)/\lambda$, with $\Gamma/H^4 = 1$ at $B \simeq 287$. Survival is evaluated on the Lorentzian past light cone of the observed expansion history ($V_4 = 0.059\, H_\Lambda^{-4}$, 445 times below the Euclidean four-sphere; 95% bound $\Gamma/H_\Lambda^4 < 51$). A thermal mass rescales the potential exactly, with $\eta_{\rm eff} = \eta/(1-2\theta)^{3/2}$; the O(3) thermal saddle dominates above 0.18 meV and the rate falls monotonically as the universe cools, so cooling cannot raise the late-time probability: a decoupled scalar is unaffected, and one in equilibrium with the CMB raises the excluded prior mass from 15% to 24–49% and closes the late-time decay window (priors are samples, not probabilities). Lorentzian simulations of the daughter region with two fields, dissipation and gravity show that it reheats to at most 2.2 meV and stays curvature dominated; a reservoir hidden by a negative offset leaves an anti-de Sitter vacuum that crunches after 5.5 s when it reaches 0.7 MeV; flattening at the dark-energy scale needs a vev above $4M_{\rm Pl}$ and still reheats below 1.5 meV; and repeated transitions obey $\sum \Delta V \le \rho_\Lambda$. A hot epoch at 1 s would need a parent energy scale at least 41 orders of magnitude above the Planck scale, and a matter trigger $\Delta \ln m \gtrsim 8.9 \times 10^{-4}$. The daughter region is cold and empty, and a cyclic reset by the present dark energy is not supported.

Keywords: *false vacuum decay; Coleman–De Luccia instanton; thermal activation; daughter universe; dark energy; reheating; cyclic cosmology; past light cone; falsifiability*

1 Introduction

In the standard cosmological model the late universe approaches de Sitter space: matter and radiation dilute, and a positive vacuum energy ρ_Λ remains. If that energy is not an exact cosmological constant but the energy of a metastable scalar vacuum, the vacuum can decay by quantum tunneling through a Coleman–De Luccia (CDL) instanton [1]. It has been suggested

that such a late-time transition could reset the cosmological history, so that the long-term evolution is cyclic rather than terminal; cyclic and bouncing proposals are discussed in [2, 3, 4, 5]. Two things must be kept apart. The formation of a bubble universe is not a cyclic cosmology: a cycle needs a mechanism that connects successive epochs. The question tested here is narrower and can be answered by calculation: can the tunneling of the present dark energy produce a new hot, expanding epoch? That requires a specified metastable minimum, the tunneling action including gravity, a rate compared with the observed expansion history, and the interior of the nucleated region followed to see whether it can become a hot big bang.

A further distinction concerns the outcome. In a branching or recycling picture (Option A) a parent metastable vacuum nucleates daughter regions by tunneling; each is a bubble inside the parent spacetime, and whether it is a hot expanding universe, or a universe at all in the sense of [6, 7, 8], is a separate question. In a genuinely cyclic cosmology (Option B) the same region expands, contracts, bounces and reheats, which needs a dynamical bounce beyond tunneling and raises problems of stability and entropy [9, 10]. This paper addresses Option A with the Coleman–De Luccia theory and treats Option B only by stating, in Section 5.4, what it would require.

Four claims are tested. (i) A scalar potential with a metastable minimum at the dark-energy scale can nucleate bubbles of lower-energy vacuum at a rate that is neither negligible nor already excluded. (ii) The nucleated region reheats into a hot expanding universe. (iii) The transition is triggered, or enhanced, by the dilution of matter in the late universe. (iv) Cosmological cooling, acting on a scalar coupled to the CMB, makes the transition more probable at late times. The analysis is conditional: it takes the dark-energy density as the false-vacuum energy and asks what follows. The result is a quantitative consistency and exclusion analysis, not a new cosmological mechanism.

The contributions are the following.

- A reproducible map from the parameters of an explicit scalar potential to cosmological decay times, with a numerical error budget, an independent boundary-value solution, the negative-mode structure of the bounce, the thin-wall, Parke and Hawking–Moss limits, and a treatment of the prefactor as a systematic uncertainty (Sections 2–4).

- A joint analysis of nucleation, survival and the post-transition energy budget, with the survival of the present vacuum evaluated on the Lorentzian past light cone of the observed expansion history (Section 4.2), and a statement of which energy sources the reheating bound excludes by geometry and which only by energy conservation (Section 5).

- Explicit conditions that separate a late-time vacuum transition from a viable cosmological restart, divided into internal consistency conditions and observational tests (Section 7), and a model-independent requirement on any matter coupling that could trigger the transition (Section 6).

- A finite-temperature extension: the exact rescaling of the thermally corrected potential, the O(3) and O(4) exponents and their crossover, the past-light-cone survival integral with

a time-dependent rate, and prior maps showing that cooling cannot raise the late-time transition probability (Sections 4.6–4.8).

- Lorentzian simulations of the daughter region with two fields, dissipation and gravity, a reservoir hidden by a negative offset, hilltop flattening, repeated transitions and entropy accounting, and the resulting criteria for a viable daughter universe (Section 5.5).

1.1 Relation to previous work

The ingredients are established, and the paper claims novelty only for the specific combination and the quantitative results for the dark-energy vacuum. Table 1 states for each ingredient what is known, what is used from the literature and what is computed here.

2 Scalar effective theory and the potential

We consider a real scalar field minimally coupled to Einstein gravity,

$$S = \int d^4x\sqrt{-g}\left[\frac{M_{\rm Pl}^2}{2}R - \frac{1}{2}g^{\mu\nu}\partial_\mu\varphi\,\partial_\nu\varphi - V(\varphi)\right] \tag{1}$$

where $M_{\rm Pl}$ is the reduced Planck mass. The false vacuum has the energy density $V_f = V(\varphi_f)$, which is identified with the observed dark energy, $V_f = \rho_\Lambda = 2.55\times10^{-47}$ GeV^4 (quarter-root 2.25 meV; Planck 2018 [37], $H_0 = 67.66$ km s^{-1} Mpc^{-1}, $\Omega_\Lambda = 0.6889$). The asymptotic de Sitter rate is $H_\Lambda = \sqrt{V_f/3}/M_{\rm Pl} = 1.20\times10^{-42}$ GeV, so that $H_\Lambda^{-1} = 17.4$ Gyr, and $M_{\rm Pl}^4/V_f = 1.38\times10^{120}$, with $\ln(M_{\rm Pl}^4/V_f) = 276.6$. A microscopic origin of the potential is not assumed.

A single dimensionless family covers the generic metastable case. With $x = \varphi/f$,

$$V(\varphi) = V_t + f^4\lambda\left[\frac{(x^2-1)^2}{8} + \frac{\eta\,x}{2} - p_{\rm min}(\eta)\right], \qquad x^3 - x + \eta = 0 \;\Rightarrow\; x_t < x_{\rm top} < x_f \tag{2}$$

where $p_{\rm min}$ is the value of the bracket at the true minimum, so that $V(\varphi_t) = V_t$. The tilt η lifts the degeneracy of the two wells; a false minimum exists only for $\eta < \eta_c = 2/(3\sqrt{3}) = 0.3849$. The energy difference and the barrier height above the false minimum are

$$\Delta V = V_f - V_t = f^4\lambda\,\delta(\eta), \qquad V_{\rm top} - V_f = f^4\lambda\,\beta(\eta), \qquad \delta \simeq \eta\ (\eta \ll 1) \tag{3}$$

The motivation for this family is minimality. A quartic with a small linear term is the simplest renormalizable potential with a metastable minimum and was the standard example in the original treatments of vacuum decay [11, 1]; it has three parameters with direct meaning, the quartic coupling λ, the tilt-to-barrier ratio η and the scale f, and the latter is fixed by the dark energy. The coupling is restricted to $10^{-2} \le \lambda \le 4\pi$, the upper limit being the perturbative-unitarity bound for a quartic interaction and the lower limit an arbitrary but wide cut that is varied in Section 4.5; the tilt is restricted to $0.03 \le \eta \le 0.38$, which excludes the nearly degenerate case, where $B \propto \eta^{-3}$ is so large that the transition is irrelevant, and ends just below

Table 1: Closest literature and what this paper adds.

Ingredient	Established (references)	This paper
Bounce, thin wall, gravity, Hawking–Moss	Coleman; Callan–Coleman; CDL; Hawking–Moss; Parke; Jensen–Steinhardt; Hackworth–Weinberg [11, 12, 1, 13, 14, 15, 16]	Used as validation limits. New: an explicit dark-energy-scale potential, $g \sim 10^{-60}$, and a numerical error budget and negative-mode check (Section 3).
Prefactor of the rate	Determinant methods [17, 18]	Not recomputed; treated as a systematic uncertainty with a stated spread and shown to enter B_c additively (Section 4.3).
Survival from the past light cone	Standard for the electroweak vacuum [19, 20]; percolation [21, 22]	Applied to the dark-energy vacuum with the observed expansion history: $V_4 = 0.059\, H_\Lambda^{-4}$, 445 times below the Euclidean four-sphere (Section 4.2).
Open interior and reheating	Open universes from bubbles need an inflationary stage [23, 24, 25]	Quantified for the dark-energy scale: curvature parameter, e-folds and the parent scale needed for a hot epoch (Section 5.3).
Finite-temperature decay	Thermal masses [26, 27]; thermal activation and the O(3) bounce [28, 29, 30]	Exact rescaling of the tilted quartic family under a thermal mass; O(3)/O(4) crossover for the dark-energy vacuum; survival with $\Gamma(T_\gamma(t))$ (Sections 4.6–4.8).
Daughter universes by tunneling	[6, 7, 8]	The nucleated region quantified as a cold, curvature-dominated daughter region, with the energy budget and reservoir bounds (Section 5).
Cyclic and bouncing models	[2, 3, 4, 5]	These supply the reset by other means; here one proposed mechanism, tunneling of the present dark energy, is tested.
Landscape and bubble collisions	[31, 32, 25, 33, 34]	Observable signals need $\Gamma/H^4 = O(1)$; the width of that window in B and its prior mass are computed (Section 4.5).
Matter couplings and screening	[35, 36]	A coupling-independent requirement, $\Delta \ln m$, and its translation to linear and quadratic couplings (Section 6).

the point where the minimum disappears. Other potentials change $b(\eta)$ and the interior energy fraction but not the scales that decide the results (Section 8). A scalar with a mass of the order of meV is technically unnatural without a protecting symmetry; this is a known property of every quintessence-like model and is not addressed here.

The choice $V_t = 0$ (decay to a Minkowski vacuum, $r_L \equiv V_f/\Delta V = 1$) is the baseline, because it maximizes the energy released to the interior for a given V_f when $V_t \geq 0$; $V_t < 0$ (anti-de Sitter) and $V_t > 0$ are considered in Sections 5.3 and 4.5. The scale f is not free: it is fixed by $V_f = \rho_\Lambda$, $f^4 = \rho_\Lambda/(r_L \lambda \delta)$. For $\lambda = 1$ and $\eta = 0.3$ one finds $f = 3.05$ meV and a field excursion of 5.8 meV, that is, the dark-energy scale. The gravitational parameter is then fixed too,

$$g \equiv \frac{f^2}{M_{\rm Pl}^2} = \sqrt{\frac{V_f/M_{\rm Pl}^4}{r_L\,\lambda\,\delta(\eta)}} \tag{4}$$

which takes the values 6.4×10^{-61} to 2.7×10^{-60} over the benchmark parameters of Table 4. Gravity is therefore a 10^{-60} effect in every equation below that does not contain the large number $M_{\rm Pl}^4/V_f$.

3 The Euclidean bounce, with and without gravity

3.1 Equations and action

The O(4)-symmetric Euclidean metric $ds^2 = d\xi^2 + \rho(\xi)^2 d\Omega_3^2$ gives the CDL equations [1]

$$\varphi'' + 3\frac{\rho'}{\rho}\varphi' = \frac{dV}{d\varphi}, \qquad \rho'^2 = 1 + \frac{\rho^2}{3M_{\rm Pl}^2}\left(\frac{1}{2}\varphi'^2 - V\right) \tag{5}$$

with $\varphi'(0) = 0$ and regularity at the second pole of the compact de Sitter geometry. On shell, the action of the bounce relative to the false vacuum is

$$B = S_E[\text{bounce}] - S_E[\text{false}], \quad S_E = 4\pi^2 \int d\xi\, \left(\rho^3 V - 3M_{\rm Pl}^2 \rho\right), \quad S_E^{\rm dS} = -\frac{24\pi^2 M_{\rm Pl}^4}{V_f} \tag{6}$$

In the flat-space limit $B = 2\pi^2 \int dr\, r^3\, [\frac{1}{2}\varphi'^2 + V - V_f]$, and for a thin wall $B = 27\pi^2\sigma^4/(2\Delta V^3)$ with $R_c = 3\sigma/\Delta V$, $\sigma = \int \sqrt{2V}\, d\varphi$ [11]. With the field in units of f and lengths in units of $(\sqrt{\lambda} f)^{-1}$ the flat-space action is $B = b(\eta)/\lambda$ exactly, and gravity enters only through g.

The Hawking–Moss action is $B_{\rm HM} = 24\pi^2 M_{\rm Pl}^4 (1/V_f - 1/V_{\rm top})$ [13]; an oscillating CDL bounce requires $|V''_{\rm top}| > 4H^2$ [15, 16], which fails for sufficiently flat potentials or strong gravity.

3.2 Numerical method and validation

The bounce is found by shooting on the central field value with overshoot/undershoot classification (adaptive eighth-order Runge–Kutta, relative tolerance 10^{-11}) [38, 39, 40], with the equations integrated through the wall and the action accumulated along the solution. For the gravitational problem the tail of the bounce outside the wall follows the false-vacuum de Sitter

geometry $\rho = \sin(H\xi)/H$ toward the second pole, and the difference from the background action is taken analytically there, which avoids the cancellation of terms of order $1/g^2$. In the thin-wall regime shooting is exponentially sensitive to the initial condition, so an independent collocation boundary-value solver is used for $\eta < 0.1$, and the two agree to 2×10^{-9} where both apply. The analytic thin-wall formula with gravity [14] serves as a further check of the gravitational suppression. Table 2 lists the checks of the bounce; the complete suite has 24 checks, all executed by a script distributed with the paper.

Table 2: Executable consistency checks of the numerical bounce.

Check	Result
Thin-wall limit: B/B_{thin} as $\eta \to 0$	0.9943 at $\eta = 0.03$, 0.9773 at 0.06 (monotone)
Shooting versus boundary-value solver	max relative difference 1.8×10^{-9} (7 cases)
Scaling $B = b(\eta)/\lambda$	max relative difference 9.8×10^{-13}
de Sitter action $-24\pi^2 M^4/V_f$	relative difference 4×10^{-16}
Friedmann constraint along the bounce	max residual 5.3×10^{-12}
Gravitational suppression against the analytic thin-wall (Parke) value	agree to 1.5% in B/B_{flat} (4 points)
Hawking–Moss consistency	84 bounces found in 98 grid points; none with $\lvert V''\rvert < 4H^2$; none above B_{HM}
Interpolation of $b(\eta)$ off the table	max relative error 6.6×10^{-5}
Derrick identity $2T + 4U = 0$ for the bounce	residual 1.4×10^{-7} at the worst of four tilts




Figure 1: (a) The potential for $\lambda = 1$, $\eta = 0.30$. (b) The bounce profile in flat space and with gravity ($g = 0.1$). (c) The numerical action $b(\eta) = B\lambda$ against the thin-wall formula; the false minimum disappears at $\eta_c = 0.385$.

3.3 Error budget and negative modes

Table 3 lists how the flat-space action changes when each numerical choice is varied: the integrator tolerance and the integrator itself, the quadrature of the action integral, the tolerance of the bisection on the central field value, the truncation of the domain, and, for the boundary-value solution, the tolerance, the initial mesh and the padding of the domain. At the production settings the numerical uncertainty of B is below 1×10^{-8}, many orders below the uncertainty of the prefactor ($\delta \ln \Gamma = O(1)$) and below the precision of any quantity compared with the data. The only sensitive setting is the bisection: a tolerance of 10^{-6} in the central field value changes

B by up to 1×10^{-4} in the thin-wall regime, which is why 10^{-13} is used.

Table 3: Numerical error budget of the flat-space action (largest relative change of B over the tilts studied).

Source	Range explored	Largest change
Integrator tolerance (DOP853)	10^{-8} to 10^{-13} (production 10^{-11})	3×10^{-9} (2×10^{-11} for tolerance $\leq 10^{-10}$)
Integrator	DOP853, RK45, LSODA, Radau	3×10^{-10}
Quadrature of the action	2×10^4 to 2×10^6 points	2×10^{-14}
Bisection on the central value	10^{-6} to 10^{-13} (production 10^{-13})	1×10^{-4} (1×10^{-8} for tolerance $\leq 10^{-10}$)
Truncation of the domain	action cut at $r_{1/2} + k/m_f$, k up to 16	1×10^{-6} at the largest radius reached
Boundary-value solver: tolerance, mesh, padding	10^{-6} to 10^{-10}; 500 to 8000 nodes; 10 to 120	3×10^{-12}
Shooting against boundary-value solution	$\eta = 0.10$	2×10^{-9}
Gravity coefficient $\kappa(g \to 0)$	integrators, tolerances, quadrature	$\kappa = 3.6407$, spread 3×10^{-5}

A CDL bounce is a saddle point with exactly one negative mode, which is what allows $\mathrm{Im}\, E$ to be interpreted as a decay rate [11, 12]. For $\eta = 0.05$, 0.15, 0.30 and 0.36 the spectrum of $-\nabla_4^2 + V''(\varphi_b)$ was computed in the angular sectors $l = 0, \ldots, 4$ (with $\psi = r^{-3/2}u$ the radial operator is $-u'' + [(l(l+2)+3/4)/r^2 + V'']u$, discretized on 30000 points). In every case the $l = 0$ sector has exactly one negative eigenvalue (lowest values -0.002, -0.024, -0.21, -0.74 in units of the false-vacuum mass squared), the $l = 1$ sector has the lowest eigenvalue consistent with the four translational zero modes (at most 2×10^{-7} of m_f^2 in magnitude) and none is negative, no eigenvalue is negative for $l \geq 2$, the field is monotonic, so that $\varphi - \varphi_f$ has no node, and the Derrick identity holds to 1×10^{-7}. With gravity the negative-mode problem is subtle and has been discussed in the literature [41]; here $g \sim 10^{-60}$, the operator differs from its flat-space form by a relative $O(g)$, and the flat-space structure applies. The gravitational spectrum was not computed for large g.

Corrections of relative size 10^{-59} cannot be resolved in double precision, and the paper does not claim to resolve them. The flat-space action is computed with no gravity at all; the gravitational correction is the linear response $B = B_{\mathrm{flat}}(1 - \kappa g)$ with $\kappa(g \to 0) = 3.641$, obtained from $g = 10^{-3}$ and 2×10^{-3} with the linear extrapolation $\kappa_0 = 2\kappa(10^{-3}) - \kappa(2 \times 10^{-3})$ (the two values differ by 2.6%, the curvature of $B(g)$) and stable to 3×10^{-5} across integrators. At $g \sim 10^{-60}$ the correction $\kappa g \sim 9.7 \times 10^{-60}$ is some 59 orders of magnitude below every other uncertainty of the rate, so what matters is only that gravity enters as a small smooth perturbation, which the Hawking–Moss criterion confirms independently (Section 3.4).

3.4 Gravitational corrections and the Hawking–Moss limit

Fig. 2a shows the ratio $B(g)/B_{\mathrm{flat}}$ computed on a grid of g and η. For small g the suppression is linear, $B = B_{\mathrm{flat}}(1 - \kappa g)$ with $\kappa = 3.5$ to 6.7 for $0.12 \leq \eta \leq 0.37$ (measured at $g = 10^{-3}$; below that the cancellation of the $1/g^2$ terms limits the numerical precision to about 10^{-4} in B/B_{flat}).

The suppression grows with g as the bubble approaches the Hubble radius, and for $g \gtrsim 1$ no oscillating bounce is found: the numerical existence agrees with the condition $|V''_{\rm top}|/4H^2 > 1$ (Fig. 2b), which is satisfied everywhere a bounce was found.




Figure 2: (a) Gravitational suppression of the action for several tilts; crosses mark the grid points at which no oscillating bounce was found. (b) The Hawking–Moss criterion $|V''_{\rm top}|/4H^2$ against g (lines, analytic; markers, numerical bounce found/not found). The shaded band is the physical value fixed by the dark energy, $g \sim 10^{-60}$.

At the physical $g \sim 10^{-60}$ the relative gravitational correction is $\kappa g \sim 9.7 \times 10^{-60}$, the Hawking–Moss criterion is satisfied by a factor $\sim 10^{59}$–10^{60}, the Hawking–Moss action is $\ln B_{\rm HM} \simeq 279$ (an exponent of 10^{121}), and the bubble radius is $\sim 2 \times 10^{-30}$ of the Hubble radius. Gravity and the Hawking–Moss channel play no role in a decay of the present dark energy within this class of potentials. They matter only if $f \gtrsim 0.3 M_{\rm Pl}$, which with V_f fixed requires $V_f/f^4 \lesssim 9 \times 10^{-119}$, a barrier $\gtrsim 1 \times 10^{118}$ times the vacuum energy, and an action beyond any consequence for the transition.

4 Nucleation rate, cooling and survival in the observed universe

4.1 The rate

The rate per unit four-volume is $\Gamma/V_4 = A\, e^{-B}$ [11, 12]. In the rescaled variables $\Gamma/V_4 = K f^4 \lambda^2 e^{-B}$, where $K = (B/2\pi)^2 D$ contains the four translational zero modes and an unknown dimensionless factor D from the remaining fluctuation determinant. Dividing by $H^4 = V_f^2/(9M_{\rm Pl}^4)$ and using $f^4 = V_f/(r_L \lambda \delta)$,

$$\frac{\Gamma}{H^4} = \frac{9K\lambda}{r_L\, \delta(\eta)} \frac{M_{\rm Pl}^4}{V_f}\, e^{-B(\lambda,\eta)} \tag{7}$$

The large number $M_{\rm Pl}^4/V_f = 1.38 \times 10^{120}$ is the number of Planck-volume cells in a Hubble four-volume. It compensates an action of $B \simeq \ln(M_{\rm Pl}^4/V_f) = 276.6$ plus the logarithm of the prefactor. Setting $\Gamma/H^4 = 1$ gives the critical action

$$B_c = \ln\frac{M_{\rm Pl}^4}{V_f} + \ln\frac{9K\lambda}{r_L\,\delta}, \qquad B_c \simeq 287 \;\; \text{(median over the prior; 5–95\% range 280–294)} \tag{8}$$

For a mean time to nucleation, an observer in the asymptotic de Sitter space sees the past-cone four-volume grow at the rate $(4\pi/3)H^{-3}$, so the mean time for one Hubble patch to nucleate a bubble is $t_{\rm dec} = H^{-1}/[(4\pi/3)\Gamma/H^4]$ for $\Gamma/H^4 \ll 1$. For $\lambda = 1$, $\eta = 0.30$ the action is $B = 441$ and the mean decay time is 10^{76} yr; at $\eta = 0.32$ ($B = 301$) it is 10^{15} yr; at $\eta = 0.34$ ($B = 194$) the rate is so large that the vacuum would not have survived and the model is excluded (Table 4).

Table 4: Benchmark potentials with the dark-energy density as false-vacuum energy ($r_L = 1$). f is the field scale, $g = f^2/M_{\rm Pl}^2$, δ_B the relative gravitational correction to B, and R_b/R_H the bubble radius over the Hubble radius.

λ	η	f (meV)	B	$\ln\Gamma/H^4$	$\log_{10} t_{\rm dec}$ (yr)	g	δ_B	$\lvert V''_{\rm top}\rvert/4H^2$	R_b/R_H
1	0.30	3.05	441	-153	76	1.6×10^{-60}	5.6×10^{-60}	5.3×10^{59}	2.6×10^{-30}
1	0.32	3.00	301	-13	15	1.5×10^{-60}	5.4×10^{-60}	4.6×10^{59}	2.5×10^{-30}
1	0.34	2.96	194	93	-31	1.5×10^{-60}	5.3×10^{-60}	3.8×10^{59}	2.3×10^{-30}
0.3	0.34	4.00	645	-357	165	2.7×10^{-60}	9.7×10^{-60}	2.1×10^{59}	3.2×10^{-30}
3	0.25	2.42	349	-60	36	9.9×10^{-61}	3.8×10^{-60}	1.2×10^{60}	2.3×10^{-30}
10	0.18	1.94	363	-72	41	6.4×10^{-61}	3.2×10^{-60}	2.9×10^{60}	2.2×10^{-30}

4.2 Survival of the present vacuum: the past light cone

A bubble nucleated at an event p expands at essentially the speed of light and reaches an observer at O if and only if p lies in the causal past of O. If nucleations are Poisson distributed with rate Γ per unit four-volume, the probability that none has occurred in the past light cone of the present observer is [19, 20]

$$P_{\rm surv} = e^{-\Gamma V_4}, \qquad V_4 = \frac{4\pi}{3}\int_0^{t_0} dt\, a(t)^3\chi(t)^3, \qquad \chi(t) = \int_t^{t_0}\frac{dt'}{a(t')} \tag{9}$$

with χ the comoving distance of the light cone at time t in a flat universe. A common order-of-magnitude shortcut uses the Euclidean volume of the de Sitter four-sphere, $8\pi^2/(3H^4)$; that is not the four-volume of any region of the observed spacetime, and it overstates V_4 by the factor given below. The observed history is $H^2 = H_0^2(\Omega_m a^{-3} + \Omega_r a^{-4} + \Omega_\Lambda)$ with $\Omega_\Lambda = 0.6889$, $\Omega_r = 9.14\times10^{-5}$ (from $T_{\rm CMB}$ and $N_{\rm eff} = 3.046$) and $\Omega_m = 0.3110$. Two independent quadratures give $V_4 = 0.12457\,H_0^{-4} = 0.0591\,H_\Lambda^{-4}$, agreeing to 1×10^{-13}; the result is converged in the lower cutoff of the scale factor and in the grid to 10^{-11} and reproduces the analytic Einstein–de Sitter value to 2×10^{-13} (check V14). The Euclidean volume is $26.3\,H_\Lambda^{-4}$, 445 times larger. Of V_4, 35% comes from events at redshift below 1, 73% below 2 and 97% below 5 (Fig. 3); the uncertainty from Ω_Λ at the Planck 1σ level is about 3.5%.

The observation that no bubble has reached us is excluded at 95% confidence if $\Gamma V_4 > -\ln 0.05 = 3.00$, that is for $\Gamma/H^4 > 50.7$ or $B < B_c - 3.9$. This is a weak condition: it is 1334 times weaker than the Euclidean estimate ($\Gamma/H^4 > 0.038$), it moves the boundary of the allowed region by only




Figure 3: Four-volume of the past light cone of the present observer, accumulated over events at redshift below z, for the observed expansion history, against the Euclidean four-sphere volume.

$\Delta B = 7.2$ out of about 287, and it bounds the future mean decay time only by $t_{\rm dec} \gtrsim 8.2 \times 10^7$ yr (the formula for $t_{\rm dec}$ is used here as an order-of-magnitude statement, since Γ/H^4 is not small). Because the prior mass sits overwhelmingly at $B \ll B_c$ or $B \gg B_c$, the fraction of the prior excluded changes from 15.0% (Euclidean estimate) to 14.8% (Section 4.5). The criterion assumes that Γ was constant, with the scalar in its false minimum throughout; it does not require observers to be present before the transition, and it says nothing about bubbles nucleated outside our past cone. Percolation of bubbles in de Sitter space requires a rate of order unity in Hubble units [21, 22], since otherwise the false-vacuum volume grows without bound and no wall collides with another. With the corrected survival bound this no longer conflicts with survival: the window $1 \lesssim \Gamma/H^4 < 51$ has the width $\Delta B = 3.9$ and contains 0.12% of the prior volume (0.18% for a percolation threshold of 0.1 and 0.05% for 10).

4.3 The prefactor and the validity of the semiclassical rate

The factor D is the ratio of the fluctuation determinants with the zero modes removed; methods to compute it for specific potentials exist [17, 18] and are not repeated here. It is treated as a systematic uncertainty. Because $\ln D$ enters B_c additively, a change of D by n decades moves B_c by $2.3\,n$: three decades give ± 7 out of 287, six decades give ± 14 (5%), and the prior mass of every regime changes by less than 10^{-3} over spreads of 0, 3 and 6 decades (Table 5). The semiclassical expansion is controlled by $1/B$, which is 3.5×10^{-3} at B_c, so that higher-loop corrections change $\ln\Gamma$ by $O(1)$ and not $O(B)$, which the prefactor spread already covers. The regime in which the approximation fails is $B = O(1)$ to $O(10)$, where the wall is not thin on the scale of the field fluctuations; 0.5% of the prior has $B < 10$ and 2.5% has $B < 30$, and none of these draws survives the survival condition (all have $\Gamma/H^4 \gg 51$), so the failure of the semiclassical approximation affects only the excluded region and cannot change the classification of any draw that is allowed.

4.4 The falsification diagram

Fig. 4 maps the decay time over the (η, λ) plane for $r_L = 1$. The region excluded by survival and the region of effective stability ($t_{\rm dec} > 10^{100}$ yr) are separated by a band: the decay time runs from the survival bound of 8.2×10^7 yr to 10^{20} yr within a change of $\Delta B \simeq 28$ in the action and to 10^{100} yr within $\Delta B \simeq 212$. The model is thus either excluded, or it describes a vacuum that decays on a time scale of 10^{20} yr or longer unless B is tuned to within about 28 of the survival bound, that is within 10% of its value.




Figure 4: Mean decay time of a Hubble patch in the (η, λ) plane for $V_f = \rho_\Lambda$ and a Minkowski true vacuum. Contours: survival bound (8.2×10^7 yr), 10^{12}, 10^{20}, 10^{50} and 10^{100} yr. The upper edge $\lambda = 4\pi$ is the perturbative-unitarity limit used for the prior.

4.5 Prior-sensitivity Monte Carlo

To quantify how much of the parameter space the window occupies, $1,000,000$ parameter sets are drawn from physically consistent priors ($\log_{10} \lambda$ uniform on $[-2, \log_{10} 4\pi]$, η uniform on $[0.03, 0.38]$, $\ln D$ Gaussian with a spread of 1.5 decades; the false minimum exists for every draw) and each is converted to a rate with the interpolated numerical action. The regime fractions are given in Table 5 together with batch-means standard errors (20 batches), convergence with the number of draws (Fig. 5c) and sensitivity to the priors. 14.8% of the prior volume lies above the survival bound, 0.77% decays within 10^{20} yr, 4.8% within 10^{100} yr and 80.4% shows no decay within 10^{100} yr. The fractions do not depend on the prefactor and the choice of r_L

is likewise immaterial. They do depend on the prior range of λ and η (Table 5; for $\lambda \leq 1$ the excluded fraction falls to 2.8%). They are therefore fractions of the stated prior, a property of the parameter space of this model and not measured probabilities, and they do not indicate how likely a transition is in our universe. $V_t < 0$ would enter through $r_L = 1/(1 + |V_t|/V_f) < 1$ and shifts B_c upward by $\ln(1 + |V_t|/V_f)$; the case $r_L \geq 1$ is the one sampled.

Table 5: Prior mass (%) of each regime. Decay fractions are cumulative and refer to survival-consistent draws; the last column is the fraction with no decay within 10^{100} yr. These are fractions of the stated prior, not probabilities for our universe.

Prior	above survival bound	decays by 10^{12} yr	by 10^{20} yr	by 10^{100} yr	no decay in 10^{100} yr
baseline	14.77 ± 0.04	0.269 ± 0.005	0.77 ± 0.01	4.78 ± 0.02	80.4 ± 0.0
$\sigma_{\log D} = 0$	14.74	0.270	0.76	4.80	80.5
$\sigma_{\log D} = 3$	14.75	0.257	0.75	4.79	80.5
$\sigma_{\log D} = 6$	14.74	0.267	0.76	4.79	80.5
r_L logU(1, 10)	14.71	0.264	0.76	4.80	80.5
$\lambda \leq 1$	2.76	0.122	0.35	2.50	94.7
η in [0.10, 0.38]	18.45	0.323	0.95	5.96	75.6
log10 λ in [-4, 2]	18.18	0.190	0.55	3.35	78.5




Figure 5: (a) Cumulative distribution of the decay time over survival-consistent draws. (b) Prior mass by regime with 3σ batch errors. (c) Convergence of the regime fractions with the number of draws.

4.6 Cooling and the thermally corrected potential

The calculation above treats the universe as a clock: the CMB temperature $T_\gamma = T_0/a$ falls with the expansion and enters nothing. It can enter the decay only if the scalar interacts with a thermal bath, and then through a thermal correction to the potential whose coefficient c is fixed by those interactions and is not a universal constant. Two regimes are therefore kept apart throughout: the scalar is decoupled and the decay is the zero-temperature O(4) channel of the previous sections, or it is in thermal equilibrium with the CMB, in which case the leading high-temperature term is

$$V_{\rm eff}(\varphi, T) = V_0(\varphi) + \tfrac{1}{2}\, c T^2 \varphi^2, \qquad \theta \equiv \frac{c T^2}{\lambda f^2} \tag{10}$$

The thermal term is a rescaling of the zero-temperature family, which makes the analysis exact. With $x = sy$ and $s = \sqrt{1-2\theta}$ the potential $(x^2-1)^2/8+\eta x/2+\theta x^2/2$ becomes $s^4[(y^2-1)^2/8+(\eta/s^3)y/2]$ up to a constant. Hence a false minimum exists only for

$$\eta_{\rm eff} \equiv \frac{\eta}{(1-2\theta)^{3/2}} < \eta_c \quad \Longleftrightarrow \quad \theta < \theta_c(\eta) = \frac{1}{2}\left[1-\left(\frac{\eta}{\eta_c}\right)^{2/3}\right] \tag{11}$$

and the bounce actions are those of the zero-temperature table at the shifted tilt: $B_4 = b(\eta_{\rm eff})/\lambda$ and $S_3 = (fs/\sqrt{\lambda})\, s_3(\eta_{\rm eff})$, where $s_3 = 4\pi \int \rho^2[x'^2/2 + w - w_f]d\rho$ is the O(3) action at $\lambda = 1$, $\theta = 0$ (tabulated here for the first time for this family, with the boundary-value solver for $\eta < 0.1$, and tending to the thin-wall value $16\pi\sigma^3/3\Delta V^2$ as $\eta \to 0$: the ratio is 0.995 at $\eta = 0.03$). Direct solutions of the O(4) and O(3) bounce with the thermal term in the potential agree with these scalings to 2×10^{-4} (check V19). For the benchmark $\lambda = 1$, $\eta = 0.30$: $f = 3.05$ meV, $\theta_c = 0.0765$, and today ($T_0 = 0.235$ meV) $\theta = 0.0059\, c$.

Cooling relaxes the thermal term, and the thermal history has a consequence that does not depend on the decay rate: the false minimum disappears at $T_c = f\sqrt{\lambda\theta_c/c}$, which is 2.7 meV (redshift 10) for $c = 0.1$ and 0.84 meV (redshift 2.6) for $c = 1$. A scalar in equilibrium with the CMB at higher temperatures would have settled into the true vacuum at the symmetric-to-broken transition, and the present dark energy would not be the false-vacuum energy. The observed universe in the false vacuum therefore requires that the scalar either decoupled before T_c or has $c < 1.1 \times 10^{-5}$ if the bath is coupled all the way back to recombination ($z = 1100$). The high-temperature form is an upper bound on the effect for particles heavier than T in the loop, since the true correction is then Boltzmann suppressed; the results below are therefore the largest thermal effect that the coefficient c can produce.




Figure 6: (a) Extrema of the thermally corrected potential for $\eta = 0.30$ against the thermal strength θ; the false minimum disappears at θ_c. (b) The quantum exponent B_4 and the thermal exponent S_3/T against the bath temperature for the benchmark; the O(3) saddle dominates above the crossover temperature $T_\times$ (where $S_3/T = B_4$) and the O(4) saddle below. Solid: $c = 0$ (equilibrium without thermal mass); dashed: $c = -1$; dotted: $c = +0.1$.

4.7 The cooling history and the three decay channels

The temperature follows the observed history, $T_\gamma(a) = T_0/a$ with $H(a)$ of Section 4.2, and cosmic time $t(a) - t_0 = \int_1^a da'/[a'H(a')]$. Three channels compete. The Hawking–Moss transition is the horizon-scale channel of de Sitter space; its action is $\ln B_{\rm HM} \simeq 279$ (Section 3.4) and it plays no role. The de Sitter horizon temperature $T_{\rm dS} = H_\Lambda/2\pi = 1.91 \times 10^{-31}$ meV is 8.1×10^{-31} of T_0, would equal the CMB temperature only after 69 e-folds ($\sim 10^{12}$ yr), and is not a homogeneous radiation bath; it is therefore kept separate and not combined with T_γ by a maximum. The thermal exponent at the horizon temperature, $S_3/T_{\rm dS} = 4.2 \times 10^{32}$, rules out any activation by it. That leaves quantum tunneling (the O(4) bounce with action B_4) and thermal activation (the O(3) bounce with S_3/T). At a given temperature the saddle with the smaller exponent dominates, and the rate takes its exponent and prefactor; the two rates are not added, because they are the two ends of one family of periodic instantons. The thermal prefactor is $\Gamma_T = D_T\, T^4 (S_3/2\pi T)^{3/2} e^{-S_3/T}$ with D_T of order unity (a systematic of 1.5 decades, as for the quantum prefactor).

For the benchmark the crossover is at $T_\times = 0.181$ meV, so that today the thermal saddle already dominates ($S_3/T_0 = 341 < B_4 = 441$), the crossover being reached when the CMB has cooled by a factor 1.29, and in the past the thermal exponent was far smaller (Table 6, Fig. 6b). Cooling lowers the rate for $c \geq 0$ at every epoch: the thermal exponent grows as $1/T$ and the thermal mass relaxes, so $\ln\Gamma$ falls monotonically with time (check V20). The late-time decay is therefore the zero-temperature O(4) decay of Section 4, because the thermal contribution switches off exponentially fast as $T \propto e^{-H_\Lambda t}$. For $c < 0$ (symmetry non-restoration, outside the usual sign of the thermal mass) the barrier grows with T and cooling lowers B_4, so the sign of the effect is reversed for the quantum channel, but the thermal exponent still has a minimum near $z \simeq 2$ for $c = -1$ and the past rate is not small.

Table 6: Exponents and rates for the benchmark ($\lambda = 1$, $\eta = 0.30$) as the universe cools. B_4 and S_3/T at $c = 0$; $\ln\Gamma/H_\Lambda^4$ for the decoupled scalar (O(4) only) and for the equilibrated scalar with the dominant saddle in brackets.

z	T (meV)	B_4	S_3/T	decoupled	$c = 0$	$c = 0.1$	$c = -1$
0	0.235	441	341	-153	-65 (O(3))	-62 (O(3))	-95 (O(3))
0.5	0.352	441	227	-153	50 (O(3))	54 (O(3))	4 (O(3))
1	0.470	441	170	-153	107 (O(3))	113 (O(3))	46 (O(3))
2	0.705	441	114	-153	165 (O(3))	173 (O(3))	66 (O(3))
3	0.939	441	85	-153	194 (O(3))	205 (O(3))	49 (O(3))

4.8 Survival with a time-dependent rate and the thermal prior maps

With a rate that depends on the temperature, the expected number of nucleations in the past light cone is

$$I = \frac{4\pi}{3} \int_{t_i}^{t_0} dt\, \Gamma(T_\gamma(t))\, a(t)^3 \chi(t)^3, \qquad P_{\rm surv} = e^{-I} \tag{12}$$

with χ as in Eq. (9). It assumes a flat universe and a homogeneous rate per physical four-volume, and walls at the speed of light. For a constant rate it reduces to Eq. (9), reproduced by the quadrature to 7×10^{-9}. For the benchmark, $\ln I = -155$ for the decoupled scalar (the vacuum survives), while an equilibrated scalar gives $\ln I = 99$ ($c = 0$, bath coupled for $z \leq 1$), 186 ($z \leq 3$) and 240 ($z \leq 10$): the benchmark is excluded by many orders of magnitude as soon as it is in equilibrium with the CMB. The same calculation over the priors of Section 4.5 (2×10^5 draws, the same seed, plus a thermal prefactor spread of 1.5 decades) gives Table 7 and Fig. 7.

Table 7: Prior mass (%) by thermal scenario (2×10^5 draws; z is the redshift below which the scalar is in equilibrium with the CMB). "Inconsistent" draws have no false minimum at some redshift of the equilibrium epoch. Decay fractions refer to the allowed draws and to the zero-temperature decay of the future.

Scenario	above survival bound	no false minimum in the past	allowed	decays by 10^{20} yr	no decay ir
decoupled	14.7	0.0	85.3	0.79	80.5
$c = 0$, $z \leq 1$	24.2	0.0	75.8	0.00	75.8
$c = 0$, $z \leq 3$	33.9	0.0	66.1	0.00	66.1
$c = 0$, $z \leq 10$	49.3	0.0	50.7	0.00	50.7
$c = 0.1$, $z \leq 1$	24.9	1.3	73.8	0.00	73.8
$c = 0.1$, $z \leq 3$	33.6	7.4	59.0	0.00	59.0
$c = 1$, $z \leq 1$	23.2	17.6	59.2	0.00	59.2
$c = -0.1$, $z \leq 1100$	38.0	0.0	62.0	0.00	62.0
$c = -1$, $z \leq 1100$	23.8	0.0	76.2	0.17	75.4




Figure 7: (a) Rate history of the benchmark: the decoupled O(4) rate is constant, the equilibrated rate falls as the universe cools; the dashed line is the survival bound on Γ/H_Λ^4. (b) Prior mass by thermal scenario.

The results answer the question posed by cooling. First, a scalar that is not coupled to the CMB is unaffected; the decay of the previous sections stands. Second, a scalar in equilibrium with the CMB is constrained more strongly, not less: the prior mass above the survival bound rises from 14.7% to 24% ($z \leq 1$), 34% ($z \leq 3$) and 49% ($z \leq 10$) for $c = 0$, and a further 1.3% to 18% of the draws have no false minimum in the equilibrium epoch for $c = 0.1$ and 1. Third, in no equilibrated scenario with $c \geq 0$ does a survival-consistent draw decay within 10^{20} yr: the decay window of Section 4.4 closes completely. The reason is that the allowed draws must have a thermal exponent large enough at $z \sim 1$, and because $S_3/T < B_4$ above $T_\times$ this requires B_4 well

above the critical action, so that the zero-temperature decay time of the future exceeds 10^{100} yr. Fourth, only the non-standard sign $c = -1$ leaves a window (0.17% of the prior against 0.79% for the decoupled scalar). In short, cooling cannot raise the probability of a late-time transition: it lowers the thermal channel monotonically, the late-time decay is the zero-temperature O(4) decay, and the thermal coupling only adds past constraints.

5 What the nucleated region can become

5.1 Energy budget

A thin-wall bubble of true vacuum is bounded by a wall that carries the released energy; its interior is, to leading order, the true vacuum with no excess energy to reheat. In the thick-wall regime relevant here the field at the centre of the bounce, x_0, lies above the true minimum, and the energy the interior can release is $\epsilon_{\rm in}\Delta V = V(x_0) - V_t$. Fig. 8c shows $\epsilon_{\rm in}$ against η: at $\eta = 0.30$, $\epsilon_{\rm in} = 0.018$, so 1.8% of ΔV resides in the interior and the remainder in the wall; $\epsilon_{\rm in}$ approaches unity only near η_c, where the barrier is gone.

5.2 Interior geometry and reheating

The interior of a CDL bubble has open spatial slices, $ds^2 = -dt^2 + a(t)^2 dH_3^2$ [23, 24], with the Friedmann constraint $(da/dt)^2 = 1 + a^2\rho/(3M_{\rm Pl}^2)$. Without a long inflationary stage inside the bubble (Section 5.3), the energy density is negligible against the curvature term at early times: $\alpha^2\rho \leq 1.1 \times 10^{-61}$ in units where $H_\Lambda = 1$, so the interior is a Milne universe, $a = t$. The condensate begins to oscillate at $t_0 = 1/m$ with $m = 3.61$ meV, behaves as matter and decays to radiation at a rate $\Gamma_\varphi \leq m$. With $\tau = H_\Lambda t$ the radiation density has the closed form $\rho_r(\tau) = \epsilon\,\tau_0^3\,\gamma\,\tau^{-4}\int_{\tau_0}^{\tau} s\,e^{-\gamma(s-\tau_0)}ds$ in units of V_f, which we compare with a numerical integration of the full system (relative difference 6×10^{-7}). The radiation temperature is $T = (30\rho_r/\pi^2 g_*)^{1/4}$ with $g_* = 3.36$.

Energy conservation bounds the reheating temperature independently of the details of the decay,

$$T_{\rm reh} \leq \left[\frac{30\,\epsilon_{\rm th}\,\epsilon_{\rm in}\Delta V}{\pi^2 g_*}\right]^{1/4}, \qquad \epsilon_{\rm th} \leq 1 \tag{13}$$

which for the largest release allowed when $V_t \geq 0$, $\epsilon_{\rm in}\Delta V = \Delta V \leq V_f = \rho_\Lambda$, gives $T_{\rm reh} \leq 2.19$ meV (25 K), against the CMB temperature of 0.235 meV today and 4.6×10^8 times below the ~ 1 MeV required for nucleosynthesis [42]. Equivalently, a thermal bath at 1 MeV needs an energy density 1.4×10^{35} times the dark-energy density. The closed form shows that the situation is worse than the bound: the radiation temperature after its maximum falls as $T \propto 1/\tau$ because the open interior keeps expanding, with $T\tau = 5.46 \times 10^{-43}$ GeV, so $T = 1$ MeV would have to occur at $t = 3.0 \times 10^{-22}$ s, before the condensate even starts to oscillate ($t_0 = 1.8 \times 10^{-13}$ s). Table 8 gives the maximum temperature for decay rates from $\Gamma_\varphi = m$ down to $10^{-20}m$ and the temperature one Hubble time after nucleation, $\sim 10^{-30}$ meV.

Table 8: Reheating of the bubble interior. ϵ: fraction of ΔV in the interior; Γ_φ/m: decay rate of the oscillating condensate; $T_{\max}$: maximum radiation temperature; the fourth column is $T_{\max}$ over the bound of Eq. (13); the fifth is the temperature one Hubble time $1/H_\Lambda$ after nucleation; the last is $T_{\rm BBN}/T_{\max}$.

ϵ	Γ_φ/m	$T_{\max}$ (meV)	$T_{\max}/T_{\rm bound}$	$T(1/H_\Lambda)$ (meV)	$T_{\rm BBN}/T_{\max}$
1 (upper bound)	1	1.2445	0.57	8.7×10^{-31}	8.0×10^{8}
1 (upper bound)	0.001	0.2318	0.11	4.1×10^{-30}	4.3×10^{9}
1 (upper bound)	1e-09	0.0073	0.00	1.3×10^{-28}	1.4×10^{11}
1 (upper bound)	1e-20	0.0000	0.00	7.3×10^{-26}	7.7×10^{13}
0.018 ($\eta=0.30$)	1	0.4558	0.57	3.2×10^{-31}	2.2×10^{9}
0.018 ($\eta=0.30$)	0.001	0.0849	0.11	1.5×10^{-30}	1.2×10^{10}
0.018 ($\eta=0.30$)	1e-09	0.0027	0.00	4.7×10^{-29}	3.7×10^{11}
0.018 ($\eta=0.30$)	1e-20	0.0000	0.00	2.7×10^{-26}	2.1×10^{14}




Figure 8: (a) Maximum reheating temperature against the released energy density; the dark-energy density gives about 2 meV, far below the energy density needed for 1 MeV. (b) Temperature of the open interior after reheating ($\Gamma_\varphi = m$): the radiation cools as $1/t$ and never approaches nucleosynthesis. (c) Fraction of ΔV carried by the interior against the tilt.

5.3 Scope of the bound: what is excluded by the model and what only by energy conservation

The bound of Eq. (13) is a statement about a locally homogeneous conversion of a vacuum-energy difference that does not exceed the dark-energy density. It is not a universal prohibition on high-temperature bubble interiors, and it is stated here with its assumptions: (A1) one real scalar field minimally coupled to gravity, with the potential of Section 2; (A2) the false-vacuum energy is the observed dark energy; (A3) the interior is homogeneous and isotropic on open slices; (A4) no inflationary stage inside the bubble; (A5) all energy densities are sub-Planckian. Models that violate one of these need a separate analysis. Table 9 lists the possible additional energy sources and says which of them the model excludes by more than energy conservation.

The geometric argument is the following. In the open slicing the curvature parameter $Y \equiv a^2\rho/(3M_{\rm Pl}^2)$ obeys $(da/dt)^2 = 1 + Y$. A hot big bang at nucleosynthesis needs $Y \sim 1$: in standard radiation domination $Ht = 1/2$, that is $\rho t^2 = 3M_{\rm Pl}^2/4$. In the curvature-dominated phase $a = t$ and $Y = \rho t^2/(3M_{\rm Pl}^2)$ can only decrease, as $1/t$ for matter and $1/t^2$ for radiation, unless the interior inflates. The energy density of the condensate and its decay products obeys the envelope

$$\rho(t) \le \epsilon\,\Delta V\,(mt)^{-3} \quad (t > t_0 = 1/m), \qquad Y(t_0) = \frac{\epsilon\,\Delta V}{3M_{\rm Pl}^2 m^2} \tag{14}$$

because matter dilutes more slowly than radiation and decay only moves energy from the slower to the faster component (checked against the closed form to machine precision, check V17). The mass at the true minimum is $m = \kappa_m \Delta V^{1/4}$ with $\kappa_m = \lambda^{1/4}[(3x_t^2 - 1)/2]^{1/2}\delta^{-1/4}$, which lies between 0.50 and 4.1 over the whole prior, so $Y(t_0) = c\sqrt{\Delta V}/M_{\rm Pl}^2$ is tiny for any sub-Planckian energy: $Y(t_0) = 2.0 \times 10^{-63}$ at $\Delta V = V_f$, and $Y(t_0) = 1$ would need $\Delta V/M_{\rm Pl}^4 = 1.8 \times 10^5$ at the benchmark, which is super-Planckian. Therefore, for any ΔV below the Planck scale, the interior is a Milne universe at all times, Y never approaches unity (Fig. 9), and the energy density at one second after nucleation cannot exceed $\epsilon\Delta V/(m \cdot 1\,\mathrm{s})^3$. Requiring $\rho(1\ \mathrm{s}) \ge \rho_{\rm BBN} = (\pi^2 g_*/30)(1\ \mathrm{MeV})^4$ ($g_* = 10.75$) demands $\Delta V^{1/4} \ge 5.1 \times 10^{61}$ GeV, that is 42, 43 and 45 orders of magnitude above the Planck mass (for the smallest κ_m in the prior, the benchmark and the largest). For the dark energy the envelope gives $\rho(1\ \mathrm{s}) = 4.4 \times 10^{-74}\,\rho_{\rm BBN}$.

Three points follow. First, the anti-de Sitter case shows that energy conservation alone is not sufficient: a very deep true vacuum releases enough energy for 1 MeV, but its open interior then collapses within seconds, and the longest-lived interior has $T \lesssim 4$ meV. Second, the geometric statement is independent of the energy scale and of the value of V_f, and therefore covers an additional reservoir too: it cannot raise Y above the curvature bound unless its energy density exceeds ΔV by 9.1×10^{60}. Third, the escape routes are exactly those the model lacks: a long inflationary stage inside the bubble, a parent energy scale close to the Planck scale, or an additional mechanism that raises Y at t_0. A successful cosmology would have to lie in that outside class.

A second scalar field χ does not change this. For any number of scalar fields with a potential $V(\varphi, \chi)$, the energy released in the transition of a homogeneous region is the potential difference




Figure 9: Curvature parameter $Y = a^2\rho/3M_{\rm Pl}^2$ of the bubble interior against the time since nucleation for $\Gamma_\varphi = m$. A flat-like radiation- or matter-dominated epoch needs $Y \sim 1$; the interior starts at $Y \sim 10^{-63}$ and moves away from it.

Table 9: Energy sources and assumptions behind the reheating bound, and what excludes each.

Source or assumption	What bounds it	Quantitative statement
Released vacuum energy, $V_t \geq 0$	energy conservation, $\Delta V \leq V_f$	$T_{\rm reh} \leq 2.19$ meV
Wall energy $(1-\epsilon_{\rm in})\Delta V$	already counted in ΔV; collisions redistribute it	benchmark: 98% of ΔV in the wall; the total is unchanged
Deeper true vacuum, $V_t < 0$ (anti-de Sitter)	geometry: crunch at $t_c = \pi\sqrt{3}M_{\rm Pl}/\sqrt{\lvert V_t\rvert}$ and the curvature envelope	$T_{\rm max} = 1$ MeV needs $\lvert V_t\rvert = 1.4\times10^{35}\,V_f$ and a crunch after 4.6 s; a life of 13.8 Gyr needs $\lvert V_t\rvert \leq 15.7\,V_f$ and $T \leq 4.4$ meV
Interior inflation	the potential has no plateau above V_f; field range $\Delta\varphi/M_{\rm Pl} \sim 10^{-30}$	at most 9.3×10^{-62} e-folds (largest over four tilts)
Larger parent scale ($V_f \gg \rho_\Lambda$)	geometry: the envelope $\rho \leq \epsilon\Delta V(mt)^{-3}$	$\rho(1\text{ s}) \geq \rho_{\rm BBN}$ needs $\Delta V^{1/4}$ 42 to 45 orders above $M_{\rm Pl}$
Extra reservoir (other fields, radiation)	not part of the model; subject to the same curvature constraint	$Y \sim 1$ at $t = 1/m$ needs $\rho_x = 9.1\times10^{60}\,\Delta V$
Inhomogeneity, bubble collisions	not included (A3); total energy is bounded by ΔV	local concentration possible; the average is unchanged

between the false and true vacua, $V_f - V_t$, and the false-vacuum energy $V_f = V(\varphi_f, \chi_f)$ is the whole observed dark energy ρ_Λ: any energy stored in χ while it sits in the false vacuum gravitates as dark energy and is already counted in V_f. A larger release needs $V_t < 0$ (the anti-de Sitter row of Table 9, a crunch within seconds), and the curvature argument above is independent of the field content. A reservoir that is not vacuum energy (radiation, matter, kinetic energy of χ) would have been present and gravitating throughout the cold era, where observations limit it to a fraction of the critical density, whereas the interior needs $\rho_x = 9.1 \times 10^{60}\,\Delta V$. A coupled two-field model is therefore a relabelling of ΔV and not a source of new energy, unless the second sector inflates or carries a Planck-scale parent energy, which is the outside class of Section 7. The simulations of Section 5.5 confirm it.

5.4 Other fates and the singularity question

If the true vacuum has negative energy ($V_t < 0$) the open interior collapses to a singularity in a finite time, $\tau_c = \pi/\sqrt{|V_t|/V_f}$: $|V_t| = 0.01\,V_f$: 547 Gyr; $|V_t| = 0.1\,V_f$: 173 Gyr; $|V_t| = 1\,V_f$: 55 Gyr; $|V_t| = 10\,V_f$: 17 Gyr. A decay to a de Sitter vacuum with $V_t > 0$ leaves the interior with an accelerating phase and no radiation epoch. Only $V_t \simeq 0$ leaves a regular, expanding, but empty interior. No case yields a nonsingular bounce from the parent geometry: the bubble is a region of the parent de Sitter space, causally connected to it, and the Milne interior is geodesically complete only because it has no matter.

The distinction between daughter regions and a cyclic cosmology (Section 1) applies to these fates. The nucleated region is a daughter region in the sense of Option A: a cold, curvature-dominated open region bounded by the wall inside the parent de Sitter space and still causally connected to it, not a new hot universe. The potential has no further metastable minimum at the true vacuum, so repeated transitions of the interior would need a different potential; the entropy produced in reheating is bounded by the energy ΔV and is carried by an open interior of infinite comoving volume, so no entropy statement can be made for the daughter as a closed system. A cyclic cosmology (Option B) needs a nonsingular bounce, which in spatially flat general relativity with conventional matter requires a violation of the null energy condition ($\dot{H} = -(\rho+p)/2M_{\rm Pl}^2 > 0$) and raises problems of stability and entropy [9, 10]; the present model contains no such mechanism. Entropy accounting and causality are computed for the daughter region in Section 5.5; the stability of an interior bounce is not.

5.5 Lorentzian simulations of the daughter region: two fields, reservoirs and repeated transitions

The interior was also simulated, not only bounded. The analytic continuation of the bounce gives the open homogeneous interior with the field starting at the bounce centre x_0 at rest and $a = t$ (Milne). In units of f for the fields and λf^4 for energy densities, with $g = f^2/M_{\rm Pl}^2$, the system is

$$\ddot{x}_i + \left(3\frac{\dot{a}}{a} + \Upsilon_i\right)\dot{x}_i + \frac{\partial V}{\partial x_i} = 0, \qquad \dot{\rho}_r + 4\frac{\dot{a}}{a}\rho_r = \sum_i \Upsilon_i \dot{x}_i^2, \qquad \ddot{a} = -\frac{g\,a}{6}(\rho + 3p) \tag{15}$$

with $\rho = \frac{1}{2}\sum_i \dot{x}_i^2 + V + \rho_r$, $p = \frac{1}{2}\sum_i \dot{x}_i^2 - V + \rho_r/3$ and the constraint $\dot{a}^2 = 1 + g\,a^2\rho/3$, whose relative residual is the energy-conservation check (eighth-order Runge–Kutta, relative tolerance 10^{-11}). The friction Υ_i is a phenomenological dissipation that moves field energy to radiation with total energy conserved. The simulations are the following (check V22–V24 and Table 10–13).

(A) Single field, physical benchmark ($\lambda = 1$, $\eta = 0.30$, $f = 3.05$ meV, $g = 1.6 \times 10^{-60}$). The full dynamics give a maximum radiation temperature within 7% of the closed form of Section 5.2 for Γ_φ/m from 1 to 10^{-3} (ratios 0.93, 1.04, 1.06 and 1.06), so the assumption that the condensate starts to oscillate at $1/m$ and dilutes as matter is sound. The constraint residual is at most 9×10^{-9} for $g \geq 10^{-3}$ (it scales with g and is trivially small at $g = 10^{-60}$, so the gravitational scan below is the real test). Scanning g from 10^{-60} to 10^3, the curvature parameter $Y = ga^2\rho/3$ reaches 3.9×10^{-2} at $g = 10^1$ and 0.48 at $g = 10^2$; a flat-like radiation epoch ($Y \geq 1/4$ with radiation dominant) first appears at $g = 10^2$, that is $f = 10\,M_{\rm Pl}$. For the dark-energy scale $g = 10^{-60}$ and $Y_{\max} = 3.9 \times 10^{-63}$.

(B) Two fields with an energy reservoir. A second field χ (here $V_\chi = \frac{\lambda_\chi}{4}(\chi^2 - w_\chi^2)^2$ with $\lambda_\chi = 1$, $w_\chi = 1.5f$, coupled by $\frac{1}{2}g_c^2(\varphi - \varphi_t)^2\chi^2$ so that it is stable in the false vacuum and tachyonic in the true one) stores 1.27 λf^4, 4.3 times the energy ΔV released by φ. A constant offset $V_{\rm off} = -(\Delta V + E_\chi)$ makes the false vacuum have the observed dark-energy density, which is the only way the reservoir can sit in the parent without gravitating. The true vacuum is then anti-de Sitter with $|V_t| = \Delta V + E_\chi$, and the simulation (dissipation $\Upsilon = m/2$ in both fields, χ seeded at $10^{-6}f$) transfers the reservoir to radiation. The depth is set in units of the dark energy, $|V_t| = \kappa\rho_\Lambda$, which fixes f and g (Table 10). For $\kappa = 10^{35}$ (enough for 1 MeV) the radiation reaches 0.72 MeV after 8.9×10^{-21} s, and the interior crunches after 5.46 s, equal to the pure anti-de Sitter value $\pi\sqrt{3}M_{\rm Pl}/\sqrt{|V_t|}$ to better than 10^{-12}. The temperature then falls as $1/t$, with $Tt = 10.9$ for this reservoir (between 10.7 and 13.5 for reservoirs from 0.02 to 20 in units of λf^4), so that at one second $T = 7.6 \times 10^{-24}$ GeV for every depth, 1×10^{20} times below 1 MeV. The envelope of Section 5.3 holds with $\rho(mt)^3/E_{\rm rel} \leq 11$ for the single field and 4.3×10^3 for the reservoir, whose hilltop delays the roll by $\sim 16/m$; with that factor the parent scale needed for 1 MeV at 1 s falls by 3.6 orders of magnitude to still 38 orders above the Planck scale. The reservoir therefore does not make a hot, lasting epoch: it makes a hot instant followed by a crunch.

Table 10: Two-field reservoir compensated by a negative offset: the true vacuum has $|V_t| = \kappa\rho_\Lambda$. f: field scale; $T_{\max}$ and its time; crunch time of the interior; temperature one second after nucleation.

κ	f (GeV)	$T_{\max}$ (MeV)	time of $T_{\max}$ (s)	crunch (s)	$T(1\text{ s})$ (GeV)
1×10^{0}	2.0×10^{-12}	1.3×10^{-9}	5.0×10^{-12}	1.7×10^{18}	7.2×10^{-24}
1×10^{10}	6.4×10^{-10}	4.0×10^{-7}	1.6×10^{-14}	1.7×10^{13}	7.2×10^{-24}
1×10^{20}	2.0×10^{-7}	1.3×10^{-4}	5.0×10^{-17}	1.7×10^{8}	7.2×10^{-24}
1×10^{30}	6.4×10^{-5}	4.0×10^{-2}	1.6×10^{-19}	1.7×10^{3}	7.2×10^{-24}
1×10^{35}	1.1×10^{-3}	7.2×10^{-1}	8.9×10^{-21}	5.5×10^{0}	7.6×10^{-24}

(C) Flattening at the dark-energy scale. With a field of vev w and $V = \frac{\lambda_c}{4}(\chi^2 - w^2)^2$, $\lambda_c w^4/4 = 3H_\Lambda^2 M_{\rm Pl}^2$ (so that the false-vacuum energy is ρ_Λ) and χ seeded at $10^{-5}w$, hilltop inflation inside the open interior gives the e-folds and curvature of Table 11 (gravity fully included, $g = 1$ in these

units, constraint residual at most 5×10^{-9}). Flatness ($N \geq 60$, $\Omega_k \lesssim 10^{-52}$) needs $w \gtrsim 4M_{\rm Pl}$: a super-Planckian vev. The reheating temperature is 0.67–1.45 meV in every case, below the bound of Eq. (13) by construction, since the energy released is ρ_Λ. A hot epoch at 1 MeV needs the same dynamics at a parent energy 1.4×10^{35} times the observed dark energy.

Table 11: Hilltop inflation of a field with vev w inside the open interior at the dark-energy scale ($H_\Lambda = M_{\rm Pl} = 1$): e-folds with $q < -0.1$, flatness $N = -\frac{1}{2}\ln\Omega_{k,\rm min}$, and the maximum radiation temperature.

$w/M_{\rm Pl}$	λ_c	$N_{\rm accel}$	$N_{\rm flat}$	$\Omega_{k,\rm min}$	$T_{\rm max}$ (meV)
1	12.000	8.2	6.8	1.2×10^{-6}	1.45
2	0.750	20.5	18.9	3.6×10^{-17}	1.18
3	0.148	38.1	36.3	3.2×10^{-32}	1.02
4	0.047	61.0	59.0	6.1×10^{-52}	0.91
5	0.019	89.3	87.1	2.2×10^{-76}	0.83
6	9.3×10^{-3}	123.1	120.7	1.4×10^{-105}	0.77
8	2.9×10^{-3}	207.0	204.4	2.8×10^{-178}	0.67




Figure 10: Simulated daughter region. (a) Radiation temperature after nucleation for a reservoir hidden by an anti-de Sitter vacuum of depth $|V_t| = \kappa\rho_\Lambda$ (dots: maximum; dashed: crunch); the green band is the nucleosynthesis window, reached by neither. (b) Hilltop flattening at the dark-energy scale: e-folds against the vev (left) and the maximum temperature (right), with the bound of Eq. (13).

(D) Repeated transitions, entropy and causality. A chain of vacua $(\varphi_f, \chi_f) \to (\varphi_t, \chi_f) \to (\varphi_t, \chi_t)$ with a Minkowski end point releases ΔV_1 and then ΔV_2 out of the parent of energy ρ_Λ. The energy released over all generations telescopes, $\sum_i \Delta V_i = \rho_\Lambda - V_{\rm final} \leq \rho_\Lambda$, so the total radiation, and its temperature $T \leq 2.19$ meV, is the same as for a single transition. Each generation has its own decay time (Table 12, from the rate of Eq. (7) with the parent energy of that generation); the second generation of the daughter has a Hubble rate $\sqrt{\Delta V_2/\rho_\Lambda}$ times that of the parent and decays no sooner than 10^{21} yr in the cases that survive. Repeated transitions are therefore possible in principle, but they add no energy. The entropy per comoving curvature volume produced by the benchmark interior is 0.31 in units of f^{-3} and is constant after reheating; the entropy inside a sphere of the bubble-wall radius at the time of the temperature maximum is 4.9, 2×10^{-63} of the area bound $8\pi^2 M_{\rm Pl}^2 R_b^2 = 2.2 \times 10^{63}$ and 2×10^{-122} of the de Sitter horizon

entropy of the parent (3.3×10^{122}), so the generalized second law is satisfied with enormous margin and gives no constraint. The interior is the future light cone of the nucleation event, with infinite comoving volume and a logarithmically divergent conformal time, so it contains no region causally connected to the parent except through the wall.

Table 12: Chain of two transitions out of the dark-energy vacuum: tilt/coupling of each step, energy released in units of ρ_Λ, decay time of each generation, total released and the resulting temperature bound.

step 1 (η/λ)	step 2 (η/λ)	$\Delta V_1/\Delta V_2$	$\log_{10} t_1$ (yr)	$\log_{10} t_2$ (yr)	$\sum \Delta V/\rho_\Lambda$	$T_{\max}$ (meV)
0.30 / 1.0	0.30 / 1.0	0.6 / 0.4	76	76	1.00	2.19
0.32 / 1.0	0.28 / 2.0	0.6 / 0.4	16	21	1.00	2.19
0.30 / 1.0	0.33 / 0.5	0.5 / 0.5	76	96	1.00	2.19
0.27 / 3.0	0.27 / 3.0	0.5 / 0.5	-7	-8	1.00	2.19

Table 13: Criteria for a viable daughter universe, by scenario: parent vacuum energy in units of the observed dark energy, maximum temperature, a hot epoch (1 MeV at 1 s), flatness $\Omega_{k,\min}$, lifetime, and the criterion that fails.

Scenario	parent energy/ρ_Λ	$T_{\max}$	hot at 1 s	$\Omega_{k,\min}$	lifetime	fails
Single field, decoupled (benchmark)	1	0.43 meV	no	1 (Milne)	unbounded	none
Reservoir in χ, compensated ($\kappa = 10^{35}$)	1, with a cancellation of 10^{35}	0.72 MeV	no (7.6×10^{-24} GeV at 1 s)	1 (Milne)	5.5 s	crunch
Hilltop at ρ_Λ ($w = 5M_{\rm Pl}$)	1	0.83 meV	no	2×10^{-76}	unbounded	energy
Hilltop at a parent $1.4 \times 10^{35}\,\rho_\Lambda$ ($w = 5M_{\rm Pl}$)	1.4×10^{35} (excluded)	1 MeV	yes	2×10^{-76}	unbounded	survival

Table 13 states the conditional criteria for renewal in the sense of Option A. A daughter region is a viable hot universe only if the parent vacuum energy is at least 1.4×10^{35} times the observed one (excluded by the data), or if a reservoir is hidden by a cancellation of the same size, which leaves a true vacuum that crunches within 5.5 s. No scenario with the observed dark energy as the parent satisfies the energy, flatness and lifetime criteria together.

6 Matter dilution and decoherence as a trigger

The original motivation for a late-time transition was that dilution of matter reduces environmental decoherence and thereby permits a macroscopic quantum transition. In the baseline model nothing of the kind appears: the potential $V(\varphi)$, hence B, does not depend on the matter density, and the Gibbons–Hawking temperature $T_H = H/2\pi = 1.9 \times 10^{-43}$ GeV [43] is 6×10^{-32} of the field scale; thermal-type corrections to B are of relative size $\sim g\,(\kappa \simeq 3.5)$, that is 10^{-59} (Section 3.4). A dependence on matter must be introduced by a coupling, and a conclusion that held only for one coupling would be weak. The requirement can be stated without choosing one.

Any coupling of non-relativistic matter of mass $m_{\rm mat}(\varphi)$ adds $\rho_m \ln m_{\rm mat}(\varphi)$ to the effective potential, so that the energy difference between the vacua changes by $\rho_m\,\Delta \ln m$, where $\Delta \ln m$

is the fractional change of the matter mass between the two field values and $\rho_m = (\Omega_m/\Omega_\Lambda)V_f$ today. Write $\varepsilon_m \equiv \rho_m \Delta \ln m/\Delta V$ for the relative shift. To first order in the tilt shift $\Delta\eta$,

$$\frac{\Delta\eta}{\eta} = \frac{2\,\delta(\eta)}{\eta\,(x_f - x_t)}\,\varepsilon_m, \qquad \Delta B = B\left|\frac{d\ln b}{d\ln\eta}\right|\frac{\Delta\eta}{\eta} \tag{16}$$

with $\delta(\eta) = \Delta V/(\lambda f^4)$ of Eq. (3). A change of the rate by a factor e needs $\Delta B = 1$; moving a benchmark from $B = 441$ to the critical action $B_c = 287$ needs $\Delta B = 154$ ($\Delta B = 13.7$ for $\eta = 0.32$). With $|d\ln b/d\ln\eta| = 5.5$ at $\eta = 0.30$ this gives $\Delta \ln m = 8.9 \times 10^{-4}$ and 1.4×10^{-1} respectively: the matter mass must differ by between about 0.1% and 10% between the two vacua. Table 14 translates this to specific couplings.

Table 14: Requirement on a matter coupling that changes the rate by a factor e (left end) or moves the benchmark into the reset window (right end).

Coupling	Requirement	Comment
General	$\Delta \ln m \geq 8.9 \times 10^{-4}$ to 1.4×10^{-1}	independent of the coupling function; $\eta = 0.30$
Linear, $m = m_0 e^{\beta\varphi/M_{\rm Pl}}$	$\beta \geq 3.7 \times 10^{26}$ to 5.7×10^{28} (for $\eta = 0.32$: 4.7×10^{26} to 6.4×10^{27})	$\Delta\varphi/M_{\rm Pl} = 2.4 \times 10^{-30}$; unscreened tests restrict $\beta \lesssim O(1)$; screening (chameleon type) lowers the local coupling, not the cosmological one [35]
Quadratic, $m = m_0(1 + \varphi^2/2\Lambda_q^2)$	$\Lambda_q \leq 58$ to 5 meV ($\eta = 0.30$)	$f = 3.0$ meV, so $\Lambda_q \sim$ (2–20)f: a new interaction at the scale of the potential; symmetron-type screening [36] can hide it in dense environments
Density-dependent effective potential	same $\Delta \ln m$	the model is then specified by $\rho_m F(\varphi)$ with F changing by that amount across $\Delta\varphi$

The conclusion is therefore conditional. For the simplest linear coupling the required strength lies 26 to 28 orders of magnitude above what unscreened laboratory and solar-system tests allow. Quadratic or screened couplings are not excluded by this paper: they evade local tests by construction, but they are tuned, because they require a coupling scale of the same size as the dark-energy potential and a mass change of at least 0.1% of the matter between vacua. The dilution of matter can act as a trigger only through such an added, tuned structure, not through the framework itself. A noise-assisted mechanism would need an explicit open-system influence functional and a calculated change of the tunneling exponent, which is not assumed here.

7 Consistency conditions and falsification tests

A model of this class must satisfy internal consistency conditions, which are mathematical or follow from the framework, and it can be confronted with observational tests. The two are kept apart because failing the first means the model is ill-defined, while failing the second means that it is disfavored by data. All are computed in this paper and can be recomputed for a different potential with the supplied code.

7.1 Internal consistency conditions

- C1 (existence). A metastable minimum requires $\eta < \eta_c = 0.385$; for larger tilt there is nothing to decay.
- C2 (semiclassical validity). The loop expansion parameter $1/B$ must be small. All draws with $B < 30$ (2.5% of the prior) have rates far above the survival bound, so the classification of allowed draws is unaffected (Section 4.3).
- C3 (negative mode and flat limit). The bounce has one negative mode and gravity enters as an $O(g)$ perturbation, $g \sim 10^{-60}$ (Section 3.3).
- C4 (energy). With $V_t \geq 0$ the released energy density cannot exceed ρ_Λ, so $T_{\rm reh} \leq 2.2$ meV; with $V_t < 0$ the interior crunches (Section 5.3).
- C5 (geometry). For any sub-Planckian energy the open interior stays curvature dominated, $Y(t) \leq Y(t_0) = 2.0 \times 10^{-63}$, unless it inflates; the single-field potential gives at most 9.3×10^{-62} e-folds (Section 5.3).
- C6 (percolation). A global transition requires $\Gamma/H^4 = O(1)$ or larger, a window of width $\Delta B = 3.9$ below the survival bound that holds 0.12% of the prior volume.
- C7 (thermal consistency). A scalar in equilibrium with the CMB at redshift z keeps its false minimum only if $c < \theta_c \lambda f^2 / T_\gamma(z)^2$, i.e. $c < 1.1 \times 10^{-5}$ for a bath coupled back to recombination, and the survival condition then applies to the rate history $\Gamma(T_\gamma(t))$ (Section 4.8).
- C8 (repeated transitions). Over any chain of transitions the energy released telescopes, $\sum_i \Delta V_i = \rho_\Lambda - V_{\rm final} \leq \rho_\Lambda$, so a second generation cannot raise the temperature above 2.2 meV (Section 5.5).

7.2 Observational tests

- O1 (survival). The absence of a bubble in our past light cone requires $\Gamma V_4 < 3.0$, i.e. $\Gamma/H^4 < 51$ at 95% confidence, which excludes 15% of the prior volume of the benchmark family and constrains the future mean decay time only to $t_{\rm dec} \gtrsim 8.2 \times 10^7$ yr.
- O2 (reset window). A transition on cosmological time scales requires $B_c \lesssim B \lesssim B_c + 28$ for decay within 10^{20} yr; this holds for 0.77% of the prior volume. An action above $B_c + 212$ puts the decay beyond 10^{100} yr.
- O3 (hot epoch). A hot universe in the nucleated region requires $\rho_{\rm reh} \gtrsim (1\ {\rm MeV})^4$, which is 1.4×10^{35} times the dark-energy density; the bound of Section 5 excludes it for decay of the present dark energy in this framework.
- O4 (trigger). Matter dilution as the trigger needs $\Delta \ln m \gtrsim 10^{-3}$ between the vacua, a linear coupling of $\beta \sim 10^{26}$–10^{29} or a screened coupling at a scale of tens of meV (Section 6), which local gravity and fifth-force tests can probe if the screening is incomplete.

- O5 (signals). A solitary transition produces no signal in our patch; bubble collisions [25, 33, 34] require a population and $\Gamma/H^4 = O(1)$ (C6). The vacuum is indistinguishable from a cosmological constant ($w = -1$) until it decays. A measured evolution of the dark energy, which recent baryon-acoustic-oscillation analyses combined with the CMB and supernovae prefer at the level of a few standard deviations depending on the supernova sample [44], would favor a rolling field and not a metastable vacuum. No observational discriminant of the cyclic reset is available from this model.

- O6 (cooling). If the scalar is coupled to the CMB the decay probability was larger in the past and falls as the universe cools: equilibrated scenarios with $c \geq 0$ lift the excluded prior mass from 15% to 24–49% and leave no survival-consistent draw that decays within 10^{20} yr, so the absence of a transition so far constrains the coupling of the scalar to the CMB as well as its potential (Section 4.8).

A transition producing a successful cosmology would have to lie outside the class studied here. It would need a parent vacuum energy of at least $\sim 10^{34}\rho_\Lambda$ (an earlier, higher-energy phase of the universe rather than the present one) and, by Section 5.3, a post-nucleation region that inflates before it reheats, so that the curvature term and the horizon are handled, and a mechanism to restore the high-energy parent from the late universe. The cyclic models that have been constructed provide such mechanisms by other means [4, 5]; this paper shows that tunneling of the present dark energy does not.

8 Discussion and limitations

- A single scalar field and a quartic-tilt potential are used. Other potentials change $b(\eta)$ and the interior energy fraction, but not the scales: the rate depends on the combination $M_{\rm Pl}^4/V_f$ and the interior energy on V_f, both fixed by the data, so O1, O3 and the geometric argument of Section 5.3 are generic.

- The one-loop prefactor is a systematic uncertainty, not a result; the conclusions are insensitive to it over six decades (Section 4.3).

- The decay is treated as a flat-space process because $g \sim 10^{-60}$; the gravitational code is validated against analytic limits and used for the map of the regime in which gravity matters, which the dark-energy scale excludes.

- The survival criterion assumes a constant rate and a false vacuum throughout the history of the observed universe; a field that evolved or was thermally affected in the early universe would change the early part of the cone, which carries a small part of V_4.

- The interior evolution assumes a homogeneous open slicing and a perturbative decay of the condensate to radiation; bubble-wall collisions can concentrate energy locally, but the average energy density is bounded by ΔV (Table 9).

- Up-tunneling from the false to a higher vacuum, the decay of the true vacuum in curved space [45], thermal activation of the Hawking–Moss type [30] and decays of the present vacuum into other vacua are not considered.
- The priors of the Monte Carlo are documented assumptions, not measurements; the regime fractions describe the sample space of the model, not the probability of a transition in our universe.
- The matter-coupling analysis is first order in the tilt shift and does not simulate any specific screened model.
- The thermal analysis assumes equilibrium with the CMB (an upper bound on the thermal channel), the leading high-temperature term with one coefficient c, and a bath coupled below a stated redshift; it neglects higher-order thermal corrections and the interpolation between the O(4) and O(3) saddles near $T_\times$, which a periodic-instanton calculation would refine.
- The simulations of Section 5.5 are homogeneous and use a phenomenological dissipation: bubble-wall dynamics, inhomogeneous wall collisions, the microphysics of reheating and the gravitational backreaction of the wall are not simulated, and a nonsingular bounce (Option B) is not modelled.

9 Conclusions

A late-time false-vacuum transition of the present dark energy is a well-defined calculation, and for the explicit potential studied it can be done to a stated precision. The quantum decay is a flat-space process controlled by the action $B = b(\eta)/\lambda$, which must lie near $B_c \approx 287$ for the transition to be neither excluded nor effectively absent; with survival evaluated on the past light cone the bound is weak ($\Gamma/H^4 < 51$) and the window is set by the action itself. Cosmological cooling does not help: a thermal mass rescales the potential exactly, the thermal O(3) channel dominates above 0.18 meV and falls monotonically as the universe cools, and a scalar in equilibrium with the CMB is more strongly constrained (24–49% of the prior excluded against 15%) with no late-time decay window left. The nucleated daughter region carries at most the dark-energy density, reheats to at most 2.2 meV and cools as $1/t$ in a curvature-dominated open universe; this rests on geometry and not on energy conservation alone, holds for any sub-Planckian parent energy and any number of scalar fields, and a trigger by matter dilution would need a change of the matter mass between the vacua of at least 0.1%. A daughter region of the present dark energy is cold and empty, and the cyclic-reset hypothesis, in the form tested, is not supported; the framework identifies what would have to change.

Lorentzian simulations of the daughter region (Section 5.5) support these statements with two fields, dissipation and gravity: the closed-form interior is reproduced to 7%, a reservoir hidden by a negative offset leaves an anti-de Sitter vacuum that crunches after 5.5 s when it reaches 0.7 MeV, flattening at the dark-energy scale needs a vev above $4M_{\rm Pl}$ and reheats below 1.5 meV, and repeated transitions release no more than ρ_Λ in total. A viable hot daughter universe needs

a parent vacuum energy at least 1.4×10^{35} times the observed dark energy, which is excluded, or a cancellation of the same size, which ends in a crunch.

Declarations

Funding. This research received no specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Conflict of interest. The author declares no competing interests.

Ethics approval, consent to participate, consent for publication. Not applicable.

Data availability. The data underlying every figure and table (bounce tables, error budget, survival calculation, falsification map, Monte Carlo summary, interior evolution, scope and coupling estimates, verification results) are deposited with the code at https://github.com/sandlerleon/late-time-vacuum-decay and archived at https://doi.org/10.5281/zenodo.23752471.

Code availability. All code (Python, NumPy, SciPy, Matplotlib) is released under the MIT licence at https://github.com/sandlerleon/late-time-vacuum-decay, archived at https://doi.org/10.5281/zenodo.2
Running the scripts in the stated order regenerates every number in this paper, and verify.py runs 24 executable checks. The manuscript is archived at https://doi.org/10.5281/zenodo.23752470.

Authors' contributions. L.S.: conceptualization, methodology, software, formal analysis, investigation, visualization, writing.

Use of generative AI. During the preparation of this work the author used Claude (Anthropic) to assist with the design of the numerical analysis, writing of the code, literature checking and drafting. The author reviewed and edited the content, checked the code against analytic limits and executable tests, verified the cited references, and takes full responsibility for the publication. No AI tool is an author.

References