

Self-Maintained Order and Hysteretic Collapse in a Non-Equilibrium Rotational Lattice

Order-dependent degradation and repair as a source of nonlinear feedback, bistability, and active maintenance

Leon Sandler

Independent Researcher, Northbrook, Illinois, USA

Corresponding author: sandler.leon@gmail.com

ORCID: <https://orcid.org/0009-0007-4584-808X>

Abstract

We study the maintenance of directional order in a non-equilibrium rotational lattice: a two-dimensional array of interacting rotors whose ordering couplings are removed by an order-dependent flux and restored by repair. Because degradation accelerates with local orientational disorder, the stationary state is a non-equilibrium steady state (NESS). In an annealed mean-field reduction controlled by the ratio ρ of repair to shielded degradation, the feedback makes order bistable: for cooperativity $h > 1$ and gain above a computed threshold, two saddle-node bifurcations bound a hysteresis window, whereas without feedback there is one fixed point, $I^* = 1/(1 + \rho)$. Kinetic Monte Carlo tests this. When degradation senses only the nearest neighbours the lattice is not bistable: the hysteresis loop closes as the sweep slows, its area vanishing in the quasi-static limit for $L = 16$ – 64 , and ordered and collapsed states converge throughout the window, because averaging the steep feedback over a broad distribution of local disorder flattens it. When degradation senses disorder over larger regions bistability returns, robustly up to $L = 96$ when the closure itself is imposed on the lattice. Two further results are established: an exact order–entropy relation, $dS/dU_\chi = -\kappa$ for the maximum-entropy von Mises law, and finite-size scaling ($L = 16$ – 96) estimating the field-free loss of stiffness at a diluted fraction $f_{KT} \approx 0.205$, far below site percolation (0.407), a Berezinskii–Kosterlitz–Thouless mechanism. The measured maintenance current peaks where order is most fragile. Static hysteresis is predicted only when degradation senses order over an extended range.

Keywords: nonlinear dynamics; non-equilibrium statistical mechanics; bistability and hysteresis; Berezinskii–Kosterlitz–Thouless transition; entropy production; kinetic Monte Carlo

29 1. Introduction

30 Many physical systems maintain order not as a thermodynamic ground state but as a dynamical balance: a
31 directionally ordered medium is continuously eroded by an external flux while a maintenance process
32 restores it [1–3]. Order then persists or collapses according to a competition of rates. When the erosion
33 depends on the current state of the medium, the balance is governed by feedback, which admits the
34 multistability and the dissipation characteristic of non-equilibrium systems [4–6]. Instances include the
35 orientational order of a polymer film under photodegradation with self-healing [7], the alignment of
36 magnetic domains under thermally activated flips [8] with active error-correcting rewrites, and the
37 directional order of a molecular monolayer under chemical attack with turnover [9].

38 The setting differs from the best-known orientational non-equilibrium systems. In flocking and active-
39 matter models, driving generates and stabilizes order through self-propulsion and alignment, and long-
40 range order can survive in two dimensions where the equilibrium XY model cannot [10, 11]; in the
41 equilibrium XY model, order is lost through the Berezinskii–Kosterlitz–Thouless (BKT) transition [12].
42 Here order is neither generated by the drive nor protected by an equilibrium ground state: it is maintained
43 against a state-dependent loss. Keeping information-bearing order away from equilibrium also has a
44 thermodynamic price [13, 14], of which kinetic proofreading is the classic example in which dissipation
45 buys fidelity [15], and which stochastic thermodynamics makes quantitative [16, 17].

46 We formulate this as a non-equilibrium interacting particle system [18] in which reversible rotor
47 dynamics is coupled to a non-conservative, order-dependent site process. The coupling produces a
48 nonlinear mean-field reduction that is bistable, with saddle-node bifurcations bounding a hysteresis
49 window in which a maintained and a collapsed NESS coexist (Section 4.2), a behavior absent without the
50 feedback (Section 4.1). We then test the reduction on the lattice by kinetic Monte Carlo designed to
51 separate static from rate-dependent hysteresis (Section 4.3) and find that its bistability survives only when
52 degradation senses order over an extended region; with nearest-neighbour sensing the lattice has a single
53 NESS, and we identify why (Section 4.4). Three further contributions support the picture: an exact
54 relation between a directional order parameter and configurational entropy, stated in the form needed here
55 from exponential-family theory (Section 3); a finite-size-scaling analysis locating the field-free transition
56 relative to site percolation and identifying a BKT mechanism (Section 5); and the measured maintenance
57 current, with the entropy production of an auxiliary cycle as an illustrative thermodynamic reading
58 (Section 6). The order-dependence of the degradation, a site in a disordered neighborhood degrading
59 faster, is what makes the stationary state a genuine NESS rather than a product of an uncoupled two-state
60 chain and an equilibrium rotor gas, and is the source of the nonlinear phenomenology.

61 2. The model

62 Let $\Lambda = \{1, \dots, L\}^2$ be a square lattice with $N = L^2$ sites and periodic boundaries. Each site i carries a rotor
 63 orientation $\theta_i \in [0, 2\pi)$ and a binary integrity $g_i \in \{1, g_0\}$, $0 < g_0 \ll 1$. The energy is

$$64 \quad H(\theta, g) = -J_0 \sum_{\langle ij \rangle} g_i g_j \cos(\theta_i - \theta_j) - h_0 \sum_i g_i \cos \theta_i. \quad (1)$$

65 The dynamics interleaves a reversible rotor relaxation with a non-conservative, order-dependent site
 66 process.

67 *Reversible rotor relaxation.* At fixed g , the orientations evolve by a Metropolis kernel reversible with
 68 respect to the Gibbs measure $\propto e^{-H/T}$ ($k_B T = 1$); alone, this equilibrates the rotors in the quenched integrity
 69 landscape and satisfies detailed balance.

70 *Order-dependent site process.* Each intact site ($g_i = 1$) is *diluted* ($g_i \rightarrow g_0$) at a rate that depends on its
 71 local orientational disorder,

$$72 \quad \lambda_i = \lambda_0 \left[1 + \beta \frac{\delta_i^h}{K^h + \delta_i^h} \right] \quad (2)$$

$$73 \quad \delta_i = 1 - \left| \frac{1}{4} \sum_{j \in \partial i} e^{i\theta_j} \right| \quad (3)$$

$$74 \quad \lambda_0 = k G (1 - s u_s). \quad (4)$$

75 so an aligned neighborhood shields a site ($\lambda_i \rightarrow \lambda_0$) while a disordered neighborhood accelerates its
 76 degradation ($\lambda_i \rightarrow \lambda_0(1 + \beta)$); degradation is thus *autocatalytic in disorder*. Each diluted site is *repaired*
 77 ($g_0 \rightarrow 1$, θ_i reset toward 0) at rate R . The bare susceptibility k , drive G , geometric shielding ($1 - s u_s$),
 78 cooperativity h and gain β parameterize the flux. This state dependence couples the site process to the
 79 rotor field: the two cannot be treated separately, and detailed balance is broken, driving the joint system to
 80 a NESS.

81 *Discrete steps and continuous rates.* The simulations advance in discrete steps: the rotors are relaxed,
 82 then every site is updated synchronously, with λ_i and R used as per-step probabilities (Appendix A). The
 83 rate equations of Section 4 are the continuous-time description of the same process, which the discrete
 84 chain approaches when $\lambda_i, R \ll 1$; the control ratio $\rho = R/\lambda_0$ is the same in both, and where the discrete
 85 update shifts a result (Section 4.1) the shift is stated.

86 The order parameter and the information variable are

$$U_\chi = \left| N^{-1} \sum_i e^{i\theta_i} \right|, \quad I = 1 - U_\chi. \quad (5)$$

3. The order–entropy bridge

Fixing the mean resultant length and maximizing differential entropy over circular densities yields the von Mises law $vM(\kappa)$ [19, 20], with density

$$f(\theta) = \frac{e^{\kappa \cos \theta}}{2\pi I_0(\kappa)}, \quad A(\kappa) = \frac{I_1(\kappa)}{I_0(\kappa)}. \quad (6)$$

where I_0 and I_1 are modified Bessel functions of the first kind [21].

Theorem 1 (Order–entropy bridge). *For the von Mises family: (i) $U_\chi = A(\kappa)$ is a smooth strictly increasing bijection $[0, \infty) \rightarrow [0, 1)$; (ii) $S(\kappa) = \ln(2\pi I_0(\kappa)) - \kappa A(\kappa)$ satisfies $dS/d\kappa = -\kappa A'(\kappa) < 0$ for $\kappa > 0$, hence $dS/dU_\chi = -\kappa$; (iii) S is a strictly decreasing invertible function of U_χ , with $S = \ln 2\pi$ at $U_\chi = 0$ and $S \rightarrow -\infty$ as $U_\chi \rightarrow 1$; (iv) $H_\chi = S_{\max} - S = -\frac{1}{2} \ln(1 - U_\chi) + \frac{1}{2}(\ln \pi - 1) + o(1)$ as $U_\chi \rightarrow 1$.*


Proof. (i) The von Mises family is an exponential family in $\cos \theta$ with log-partition function $\ln(2\pi I_0(\kappa))$, so $A(\kappa) = d \ln I_0(\kappa)/d\kappa = E_\kappa[\cos \theta]$, and $A'(\kappa) = \text{Var}_\kappa(\cos \theta)$, which is strictly positive for every $\kappa \geq 0$ because $\cos \theta$ is not almost surely constant under any von Mises law. The Bessel recurrences give the Riccati identity (see [21] for properties of the ratio A)

$$A'(\kappa) = 1 - \frac{A(\kappa)}{\kappa} - A(\kappa)^2, \quad (7)$$

and $0 < A(\kappa) < 1$ for $\kappa > 0$. With $A(0) = 0$ and the asymptotics $A(\kappa) = 1 - 1/(2\kappa) + O(\kappa^{-2})$ as $\kappa \rightarrow \infty$, A is a strictly increasing bijection $[0, \infty) \rightarrow [0, 1)$, which proves (i). (ii) Since $(d/d\kappa) \ln(2\pi I_0) = A$ and $(d/d\kappa)(\kappa A) = A + \kappa A'$, $dS/d\kappa = -\kappa A' < 0$; dividing by $dU_\chi/d\kappa = A'$ gives $dS/dU_\chi = -\kappa$. (iii) follows by the inverse-function theorem. (iv) $I_0(\kappa) \sim e^\kappa/\sqrt{2\pi\kappa}$ gives $S = \frac{1}{2} + \frac{1}{2} \ln 2\pi - \frac{1}{2} \ln \kappa + o(1)$, and inverting $A(\kappa) \approx 1 - 1/(2\kappa)$ near $U_\chi \rightarrow 1$ ($\kappa \sim 1/[2(1 - U_\chi)]$) yields the stated tail. ■

The identity $dS/dU_\chi = -\kappa$ is exact: the marginal information gained per unit increase in directional order equals the concentration of the underlying law, so the last increments toward perfect order are logarithmically costly (Figure 1).

Parts (i)–(iii) are instances of standard exponential-family identities: the mean parameter increases because its derivative is a variance, and the derivative of the entropy with respect to the mean parameter is minus the natural parameter [19, 20]. We do not claim the identity as new. Theorem 1 states it in the explicit form needed here, for the circular order parameter U_χ and with the logarithmic tail (iv), and uses it as an exact tool for translating directional order into configurational information in the rest of the paper.



115

116 **Figure 1.** The order–entropy bridge (Theorem 1). (A) S is strictly decreasing in U_χ . (B) $H_\chi = S_{\max} - S$ is a strictly
 117 increasing invertible image of U_χ , with the derived logarithmic tail (dashed).

118 4. Mean-field reduction, feedback and bistability

119 **4.1 The uncoupled baseline.** Setting $\beta = 0$ removes the feedback: the integrity of each site is then an
 120 exact two-state Markov chain (intact $\rightleftharpoons$ diluted at rates λ_0, R), independent of the rotors. The intact
 121 fraction obeys $\dot{p} = R(1 - p) - \lambda_0 p$, so with $\tau = \lambda_0 t$ and $\rho = R/\lambda_0$,

$$122 \quad \frac{dI}{d\tau} = (1 - I) - \rho I, \quad I^* = \frac{1}{1 + \rho}. \quad (8)$$

123 a globally stable fixed point approached at rate $(1 + \rho)$; the lattice $p(t)$ follows this master equation up to a
 124 discrete-time correction: the simulation updates all sites synchronously once per step, so its exact fixed
 125 point is $R/(\lambda_0 + R - \lambda_0 R)$ rather than $R/(\lambda_0 + R)$, a relative shift below $\lambda_0/(1 - \lambda_0) \approx 2\%$ that vanishes as λ_0 ,
 126 $R \rightarrow 0$ (it is checked to Monte Carlo accuracy in Appendix A) and U_χ is slaved to it (Figure 2). The linear
 127 baseline is recovered as the $\beta = 0$ limit.




Figure 2. Uncoupled baseline ($\beta = 0$, $L = 48$). (A) Lattice intact fraction from a fully intact start (three runs per ρ) and the exact mean of the synchronous two-state chain (dashed). (B) Stationary intact fraction p and order U_x against ρ , with the continuous-time value $\rho/(1 + \rho)$ and the discrete fixed point $R/(\lambda_0 + R - \lambda_0 R)$: U_x is slaved to p up to a small boundary-layer correction. (C) Relaxation of the chain to its fixed point against $\tau = \lambda_0 t$, with the continuous-time rate $1 + \rho$ (dashed).

4.2 Nonlinear feedback and bistability. With order-dependent degradation, the annealed mean-field identification $\delta \doteq I$ closes the feedback loop and the information variable obeys the nonlinear ODE

$$\frac{dI}{d\tau} = \mu(I)(1 - I) - \rho I, \quad \mu(I) = 1 + \beta \frac{I^h}{K^h + I^h}. \quad (9)$$

Proposition 2 (Bistable NESS). *Steady states solve*

$$\rho = F(I) := \frac{\mu(I)(1 - I)}{I}. \quad (10)$$

For cooperative feedback ($h > 1$ and $\beta > \beta_c(h, K)$) F is non-monotonic, so there is an interval $\rho \in (\rho_-, \rho_+)$ on which the ODE has three fixed points: two stable — a maintained (low- I) branch and a collapsed (high- I) branch — separated by an unstable one. The endpoints $\rho_{\pm}$ are saddle-node bifurcations, and the system exhibits hysteresis: decreasing ρ from the maintained state, order persists until ρ_- , then collapses; increasing ρ from the collapsed state, order is not recovered until ρ_+ . For $(h, K, \beta) = (6, 0.5, 20)$ the bistable window is $\rho \in (3.9, 11.3)$.

Proof sketch. $F(I) = \mu(I)(1 - I)/I \rightarrow \infty$ as $I \rightarrow 0^+$, because $\mu(0) = 1$, and $F(I) \rightarrow 0$ as $I \rightarrow 1^-$. For $h > 1$ and large β the cooperative term contributes a rising factor that makes F non-monotonic, with a local minimum and a local maximum whose values are ρ_- (minimum) and ρ_+ (maximum). Between them three real roots exist; linearizing $G(I) = \mu(I)(1 - I) - \rho I$ gives stability $\text{sgn}(-G'(I^*))$, stable on the outer branches and unstable on the middle one. The saddle-node conditions $F'(I) = 0$ locate $\rho_{\pm}$. ■

Where the feedback is strong enough. Cooperativity is necessary. For $h \leq 1$, $F(I) = (1 - I)/I + \beta I^{h-1}(1 - I)/(K^h + I^h)$ is a sum of strictly decreasing functions, so the steady state is unique for every β . For $h > 1$ the threshold $\beta_c(h, K)$, the smallest gain at which F develops its pair of turning points, follows from the saddle-node conditions as in the companion order–repair analysis [22]. At $K = 0.5$ it is $\beta_c = 219, 50.7, 19.1, 7.73$ and 2.42 for $h = 1.2, 1.5, 2, 3$ and 6 , diverging as $h \rightarrow 1^+$. The parameters used throughout, $(h, K, \beta) = (6, 0.5, 20)$, lie at $\beta/\beta_c = 8.3$, where the window is $\rho \in (3.92, 11.30)$; it shrinks to a point as $\beta \rightarrow \beta_c$ and widens to $(4.57, 21.8)$ at $\beta = 40$ (`mf_threshold.py` in the study repository).

Figure 3 shows the S-shaped steady-state manifold, the bifurcation diagram with its hysteresis window, and two trajectories at a common ρ relaxing to distinct (maintained vs collapsed) NESS. Bistability is the qualitative signature of the feedback: it is absent in the $\beta = 0$ baseline and appears only because degradation is coupled to the order it destroys.




Figure 3. Nonlinear feedback $\rightarrow$ bistability (Proposition 2). (A) The steady-state manifold $\rho = F(I)$ is S-shaped, so a range of ρ admits three fixed points. (B) Bifurcation diagram: stable (green) and unstable (red) branches with two saddle-node points bounding a hysteresis window (shaded). (C) At $\rho = 7.5$, inside the window, two initial conditions flow to a maintained and a collapsed NESS.

4.3 Testing the bistability on the lattice. Proposition 2 is a statement about the reduced ODE, obtained under one approximation: the *annealed mean-field closure* $\delta \doteq I$, which replaces the local neighbourhood disorder δ_i of each site by the global average I . Whether its central prediction survives on the short-range lattice is a numerical question, and a hysteresis loop recorded at one sweep rate cannot settle it: a loop opens whenever the control parameter moves faster than the system relaxes, whether or not two stable states exist [23]. We therefore use kinetic Monte Carlo [24] to apply two tests that separate dynamic from static hysteresis, at the parameters of Proposition 2 ($T = 0.45, h_0 = 0.30$) and in a colder, field-free variant ($T = 0.30, h_0 = 0$) in which thermal fluctuations are weaker. The protocol, run table, engine verification and error analysis are given in Appendix A.


175 *Sweep-rate extrapolation.* The ratio ρ is stepped from 16 down to 1 and back on a 31-point grid with a
 176 dwell of $\tau = 1$ to 256 steps at each value, at $L = 16\text{--}64$ (8 independent runs per size and rate). The signed
 177 loop area $A = \int (U_{x\downarrow} - U_{x\uparrow}) d\rho$, which unlike the absolute area is unbiased under noise, peaks at 2.5 for
 178 dwells of two to four steps and falls steadily as the sweep slows, to 0.09 at $\tau = 32$ for $L = 64$ (Figure 4B).
 179 From $\tau = 64$ on it is consistent with zero: $A = -0.007$ (95% bootstrap interval -0.020 to 0.005) over the
 180 dwells $\tau = 64\text{--}256$ at $L = 64$, $|A| \leq 0.018$ at every size in both variants, and every interval contains zero. A
 181 bistable system keeps a finite loop as $\tau \rightarrow \infty$, bounded by its two saddle-nodes; for the mean-field
 182 reduction that quasi-static area is 4.30.

183 *Two starts at fixed ρ .* Inside the mean-field window ($\rho = 4.5, 6, 8$ and 10) each run starts either from the
 184 ordered state or from a collapsed one (95% of sites diluted, angles random) and is followed for 3000 steps
 185 ($L = 16\text{--}64$, 8 runs per start). The stationary order reached from the two starts agrees to within 0.004 in all
 186 40 combinations of ρ , L and variant, with the bootstrap interval of the difference containing zero in 35 of
 187 them (Figure 4C). The seed-averaged trajectories meet within 15–115 steps, and the meeting time does
 188 not grow with the number of sites from $N = 256$ to 4096 (Figure A1C), whereas escape from a metastable
 189 branch slows exponentially with N . The stationary distribution of U_x is unimodal (bimodality coefficient $\leq$
 190 0.37, below the 5/9 of a uniform distribution [25]), and its integrated autocorrelation time is 6–11 steps,
 191 again independent of L (Figure A1B).

192 The five combinations whose interval excludes zero are, with difference and 95% interval: $T = 0.30$, $L =$
 193 32 , $\rho = 6$: -0.0021 (-0.00384 to -0.00038); $T = 0.30$, $L = 32$, $\rho = 8$: +0.0014 (+0.00045 to +0.00231); $T =$
 194 0.30 , $L = 64$, $\rho = 6$: +0.0008 (+0.00001 to +0.00144); $T = 0.45$, $L = 16$, $\rho = 10$: -0.0015 (-0.00256 to
 195 -0.00047); $T = 0.45$, $L = 64$, $\rho = 10$: -0.0005 (-0.00085 to -0.00007). Their differences are at most 0.0021,
 196 two orders of magnitude below the 0.63 separation of the mean-field branches at $\rho = 6$, and they have no
 197 common sign (23 of the 40 differences overall are positive). A Welch test per combination, corrected for
 198 the 40 comparisons by Holm's procedure, leaves none significant (smallest adjusted $p = 0.70$). With eight
 199 runs per start, percentile bootstrap intervals are known to be somewhat too narrow, so a handful of
 200 exclusions among 40 intervals is expected by chance. They do not indicate a residual dependence on the
 201 initial state.

202 Both tests give the same answer. At these parameters the nearest-neighbour lattice has a single stationary
 203 state at each ρ , and the hysteresis it shows is dynamic: a lag behind a moving control parameter that
 204 vanishes in the quasi-static limit. The slow sweeps trace one continuous crossover (Figure 4A) across the
 205 range in which the reduction predicts two branches, and at $\rho = 6$, where the reduction has stable states at
 206 $U_x = 0.86$ and 0.23 , both starts settle at $U_x = 0.74$. Mean-field bistability does not survive on the lattice at
 207 these parameters, including the colder variant. The stationary currents nonetheless balance, $J_{\text{kill}} = J_{\text{rep}}$ to

208 better than one part in 10^3 at every ρ and L (Figure 4D), so the lattice reaches a genuine NESS and the
 209 cycle current used in Section 6 is the measured one.



210

211 **Figure 4.** Kinetic Monte Carlo test of Proposition 2 on the nearest-neighbour lattice ($T = 0.45$, $h_0 = 0.30$ unless
 212 stated). (A) Sweeps of ρ down (solid) and up (dashed) at $L = 64$, with 1, 16 and 256 steps per ρ value: the loop of
 213 fast sweeps closes as the sweep slows; the mean-field window is shaded and its stable branches are grey. (B) Signed
 214 loop area against dwell per ρ value for $L = 16$ – 64 and for the colder field-free variant (squares); mean $\pm$ 95%
 215 interval over 8 runs. (C) Stationary order from an ordered and a collapsed start at fixed ρ inside the window ($L =$
 216 64): the two coincide. (D) Stationary kill and repair currents per site and step at $\rho = 4.5, 6$ and 8 and $L = 16, 32, 64$;
 217 the line is equality.

218 **4.4 Why the closure fails, and where it holds.** The reduction assumes that every intact site is degraded
 219 at the rate $\mu(I)$ set by the global disorder. The lattice applies instead the average of μ over the local
 220 disorders its intact sites actually see, $\bar{\mu} = \langle \mu(\delta_i) \rangle$, and the two differ in both directions (Figure 5A). Where
 221 order is low the local disorder of intact sites stays far below the global one: at $\rho = 2$ the global I is 0.79
 222 while intact sites see $\delta = 0.45$, so the lattice kills at $\bar{\mu} = 9.1$ against $\mu(I) = 19.8$. Part of this is a ceiling:
 223 four rotors pointing in independent random directions give $\delta = 0.55$ on average, barely above $K = 0.5$, so
 224 a nearest-neighbour sensor can never engage the switch fully. Where order is high the lattice kills faster

225 than the closure assumes: at $\rho = 6$, $\bar{\mu} = 2.47$ against $\mu(I) = 1.37$, and above even $\mu(\delta) = 1.02$, the rate at the
 226 mean local disorder. That last inequality is the signature of averaging a steep, convex switch over a broad
 227 distribution of δ_i : the few sites in disordered pockets are killed at nearly the full rate even in an ordered
 228 lattice. Together the two effects flatten the effective feedback, and a flattened μ cannot bend F into the S-
 229 shape that Proposition 2 requires. Both shrink as the neighbourhood that a site senses grows, since the
 230 average over n rotors narrows the distribution of δ_i and raises its ceiling (to 0.82 for $n = 24$ and 0.90 for n
 231 $= 80$).

232 A control experiment tests this directly. Keeping the rotor dynamics, the repair process and every
 233 parameter fixed, we let the kill rate respond to the disorder of a larger block of n surrounding sites ($n = 8$,
 234 24, 80) or to the global disorder $1 - U_\gamma$, which imposes the mean-field closure on the lattice itself ($L = 48$,
 235 4 runs per case; Figure 5B–D). With $n = 4$ the slow-sweep loop vanishes (area 0.01 at $\tau = 256$) and with n
 236 $= 8$ it nearly does (0.03); the two starts agree at every ρ . With $n = 24$ a small loop remains (0.21, still
 237 shrinking between $\tau = 64$ and 256) while the starts still agree at the values of ρ tested. With $n = 80$ the
 238 loop has area 1.00 and at $\rho = 6$ the ordered start holds $U_\gamma = 0.85$ while the collapsed start stays at 0.39
 239 throughout the run. With global sensing the two starts are separated at $\rho = 4.5$ (0.82 against 0.28) and 6
 240 (0.85 against 0.37), and the slow sweep collapses at $\rho \approx 3.7$, close to the mean-field saddle-node $\rho_- =$
 241 3.92, but recovers already at $\rho \approx 7.5$, below $\rho_+ = 11.30$: at finite N fluctuations let the collapsed state
 242 escape before its own saddle-node, so the loop (area 1.93) is about half the quasi-static mean-field value
 243 4.30.

244 Because a separation over 2000 steps at one size does not by itself establish bistability, we tested both
 245 cases in size and time (Figure A2): $L = 32, 48, 64$ and 96, 4 runs from each start, 15000 steps each. With
 246 global sensing the two starts stay on separate branches at every ρ tested inside the window (4.0, 4.5, 6.0,
 247 7.0) and every size: none of the 128 runs left its branch, and the separation, 0.43–0.54, does not change
 248 with L . That is the behaviour expected of a genuine mean-field bistability, whose metastable lifetimes
 249 grow exponentially with N . With the 80-site neighbourhood the picture is that of a discontinuous
 250 transition in a finite-range system. At $\rho = 5.5$ and 6.0 the separation (0.49 and 0.46) is again independent
 251 of L and no run left its branch. At the edges of the window, states do decay: at $\rho = 5.0$ the ordered state
 252 collapsed in 12 of 16 runs, and sooner in larger systems (after 7390–11470 steps at $L = 32$ but 1200–5070
 253 at $L = 96$), consistent with a decay rate proportional to the number of sites, as nucleation of the stable
 254 phase predicts; at $\rho = 6.5$ the collapsed state recovered in 13 of 16 runs. With finite-range sensing such
 255 nucleation-limited decay is expected to narrow the interval of true coexistence as $N \rightarrow \infty$, leaving long-
 256 lived metastability and a discontinuous jump near the value of ρ at which an interface between the two
 257 phases stalls; our data are consistent with this but do not locate that point. We therefore describe the

restored bistability as strong numerical evidence rather than proof: robust in size and time for global sensing, and for the 80-site neighbourhood a discontinuous transition with hysteresis whose infinite-size, infinite-time limit remains open (problem (b)).

The bistability of Proposition 2 is therefore not an artefact of the reduction: it is realized on the lattice when degradation responds to order averaged over a large enough region, and it is the mean-field limit of that case. It is destroyed when degradation responds only to the nearest neighbours; at the parameters studied the crossover lies between $n = 8$ and $n = 80$ sensed sites. The physical content of Proposition 2 is accordingly conditional on the range over which the degrading flux senses disorder, and that range, not only the gain β and the cooperativity h , is a control parameter of the maintained state.




Figure 5. Why nearest-neighbour feedback is not bistable, and the sensing range that restores bistability. (A) Mean local disorder of intact sites $\bar{\delta}$ against the global $I = 1 - U_x$ in the stationary lattice ($L = 64$; filled: ordered start, open: collapsed start; both parameter sets); the line is the closure $\bar{\delta} = I$ and the dotted line marks K . (B) Slow sweeps (256 steps per ρ value) at $L = 48$ with the kill rate sensing the 4 nearest neighbours (the model), blocks of 8, 24 and 80 sites, or the global disorder; solid down, dashed up; the mean-field window is shaded. (C) Difference in stationary order between the ordered and the collapsed start at fixed ρ , with 95% intervals. (D) Slow-sweep loop area against the number of sites sensed, at 64 and 256 steps per ρ value.

275 5. The field-free transition is BKT-driven

276 At $h_0 = 0$ the maintained phase requires the intact sub-lattice to sustain the quasi-long-range order of the
 277 two-dimensional XY model [12]. We locate the field-free transition of the quenched problem, in which a
 278 fixed fraction f of sites is diluted to $g = 0$, by finite-size scaling at $T = 0.45$ with $L = 16, 24, 32, 48, 64$ and
 279 96 and eight independent disorder realizations per size and dilution (Appendix A), measuring the Binder
 280 cumulant U_4 and the helicity modulus (spin stiffness)

$$281 \quad Y = \frac{1}{L^2} \left\langle \sum_b J \cos \Delta\theta \right\rangle - \frac{1}{L^2 T} \left\langle \left(\sum_b J \sin \Delta\theta \right)^2 \right\rangle, \quad (11)$$

282 averaged over the two lattice directions. The Nelson–Kosterlitz criterion [26] places the BKT transition
 283 where Y crosses the universal line $2T/\pi$. Dilution lowers the effective coupling and with it the effective
 284 transition temperature, and Y crosses $2T/\pi$ at $f_{\text{KT}}(L) = 0.220$ for $L = 16$, falling slowly to 0.210 at $L = 96$,
 285 with bootstrap intervals over realizations within ± 0.005 for $L \geq 48$ (Figure 6A, B). The drift is small and
 286 of the logarithmic kind expected at a BKT point, and two extrapolations agree: a linear fit in $1/\ln^2 L$ gives
 287 $f_{\text{KT}}(\infty) = 0.205$ (95% interval 0.193 – 0.216), and a Weber–Minnhagen fit [27] of $Y_L = (2T/\pi)[1 + 1/(2 \ln L$
 288 $+ C)]$ to the sizes $L \geq 24$ gives $f_{\text{KT}} = 0.205$ (0.198 – 0.213) with $\chi^2 = 0.75$ for 4 degrees of freedom. The
 289 estimate is robust to dropping the smallest lattices: the $1/\ln^2 L$ extrapolation gives 0.205 (0.194 – 0.216) for
 290 $L \geq 16$, 0.209 (0.203 – 0.282) for $L \geq 24$, 0.207 (0.197 – 0.227) for $L \geq 32$, 0.201 (0.181 – 0.225) for $L \geq 48$,
 291 and the Weber–Minnhagen fit gives 0.205 (0.196 – 0.213) for $L \geq 24$, 0.205 (0.196 – 0.213) for $L \geq 32$,
 292 0.200 (0.195 – 0.213) for $L \geq 48$. We report $f_{\text{KT}} \approx 0.205$ as a numerical estimate at $L \leq 96$ with eight
 293 realizations per point, not as a precisely established critical value. All these estimates lie far below the
 294 square-lattice site-percolation value $1 - p_c \approx 0.407$ [28]: the intact sub-lattice still percolates geometrically
 295 at f_{KT} , so order is lost to thermal (BKT) fluctuations rather than to geometric disconnection. The Binder
 296 cumulant (Figure 6C) stays at its ordered value up to $f \approx 0.23$ and falls between $f = 0.26$ and 0.35 , again
 297 well before percolation. The magnitude is plausible: the clean model has $T_{\text{BKT}} \approx 0.893$ [29], twice the
 298 temperature used here, and a linear interpolation of $T_{\text{BKT}}(f)$ between that value and zero at percolation, a
 299 heuristic only, would bring the transition down to $T = 0.45$ at $f \approx 0.20$.




Figure 6. Finite-size scaling of the field-free ($h_0 = 0$) transition under quenched dilution ($T = 0.45$; $L = 16$ – 96 ; 8 disorder realizations per point; error bars are 95% intervals). (A) Helicity modulus $Y(f)$ and the Nelson–Kosterlitz line $2T/\pi$ (dashed). (B) Crossing points $f_{KT}(L)$ with bootstrap intervals against $1/\ln^2 L$, the linear extrapolation (square) and the Weber–Minnhagen estimate (diamond); the dotted line is the site-percolation value $1 - p_c \approx 0.407$. (C) Binder cumulant $U_4(f)$; the dotted line marks $1 - p_c$.

6. Entropy production of an auxiliary maintenance cycle

The simulated dynamics is strictly irreversible: a killed site is never un-killed, only repaired, and no transition has a reverse partner. Its Schnakenberg entropy production is therefore formally infinite, and the simulation by itself does not measure a thermodynamic cost. To give the maintenance cycle a finite thermodynamic reading we introduce an *auxiliary* model, which is not part of the simulated dynamics. A single site has two mesostates, O (intact) and D (diluted), joined by a kill edge $O \rightarrow D$ with rate $\bar{k}$, the mean kill rate of an intact site measured in the stationary lattice, and a repair edge $D \rightarrow O$ with rate R , each given a small reverse rate ε . This is a Schnakenberg network [30] with one cycle, of affinity

$$A = \ln \left(\frac{\bar{k} R}{\varepsilon^2} \right), \quad (12)$$

and to leading order in ε its stationary current is the measured lattice current $J = \bar{k}p = J_{\text{kill}} = J_{\text{rep}}$. Its entropy production rate is

$$\sigma_{\text{aux}} = J A \geq 0 \quad \text{for } \bar{k} R \geq \varepsilon^2. \quad (13)$$

Figure 7 separates what is measured from what is modelled. The current (Figure 7B) is measured: it is set by the coupled dynamics and differs qualitatively from the uncoupled chain, which it exceeds everywhere. It vanishes as $\rho \rightarrow 0$, where little remains to be killed, peaks at $J = 0.052$ per site and step near $\rho = 4$, where the feedback is most active, and falls towards $\lambda_0 = 0.02$ in the well-maintained regime, where $\mu \rightarrow 1$. The throughput of the maintenance cycle is therefore largest where order is most fragile,

not where most order is held. The entropy production (Figure 7A) is modelled: σ_{aux} inherits from the lattice only the current J and the rates $\bar{k}$ and R , and depends logarithmically on the unspecified reverse rate ε , diverging in the irreversible limit $\varepsilon \rightarrow 0$ of the simulation. Two statements survive any choice of small ε : $\sigma_{\text{aux}} > 0$ throughout, so any thermodynamically consistent completion of the dynamics with weak reverse rates assigns the maintenance cycle a positive dissipation; and its dependence on ρ follows the measured current. Its absolute value does not survive, and we do not claim that the simulation establishes a quantitative thermodynamic cost of maintaining order or information. That would require the microscopic reverse rates of a specific degradation and repair chemistry (open problem (c)).




Figure 7. Maintenance current and the auxiliary cycle. (A) Entropy production rate $\sigma_{\text{aux}} = J \ln(\bar{k}R/\varepsilon^2)$ of the auxiliary two-state cycle of Section 6 for three reverse rates ε (an illustrative, regularization-dependent quantity, not a property measured in the simulation). (B) Stationary kill current J per site and step measured in the lattice ($L = 48$, 4 runs per point; the repair current coincides) for both parameter sets, the uncoupled two-state chain (dashed) and λ_0 (dotted).

7. Generality and instances

The construction assumes only three ingredients: a directional order parameter admitting a maximum-entropy law (Theorem 1), an order-dependent flux that removes ordering couplings (the feedback of Section 4), and a repair process that restores them. Systems satisfying these ingredients are expected to exhibit analogous feedback-driven behavior, although the quantitative thresholds and bifurcation structure will depend on their microscopic dynamics — real systems may carry additional degrees of freedom (spatial correlations, memory, non-Markovian repair) that this minimal model does not capture. Instances

include the orientational order of a polymer film degraded by ultraviolet chain scission and restored by annealing or self-healing chemistry; magnetic storage domains under thermally activated flips with active error-correcting rewrites (repair); liquid-crystal directors under photochemical damage; and the chiral order of a self-assembled molecular monolayer under non-enzymatic attack with turnover, the applied problem from which this model was abstracted [9]. In each, u_s encodes the geometry throttling the flux and ρ measures repair relative to shielded, order-dependent degradation. We emphasize that these are candidate realizations of the same qualitative mechanism, not demonstrated physical instances: the present study is analytic and computational, with no experimental data used (Data and code availability), and the framework should be read as a minimal model that may be applicable to systems with order-dependent degradation and active repair rather than as a validated description of any one of them.

7.1 A falsifiable experimental signature

The mechanism makes a concrete prediction that does not require the rotor language of the model, and Sections 4.3 and 4.4 sharpen it. In any driven, actively maintained, orientationally ordered medium (a magnetic-domain array, a self-healing polymer film, an aligned colloidal or active-particle ensemble), sweep the ratio of repair to degradation down from a well-ordered state and back up at several sweep rates, holding all other conditions fixed. Every such medium shows a loop when the sweep is fast, so the discriminating observable is the loop area extrapolated to zero sweep rate, supported by the dependence of the final state on the initial one at fixed ρ . A finite static loop, with order lost below a threshold ρ_- and not regained until $\rho_+ > \rho_-$ however slowly the sweep is made, indicates feedback acting over an extended range, as in Proposition 2. A loop that closes as the sweep slows, as on the nearest-neighbour lattice, indicates feedback that is local, too weak ($\beta < \beta_c$) or not cooperative ($h \leq 1$); uncoupled two-state degradation shows only the trivial lag of Section 4.1. Where the spatial range over which degradation responds to order can be varied, for example through the diffusion length of a degrading species, the crossover of Section 4.4 is itself testable. This gives Sections 3–6 a shared empirical target across the candidate systems of this section.

8. Status of the claims and approximations

For transparency at the level expected in theoretical statistical physics, we state each principal claim, its logical status, and its support.

Table 1. Status of the principal claims, the logical status of each and its support.

Claim	Status	Support
Order–entropy bijection, $dS/dU_\chi = -\kappa$, tail law (Thm 1)	Proved	Complete analytic proof; Fig. 1
Linear reduction $\Gamma^* = 1/(1 + \rho)$ at $\beta = 0$	Exact (continuous time)	Two-state master equation; lattice agrees up to the discrete-time correction, Fig. 2, App.

		A
Bistability and saddle-nodes of the reduction (Prop. 2); threshold $\beta_c(h)$, none for $h \leq 1$	Exact for the mean-field ODE	Analytic; Fig. 3
Bistability on the nearest-neighbour lattice	Not realized at the parameters studied	Zero quasi-static loop area, converging starts, N-independent meeting times; Fig. 4, Fig. A1
Bistability with global sensing	Strong numerical evidence, $L \leq 96$	Finite slow-sweep loop; two branches, no switch in 15 000 steps at any size; Figs. 5, A2
Bistability with 80-site sensing	Discontinuous transition with metastability	Split branches mid-window; nucleation-limited decay at the edges, faster at larger L; Figs. 5, A2
Non-equilibrium steady state ($J_{\text{kill}} = J_{\text{rep}}$)	Validated numerically	Relative current imbalance below 10^{-3} ; Fig. 4D
BKT-driven field-free transition, $f_{\text{KT}} \approx 0.205 < 1 - p_c$	Numerical estimate (FSS, $L = 16-96$)	Helicity modulus vs $2T/\pi$, two extrapolations and their sensitivity, Binder cumulant; Fig. 6
Maintenance current $J(p)$, non-monotonic	Measured	Fig. 7B, Fig. 4D
Entropy production $\sigma_{\text{aux}} > 0$	Illustrative, auxiliary model	Depends on the added reverse rate ε ; Fig. 7A

373

374 The framework rests on three explicitly identified approximations. (1) The *annealed mean-field closure* δ
375 $\doteq I$ underlies Proposition 2. It is now tested directly (Section 4.4): it fails for nearest-neighbour sensing,
376 where it misstates the effective feedback in both directions, and becomes accurate as the sensing range
377 grows; whether it becomes exact in a limit in which the range grows with N is open problem (a). (2) The
378 *slaving* $U_\chi \doteq p$ in the $\beta = 0$ baseline holds up to a boundary-layer correction controlled by J_0 and h_0 . (3)
379 The entropy production of Section 6 belongs to an *auxiliary cycle* with reverse rates ε that the simulated
380 dynamics does not have; it is illustrative, and the strictly irreversible limit is singular. No other claim
381 relies on an unstated approximation.

382 9. Open problems

383 (a) *Range of the feedback and the mean-field limit*: establish rigorously the scaling limit of the coupled
384 rotor/integrity process and the conditions under which the nonlinear mean-field ODE, including its
385 bistability, is exact, for instance in a Kac-type limit in which the sensing range grows with N [18]; locate
386 the critical sensing range as a function of (β, h, K, T) ; and determine whether any region of nearest-
387 neighbour parameters, such as much larger β or smaller K, is bistable.

388 (b) *Metastability and first passage*: for finite sensing ranges, determine whether an interval of coexistence
389 survives $N \rightarrow \infty$ or shrinks to the single value of p at which an interface between the phases stalls, as the
390 nucleation-limited decay at the window edges (Section 4.4) suggests; and characterize the distribution of
391 collapse times and the large-deviation rate for noise-induced transitions, including how the lifetimes scale
392 with N and with the sensing range.

(c) *Thermodynamics and the BKT point*: a quantitative cost of maintenance requires a thermodynamically consistent model of the degradation and repair steps, with microscopic reverse rates, in place of the auxiliary cycle of Section 6, after which the fluctuation theorems for the entropy production can be tested across the bifurcation; and the Nelson–Kosterlitz analysis here ($L \leq 96$) gives $f_{KT} \approx 0.205$, which larger sizes, more realizations and temperature scans along the dilution line should refine.

10. Conclusion

Making degradation depend on the order it destroys turns a passive decay problem into a genuine non-equilibrium one. In mean field the maintained lattice is governed by a single ratio ρ through a nonlinear law that is bistable above a computed feedback threshold. On the lattice the same feedback is bistable only if degradation senses disorder over an extended region, robustly when it senses the global order and as a discontinuous transition with long-lived metastability for a finite range: with nearest-neighbour sensing the broad distribution of local disorder averages the switch away, the steady state is unique, and the hysteresis seen at finite sweep rates is dynamic and vanishes in the quasi-static limit. The range of the feedback thus joins its gain and cooperativity as a control parameter, and the loop area extrapolated to zero sweep rate is the observable that distinguishes the two regimes. The field-free loss of order is BKT-driven, at an estimated $f_{KT} \approx 0.205$, far below percolation. The maintenance current peaks where order is most fragile, and any thermodynamically consistent completion of the dynamics assigns it a positive dissipation, though its size is not fixed by the model; the exact relation $dS/dU_\chi = -\kappa$ ties directional order to configurational information. The framework is substrate-free, packaging a family of maintenance-versus-decay problems into one reduced, dissipative law together with a lattice criterion for when that law holds.

Author contributions (CRediT)

Leon Sandler: Conceptualization, Methodology, Software, Formal analysis, Writing – original draft, Writing – review & editing.

Declaration of competing interest

The author declares that he has no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial or not-for-profit sectors.

423 Declaration of generative AI and AI-assisted technologies in the writing process

424 During the preparation of this revision the author used Claude (Anthropic) to assist with typesetting the
425 equations in the journal's format, writing and testing the Monte Carlo and analysis code of Sections 4.3–5
426 and Appendix A to the author's specification, checking the reference list against Crossref records and
427 language editing. After using this tool, the author reviewed and edited the content as needed and takes full
428 responsibility for the content of the publication.

429 Data and code availability

430 All Monte Carlo results of this version, the engine, its verification suite, the analysis scripts that turn raw
431 runs into every number of Sections 4–6 and Appendix A, the raw run outputs and the scripts for every
432 figure are openly available at <https://github.com/sandlerleon/rotational-lattice-maintenance>, archived at
433 <https://doi.org/10.5281/zenodo.23131432>. Each run is reproducible on its own from its seed code
434 (Appendix A). Every figure, including the analytic Figures 1 and 3, is generated by scripts in that
435 repository (`figures_theory.py` and `analyze_study.py`). The engine reproduces exactly the update rules of
436 the open-source MyUncle framework (<https://github.com/sandlerleon/MyUncle>,
437 <https://doi.org/10.5281/zenodo.21223569>) used for earlier versions. This manuscript is archived as a
438 preprint at <https://doi.org/10.5281/zenodo.21210708>. No experimental data were used.

439 Appendix A. Monte Carlo protocol and error analysis

440 **A.1 Engine and verification.** One step consists of two checkerboard Metropolis sweeps of the rotors at
441 fixed integrity, with proposals uniform on $(-\pi, \pi]$, followed by one synchronous site update: each intact
442 site is diluted with probability λ_i , then each diluted site is repaired with probability R and its rotor reset to
443 $\theta = 0$. One step is the unit of time throughout. The engine restates the update rules of the MyUncle
444 framework used for earlier versions of this work and, before any result was computed, passed the checks
445 of its test suite: it reproduces MyUncle trajectories bit for bit for the same seed at two parameter sets; at β
446 $= 0$ the intact fraction relaxes to the discrete two-state fixed point $R/(\lambda_0 + R - \lambda_0 R)$ (0.7606 measured
447 against 0.7614 exact, and 0.7500 for the continuous-time value); $R = 0$ never repairs and $\lambda_0 = 0$ never
448 kills; the helicity modulus of the ordered undiluted lattice equals 1; and the block and global sensing
449 options of Section 4.4 average exactly the intended sites.

450 **A.2 Seeding and error estimates.** Every run is seeded from a NumPy SeedSequence of a master seed
451 (20261003) and a code naming its study, parameter set, size, control value, replicate and initial state, so
452 runs are statistically independent and each can be repeated alone. Interval estimates are 95% percentile

intervals from 2000 bootstrap resamples over independent runs or disorder realizations [31], so that variation between runs, not correlation within a run, sets the error.

A.3 Equilibration and autocorrelation. Integrated autocorrelation times are estimated with the automatic window $W \geq 6\tau_{\text{int}}$ [32]. In the stationary coupled lattice $\tau_{\text{int}}(U_\chi) = 6\text{--}11$ steps and τ_{int} of the intact fraction 9–16 steps for $L = 16\text{--}64$ and $\rho = 4.5\text{--}8$ (Figure A1A, B), with no growth with L . The 2000-step discard of the stationary runs is therefore more than $150\tau_{\text{int}}$, and the second-half averages of the 3000-step two-start runs begin after 1500 steps, more than $100\tau_{\text{int}}$. In the quenched runs of Section 5, 2000 equilibration sweeps precede 4000 measurement sweeps; the longest τ_{int} of Y at any size and dilution is 69 sweeps, so each realization contributes at least 29 effectively independent samples, and the error of each point is in any case taken from the spread over the eight disorder realizations.

A.4 Run table. Table A1 lists every study.

Table A1. Monte Carlo runs. Times are in steps (coupled lattice) or sweeps (quenched lattice); “runs” are independent replicates per parameter combination.

Study (section)	Sizes L	Control values	Runs	Length; discard; sampling
Sweep rate (4.3)	16, 24, 32, 48, 64; colder variant 24, 48	$\rho = 16 \rightarrow 1 \rightarrow 16$, 31 values; dwell $\tau = 1\text{--}256$ (9 values)	8	200-step settle at $\rho = 16$, then 62τ ; U_χ at the end of each dwell
Two starts (4.3)	16, 24, 32, 48, 64	$\rho = 4.5, 6, 8, 10$; both parameter sets	8 per start	3000; first half; every 5
Stationary (4.3, A.3)	16, 32, 64	$\rho = 4.5, 6, 8$	4	12 000; 2000; every step
Closure (4.4)	64	$\rho = 1\text{--}16$ (10 values); both parameter sets and starts	4	800 + 1200; 800; every 4
Sensing range (4.4)	48	block of 8, 24, 80 sites, global; sweeps $\tau = 64, 256$; two starts at $\rho = 3\text{--}12$	4	sweeps as above; two starts 2000, first half, every 5
Quenched dilution (5)	16, 24, 32, 48, 64, 96	$f = 0.10\text{--}0.35$ (8 values), $T = 0.45$, $h_0 = 0$	8 realizations	2000 + 4000 sweeps; 2000; every 4




Figure A1. Monte Carlo diagnostics. (A) Stationary autocorrelation of U_χ at $L = 64$ for three values of ρ , with the integrated autocorrelation time. (B) $\tau_{\text{int}}(U_\chi)$ against L : no critical slowing down. (C) Time for the seed-averaged

trajectories from the ordered and the collapsed start to agree within 0.02 (20-step moving average), against the number of sites, for both parameter sets and four values of ρ inside the mean-field window: it does not grow with N .




Figure A2. Extended-range bistability in size and time (4 runs per start, 15000 steps). (A, B) Order in the second half of each run from the ordered (filled) and the collapsed (open) start, for the 80-site neighbourhood and for global sensing, at $L = 32-96$ (points offset for visibility). (C) Ordered starts of the 80-site case at $\rho = 5.0$ decay to the collapsed branch, sooner at $L = 96$ than at $L = 32$. (D) Collapsed starts at $\rho = 6.5$ recover. Dotted: the threshold $U_X = 0.6$ used to detect a switch.

References

- [1] M.C. Cross, P.C. Hohenberg, Pattern formation outside of equilibrium, *Rev. Mod. Phys.* 65 (1993) 851–1112. <https://doi.org/10.1103/RevModPhys.65.851>
- [2] M.C. Marchetti, J.F. Joanny, S. Ramaswamy, T.B. Liverpool, J. Prost, M. Rao, et al., Hydrodynamics of soft active matter, *Rev. Mod. Phys.* 85 (2013) 1143–1189. <https://doi.org/10.1103/RevModPhys.85.1143>
- [3] S. Ramaswamy, The mechanics and statistics of active matter, *Annu. Rev. Condens. Matter Phys.* 1 (2010) 323–345. <https://doi.org/10.1146/annurev-conmatphys-070909-104101>
- [4] F. Schlögl, Chemical reaction models for non-equilibrium phase transitions, *Z. Phys.* 253 (1972) 147–161. <https://doi.org/10.1007/BF01379769>

- [5] J.E. Ferrell, Self-perpetuating states in signal transduction: positive feedback, double-negative feedback and bistability, *Curr. Opin. Cell Biol.* 14 (2002) 140–148. [https://doi.org/10.1016/S0955-0674\(02\)00314-9](https://doi.org/10.1016/S0955-0674(02)00314-9)
- [6] H. Hinrichsen, Non-equilibrium critical phenomena and phase transitions into absorbing states, *Adv. Phys.* 49 (2000) 815–958. <https://doi.org/10.1080/00018730050198152>
- [7] B. Blaiszik, S. Kramer, S. Olugebefola, J. Moore, N. Sottos, S. White, Self-healing polymers and composites, *Annu. Rev. Mater. Res.* 40 (2010) 179–211. <https://doi.org/10.1146/annurev-matsci-070909-104532>
- [8] W.F. Brown, Thermal fluctuations of a single-domain particle, *Phys. Rev.* 130 (1963) 1677–1686. <https://doi.org/10.1103/PhysRev.130.1677>
- [9] L. Sandler, A chiral-order glycation channel for synthetic cells, companion preprint, Zenodo, 2026.
- [10] T. Vicsek, A. Czirók, E. Ben-Jacob, I. Cohen, O. Shochet, Novel type of phase transition in a system of self-driven particles, *Phys. Rev. Lett.* 75 (1995) 1226–1229. <https://doi.org/10.1103/PhysRevLett.75.1226>
- [11] J. Toner, Y. Tu, Long-range order in a two-dimensional dynamical XY model: how birds fly together, *Phys. Rev. Lett.* 75 (1995) 4326–4329. <https://doi.org/10.1103/PhysRevLett.75.4326>
- [12] J.M. Kosterlitz, D.J. Thouless, Ordering, metastability and phase transitions in two-dimensional systems, *J. Phys. C: Solid State Phys.* 6 (1973) 1181–1203. <https://doi.org/10.1088/0022-3719/6/7/010>
- [13] R. Landauer, Irreversibility and heat generation in the computing process, *IBM J. Res. Dev.* 5 (1961) 183–191. <https://doi.org/10.1147/rd.53.0183>
- [14] J.M.R. Parrondo, J.M. Horowitz, T. Sagawa, Thermodynamics of information, *Nat. Phys.* 11 (2015) 131–139. <https://doi.org/10.1038/nphys3230>
- [15] J.J. Hopfield, Kinetic proofreading: a new mechanism for reducing errors in biosynthetic processes requiring high specificity, *Proc. Natl. Acad. Sci. USA* 71 (1974) 4135–4139. <https://doi.org/10.1073/pnas.71.10.4135>
- [16] U. Seifert, Stochastic thermodynamics, fluctuation theorems and molecular machines, *Rep. Prog. Phys.* 75 (2012) 126001. <https://doi.org/10.1088/0034-4885/75/12/126001>
- [17] F.S. Gnesotto, F. Mura, J. Gladrow, C.P. Broedersz, Broken detailed balance and non-equilibrium dynamics in living systems: a review, *Rep. Prog. Phys.* 81 (2018) 066601. <https://doi.org/10.1088/1361-6633/aab3ed>
- [18] T.M. Liggett, *Interacting Particle Systems*, Springer, New York, 1985.
- [19] E.T. Jaynes, Information theory and statistical mechanics, *Phys. Rev.* 106 (1957) 620–630. <https://doi.org/10.1103/PhysRev.106.620>
- [20] K.V. Mardia, P.E. Jupp, *Directional Statistics*, Wiley, Chichester, 2000.
- [21] D.E. Amos, Computation of modified Bessel functions and their ratios, *Math. Comput.* 28 (1974) 239–251. <https://doi.org/10.1090/S0025-5718-1974-0333287-7>
- [22] L. Sandler, Order–repair model: repair fidelity as a control parameter for tissue stability, software and manuscript, Zenodo, 2026. <https://doi.org/10.5281/zenodo.22851970>.
- [23] M. Rao, H.R. Krishnamurthy, R. Pandit, Magnetic hysteresis in two model spin systems, *Phys. Rev. B* 42 (1990) 856–884. <https://doi.org/10.1103/PhysRevB.42.856>
- [24] A. Bortz, M. Kalos, J. Lebowitz, A new algorithm for Monte Carlo simulation of Ising spin systems, *J. Comput. Phys.* 17 (1975) 10–18. [https://doi.org/10.1016/0021-9991\(75\)90060-1](https://doi.org/10.1016/0021-9991(75)90060-1)
- [25] R. Pfister, K.A. Schwarz, M. Janczyk, R. Dale, J.B. Freeman, Good things peak in pairs: a note on the bimodality coefficient, *Front. Psychol.* 4 (2013) 700. <https://doi.org/10.3389/fpsyg.2013.00700>
- [26] D.R. Nelson, J.M. Kosterlitz, Universal jump in the superfluid density of two-dimensional superfluids, *Phys. Rev. Lett.* 39 (1977) 1201–1205. <https://doi.org/10.1103/PhysRevLett.39.1201>
- [27] H. Weber, P. Minnhagen, Monte Carlo determination of the critical temperature for the two-dimensional XY model, *Phys. Rev. B* 37 (1988) 5986–5989. <https://doi.org/10.1103/PhysRevB.37.5986>

- 530 [28] D. Stauffer, A. Aharony, Introduction to Percolation Theory, second ed., Taylor & Francis, London, 1994.
- 531 [29] M. Hasenbusch, The two-dimensional XY model at the transition temperature: a high-precision Monte Carlo
- 532 study, J. Phys. A: Math. Gen. 38 (2005) 5869–5883. <https://doi.org/10.1088/0305-4470/38/26/003>
- 533 [30] J. Schnakenberg, Network theory of microscopic and macroscopic behavior of master equation systems, Rev.
- 534 Mod. Phys. 48 (1976) 571–585. <https://doi.org/10.1103/RevModPhys.48.571>
- 535 [31] B. Efron, Bootstrap methods: another look at the jackknife, Ann. Stat. 7 (1979) 1–26.
- 536 <https://doi.org/10.1214/aos/1176344552>
- 537 [32] N. Madras, A.D. Sokal, The pivot algorithm: a highly efficient Monte Carlo method for the self-avoiding walk,
- 538 J. Stat. Phys. 50 (1988) 109–186. <https://doi.org/10.1007/BF01022990>