

# MOSAIC<sub>bioacc</sub> REPORT

2021-04-15

---

This report is provided by the MOSAIC<sub>bioacc</sub> application available here:  
<https://mosaic.univ-lyon1.fr/bioacc>

Contact: sandrine.charles@univ-lyon1.fr

MOSAIC<sub>bioacc</sub> uses the JAGS (version 4.3.0) and R (version 4.0.2) software, and in particular packages RJags (version 4.10), jagsUI (version 1.5.1) and Shiny (version 1.6.0).

The MOSAIC<sub>bioacc</sub> application is a turn-key web tool providing bioaccumulation factors (BCF/BSAF/BMF) from a toxicokinetic (TK) model fitted to accumulation-depuration data. It is designed to fulfil the requirements of regulators when examining applications for market authorization of active substances.

---

## Data summary

File used: Oncorhynchus\_tetramethyl-benzene\_14d\_Lo2015.csv

Exposure:  $120 \mu g.g^{-1}$

Accumulation phase duration: 14 days

Number of replicates: 4

Times: 0, 2, 5, 9, 14, 14.5, 15, 17, 19, 23

Exposure routes: food

Elimination routes: excretion

## Bayesian inference

Three MCMC chains were used to estimate model parameters.

Number of iterations: 209776

Thin: 56

## TK Model

The TK model used for these calculations was:

$$\frac{dC_p(t)}{dt} = k_{uf} \times c_f - (k_{ee}) \times C_p(t) \quad \text{for } 0 \leq t \leq t_c$$

$$\frac{dC_p(t)}{dt} = - (k_{ee}) \times C_p(t) \quad \text{for } t > t_c$$

with:

$t$ : time (expressed in days )

$t_c$ : duration of the accumulation phase (expressed in days )

$C_p(t)$ : internal concentration of the parent compound at time (expressed in  $\mu g.g^{-1}$ )

$k_{ee}$ : elimination rates of excretion (expressed per days  $^{-1}$ )

$c_f$ : exposure concentration of food route (expressed in  $\mu g.g^{-1}$ )

$k_{uf}$ : uptake rate of food exposure (expressed per days  $^{-1}$ )


## Bioaccumulation factor calculation

### Calculations

$$BMF_k = \frac{k_{uf}}{k_{ee}}$$

$$BMF_{ss} = \frac{C_p(t_c)}{c_f}$$

### BMF<sub>k</sub> plot




### BMF summary

	2.5%	50%	97.5%	CV
BMFk	5.8e-03	7e-03	9e-03	0.11

## Fitting results

### Fit plot



### Quantiles of estimated parameters


	2.5%	50%	97.5%	
$k_{uf}$	4.668e-04	7.253e-04	1.046e-03	$d^{-1}$
$k_{ee}$	0.0439	0.1029	0.1693	$d^{-1}$
$\sigma_p$	0.09863	0.131	0.1887	$\mu g.g^{-1}$

## Goodness-of-fit criteria

### Posterior Predictive Check


The PPC shows the observed values against their corresponding estimated predictions (black dots), along with their 95% credible interval (vertical segments). If the fit is correct, we expect to see 95% of the data within the intervals. Ideally observations and predictions should coincide, so we would expect to see black dots along the first bisector  $y = x$  (plain black line). The 95% credible intervals are colored in green if they

overlap this line, in red otherwise.



### Priors and posteriors


The prior distribution is represented by the gray area and the posterior distribution by the orange area. The accuracy of the model parameter estimation can be visualized by comparing prior and posterior distributions: the overall expectation is to get a narrower posterior distribution compared to the prior one, what reflects that data contributed enough to precisely estimate parameters.



### Correlation between parameters

If you want to see the coloured matrix giving a summary of parameter correlations, you need to import the corresponding figure directly from the application, page bottom, section “Downloads”, then choose Download an output and select “GOF” then “parameter correlation”. You can select the output format you prefer.

Correlations between parameters are visualized by projecting the joint posterior distribution in a plot matrix with planes of parameter pairs (lower triangular elements), marginal posterior distribution of each model parameter (diagonal), and Pearson correlation coefficients (upper triangular elements). Correlations are expected to be low (reflected by “potatoid” shapes of density lines in orange); a leaning elliptical shape translates high correlations (positive if leaning to the right, negative if leaning to the left).



### Potential Scale Reduction Factors

Convergence of the MCMC chains can be checked with the Gelman-Rubin diagnostic expressed with the potential scale reduction factor (PSRF). Approximate convergence is diagnosed when the PSRF is below 1.01.

	PSRF
kuf	1.001
kee	1.001
sigmaCpred	1

### **Watanabe–Akaike information criterion**

Information criteria offer a computationally appealing way of estimating the generalization performance of the model. A fully Bayesian criterion is the widely applicable information criterion (WAIC) by Watanabe a penalized deviance statistics accounting for the uncertainty in the parameters and can be used also for singular models. WAIC is widely used in model comparison for a same dataset (e.g., with or without  $k_{ee}$ ). Sub-models with lower WAIC values will be preferred.

$$\text{WAIC} = -25.8$$


### **Deviance Information Criterion**

This criteria, denoted DIC, is a penalized deviance statistics accounting for the number of parameters for use in model comparison for a same dataset (e.g., with or without  $k_{ee}$ ). Sub-models with lower DIC values will be preferred.


$$\text{DIC} = -23.68$$

### **Traces of MCMC iterations**

A traceplot is an essential plot for assessing convergence and diagnosing of MCMC chains. It shows the time series of the sampling process leading to the posterior distribution. Different colors are used for each of the chains (here 3) to assess within-chain convergence.







## Data Table

time	conc	expf	replicate
0.0	0.0000	120	1
2.0	0.2691	120	1
2.0	0.2954	120	2
2.0	0.4370	120	3
5.0	0.4596	120	1
5.0	0.4805	120	2
5.0	0.5015	120	3
9.0	0.5171	120	1
9.0	0.5258	120	2
9.0	0.5870	120	3
14.0	0.5151	120	1
14.0	0.5553	120	2
14.5	0.5151	120	1
14.5	0.5343	120	2
14.5	0.5711	120	4
15.0	0.5081	120	1
15.0	0.5413	120	2
17.0	0.4416	120	1
17.0	0.5097	120	2
19.0	0.2859	120	1
19.0	0.3838	120	2
19.0	0.4729	120	3
23.0	0.4640	120	1
23.0	0.4570	120	2