



MOSAIC_{bioacc} REPORT

2021-06-03

This report is provided by the $MOSAIC_{bioacc}$ application available here: https://mosaic.univ-lyon1.fr/bioacc

 $Contact: \ sandrine.charles@univ-lyon1.fr$

MOSAIC_{bioacc} uses the JAGS (version 4.3.0) and R (version 4.0.2) software, and in particular packages RJags (version 4.10), jagsUI (version 1.5.1) and Shiny (version 1.6.0).

The MOSAIC_{bioacc} application is a turn-key web tool providing bioaccumulation factors (BCF/BSAF/BMF) from a toxicokinetic (TK) model fitted to accumulation-depuration data. It is designed to fulfil the requirements of regulators when examining applications for market authorization of active substances.

Data summary

File used: Daphnia pyrene 24h Wang2021 body-without-gut-food.txt

Exposure: $37.884 \ \mu q.q^{-1}$

Accumulation phase duration: 24 hours

Number of replicates: 3

Times: 0, 0.5, 2, 5, 10, 24, 26, 29, 34, 48, 72

Exposure routes: food

Elimination routes: excretion

Bayesian inference

Three MCMC chains were used to estimate model parameters.

Number of iterations: 33714

Thin: 9





TK Model

The TK model used for these calculations was:

$$\frac{dC_p(t)}{dt} = k_{uf} \times c_f - (k_{ee}) \times C_p(t) \quad for \ 0 \le t \le t_c$$

$$\frac{dC_p(t)}{dt} = -(k_{ee}) \times C_p(t) \quad for \ t > t_c$$

with:

t: time (expressed in hours)

 t_c : duration of the accumulation phase (expressed in hours)

 $C_p(t)$: internal concentration of the parent compound at time (expressed in $\mu g.g^{-1}$)

 k_{ee} : elimination rates of excretion (expressed per hours $^{-1}$)

 c_f : exposure concentration of food route (expressed in $\mu g.g^{-1}$)

 k_{uf} : uptake rate of food exposure (expressed per hours $^{-1}$)


Bioaccumulation factor calculation

Calculations

$$BMF_k = \frac{k_{uf}}{k_{ee}}$$

$$BMF_{ss} = \frac{C_p(t_c)}{c_f}$$

 BMF_k plot



BMF summary






	2.5%	50%	97.5%	CV
BMFk	0.011	0.012	0.013	0.042

Fitting results

Fit plot



Quantiles of estimated parameters

	2.5%	50%	97.5%	
$\overline{k_{uf}}$	0.00247	0.003039	0.003731	h^{-1}
		0.2524		h^{-1}
σ_p	0.03514	0.04604	0.06361	$\mu g.g^{-1}$

Goodness-of-fit criteria

Posterior Predictive Check

The PPC shows the observed values against their corresponding estimated predictions (black dots), along with their 95% credible interval (vertical segments). If the fit is correct, we expect to see 95% of the data within the intervals. Ideally observations and predictions should coincide, so we would expect to see black dots along the first bisector y = x (plain black line). The 95% credible intervals are colored in green if they






overlap this line, in red otherwise.

Parent compound:

percentage of data in CI:




Priors and posteriors

The prior distribution is represented by the gray area and the posterior distribution by the orange area. The accuracy of the model parameter estimation can be visualized by comparing prior and posterior distributions: the overall expectation is to get a narrower posterior distribution compared to the prior one, what reflects that data contributed enough to precisely estimate parameters.








Correlation between parameters

If you want to see the coloured matrix giving a summary of parameter correlations, you need to import the corresponding figure directly from the application, page bottom, section "Downloads", then choose Download an output and select "GOF" then "parameter correlation". You can select the output format you prefer.

Correlations between parameters are visualized by projecting the joint posterior distribution in a plot matrix with planes of parameter pairs (lower triangular elements), marginal posterior distribution of each model parameter (diagonal), and Pearson correlation coefficients (upper triangular elements). Correlations are expected to be low (reflected by "potatoid" shapes of density lines in orange); a leaning elliptical shape translates high correlations (positive if leaning to the right, negative if leaning to the left).







Potential Scale Reduction Factors

Convergence of the MCMC chains can be check with the Gelman-Rubin diagnostic expressed with the potential scale reduction factor (PSRF). Approximate convergence is diagnosed when the PSRF is below 1.01.

	PSRF
kuf	1.000937
kee	1.001049
sigmaCpred	1.000117





Watanabe-Akaike information criterion

Information criteria offer a computationally appealing way of estimating the generalization performance of the model. A fully Bayesian criterion is the widely applicable information criterion (WAIC) by Watanabe a penalized deviance statistics accounting for the uncertainty in the parameters and can be used also for singular models. WAIC is widely used in model comparison for a same dataset (e.g., with or without $k_{\rm ee}$). Sub-models with lower WAIC values will be preferred.

WAIC = -80.2

Deviance Information Criterion

This criteria, denoted DIC, is a penalized deviance statistics accounting for the number of parameters for use in model comparison for a same dataset (e.g., with or without $k_{\rm ee}$). Sub-models with lower DIC values will be preferred.


DIC = -79.79249

Traces of MCMC iterations

A traceplot is an essential plot for assessing convergence and diagnosing of MCMC chains. It shows the time series of the sampling process leading to the posterior distribution. Different colors are used for each of the chains (here 3) to assess within-chain convergence.


















Data Table

time	conc	$\exp f$	replicate
0.0	0.000000000	37.884	1
0.5	0.088343558	37.884	1
0.5	0.106748466	37.884	2
0.5	0.117791411	37.884	3
2.0	0.169325153	37.884	1
2.0	0.220858896	37.884	2
5.0	0.265030675	37.884	1
5.0	0.382822086	37.884	2
10.0	0.412269939	37.884	1
10.0	0.338650307	37.884	2
24.0	0.552147239	37.884	1
24.0	0.434355828	37.884	2
26.0	0.316564417	37.884	1
26.0	0.246625767	37.884	2
26.0	0.220858896	37.884	3
29.0	0.169325153	37.884	1
29.0	0.117791411	37.884	2
29.0	0.106748466	37.884	3
34.0	0.073619632	37.884	1
34.0	0.058895706	37.884	2
34.0	0.051533742	37.884	3
48.0	0.040490798	37.884	1
48.0	0.022085890	37.884	2
48.0	0.007361963	37.884	3
72.0	0.014723926	37.884	1
72.0	0.007361963	37.884	2