Euclid's Elements

Book VII

Definitions:

- A unit is that by virtue of which each of the things that exist is called one
- 2 A number is a multitude composed of units. (not one)
- A number is part of a number, the less of the greater, when it measures the greater
- 11 A prime number is that which is measured by a unit alone.
- 12 Numbers prime to one another are those which are measured by a unit alone as a common measure
- A number is said to multiply a number when that which is multiplied is added to itself as many times as there are units in the other, and thus some number is produced.
- Numbers are proportional when the first is the same multiple, or the same part, or the same parts, of the second that the third is of the fourth.

As long as algebra and geometry have been separated, their progress have been slow and their uses limited; but when these two sciences have been united, they have lent each mutual forces, and have marched together towards perfection.

Joseph-Louis Lagrange (1736 to 1813)



Table of Contents, Chapter 7

- Determine if two numbers are relatively prime
- 2 Find the greatest common divisor for two numbers
- 3 Find the largest common divisor for three numbers
- 4 Given two natural numbers, A and B, either B is part of A, or there exists a natural number (a part) that can measure both A and B
- 5 If B = $(1/q)\cdot A$ and D = $(1/q)\cdot C$, then $(B+D) = (1/q)\cdot (A+C)$
- 6 If B = $(p/q)\cdot A$ and D = $(p/q)\cdot C$, then $(B+D) = (p/q)\cdot (A+C)$
- 7 If B = A/q and D = C/q, B>D, then (B-D) = (A-C)/q
- 8 If B = $(p/q)\cdot A$ and D = $(p/q)\cdot C$, B>D, then $(B-D) = (p/q)\cdot (A-C)$
- 9 If B = (1/q)·A and D = (1/q)·C, and If B = (r/s)·D, then A = (r/s)·C

- 10 If B = $(p/q)\cdot A$ and D = $(p/q)\cdot C$, and If B = $(r/s)\cdot D$, then A = $(r/s)\cdot C$
- 11 If A:B = C:D, then (A-C):(B-D) = A:B
- 12 If A:B = C:D, then (A+C):(B+C) = A:B
- 13 If A:B = C:D, then A:C = B:D
- 14 If A:B = D:E and B:C = E:F, then A:C = D:F
- 15 If B = i·1 and E = i·D, and if D = j·1 then E = j·B
- 16 $A \times B = B \times A$
- 17 If D = A × B and E = A × C then D:E = B:C
- 18 If D = B × A and E = C × A then D:E = B:C
- 19 If A:B = C:D then $A \times D = B \times C$ If $A \times D = B \times C$ then A:B = C:D
- 20 Given the ratio A:B and C,D are the smallest numbers such that A:B = C:D then A = n·C and B = n·D

- If A,B are relatively prime, then A,B are the smallest whole numbers that can be used to describe the ratio A:B
- 22 If A,B are the smallest whole numbers that can be used to describe the ratio A:B, then A,B are relatively prime
- 23 If A,B are relatively prime and if A = n·C, then B,C are relatively prime
- 24 If A,C are relatively prime and B,C are relatively prime then the A × B is relatively prime to C
- 25 If A,B are relatively prime then A²,B are relatively prime
- If A is relatively prime to C and D, and if B is also relatively prime to C and D, then A × B is relatively prime to C × D
- 27 If A,B are relatively prime, then A²,B² are relatively prime, and A³,B³ are relatively prime, and so on



Table of Contents, Chapter 7

- 28 If A,B are relatively prime, then A,(A+B) are relatively prime
- 29 If A is prime, and B ≠ n·A, then A,B are relatively prime
- 30 If C = A×B and C = i·D where D is prime, then either A = j·D or B = j·D
- 31 If $A = B \times C$, then $A = j \cdot D$ where D is prime
- 32 If A is a number then it is either prime, or $A = j \cdot D$ where D is prime
- Find the smallest numbers X,Y,Z where the ratio X:Y:Z is equal to the given ratio A:B:C
- 34 Find the lowest common denominator of 2 numbers
- 35 If E is the lowest common denominator of A,B, and if C = n ·A = m·B, then C = i·E
- 36 Find the least common multiple of 3 numbers

- If $A = p \cdot B$, then $A = q \cdot C$ where $C = p \cdot 1$
- 38 If $A = (1/c) \cdot B$ and $C = c \cdot 1$ then $A = n \cdot C$
- Find the smallest number that has the fractions 1/a, 1/b, 1/c



If there be as many numbers as we please, and others equal to them in multitude, which taken two and two are in the same ratio, they will also be in the same ratio ex aequali



If there be as many numbers as we please, and others equal to them in multitude, which taken two and two are in the same ratio, they will also be in the same ratio ex aequali

								_	_		_				6
			Λ							. .	.				
			- /\												
			-							1	J				
										_	7				
												1		i	
		_							_	_					
									+	+					
)												
									1	1					
													_		
		_							_						
	+														

A:B = D:E

B:C = E:F

In other words

Given as many numbers as we please A,B,C

And an equal number of others, D,E,F

And let them be, two by two, in the same ratio (A is to B as D is to E, B is to C as E is to F)

If there be as many numbers as we please, and others equal to them in multitude, which taken two and two are in the same ratio, they will also be in the same ratio ex aequali

á							_								
- 7												1			1
			÷-\							+	3				
			÷⊢								י				
										_					
4					-							-			
		1	_						_	_					
		+	2						+						
		L	ָּ							-					
1															
									_						
		(_						

A:B = D:E

B:C = E:F

A:C = D:F

In other words

Given as many numbers as we please A,B,C

And an equal number of others, D,E,F

And let them be, two by two, in the same ratio (A is to B as D is to E, B is to C as E is to F)

Then, as A is to C, so is D to F

If there be as many numbers as we please, and others equal to them in multitude, which taken two and two are in the same ratio, they will also be in the same ratio ex aequali

			_							+ F	A			
			$\boldsymbol{\wedge}$								1			
		_							.	1				
		В												
	(<u> </u>							F					

A:B = D:E B:C = E:F

In other words

Given as many numbers as we please A,B,C

And an equal number of others, D,E,F

And let them be, two by two, in the same ratio (A is to B as D is to E, B is to C as E is to F)

Then, as A is to C, so is D to F

Proof

If there be as many numbers as we please, and others equal to them in multitude, which taken two and two are in the same ratio, they will also be in the same ratio ex aequali

В	· · · · · · · · · · · · · · · · · · ·	
C		
A:B = D:E		
A.D = D.E		
B:C=E:F		
1 D D D D		
A:D = B:E		

In other words

Given as many numbers as we please A,B,C

And an equal number of others, D,E,F

And let them be, two by two, in the same ratio (A is to B as D is to E, B is to C as E is to F)

Then, as A is to C, so is D to F

Proof

Since A is to B as D is to E, alternately, A is to D as B is to E (VII-13)

If there be as many numbers as we please, and others equal to them in multitude, which taken two and two are in the same ratio, they will also be in the same ratio ex aequali

-		 					
. i . i . i . i . i . i . i . i							
i i i Ai					ע		
 						_	
· + + + + D +-				++	.		
D					-		
-				-			
				ļ 			
A:B =	り:上						
B:C =	- Eith						
+++							
A:D=	R:F						
BIF =	C · F						

In other words

Given as many numbers as we please A,B,C

And an equal number of others, D,E,F

And let them be, two by two, in the same ratio (A is to B as D is to E, B is to C as E is to F)

Then, as A is to C, so is D to F

Proof

Since A is to B as D is to E, alternately, A is to D as B is to E (VII-13)

And since B is to C as E is to F, alternately, B is to E as C is to F (VII-13)

If there be as many numbers as we please, and others equal to them in multitude, which taken two and two are in the same ratio, they will also be in the same ratio ex aequali

B		E	
· · · · · · · · · · · · · · · · · · ·			
- + + + + + + + +		··· - - - - +	
A - D - D - I	-		
A:B = D:E	-		
B:C = E:I	= ;;;;		
A:D = B:E	-		
$B \cdot F = C \cdot F$	<u>-</u>		

In other words

Given as many numbers as we please A,B,C

And an equal number of others, D,E,F

And let them be, two by two, in the same ratio (A is to B as D is to E, B is to C as E is to F)

Then, as A is to C, so is D to F

Proof

Since A is to B as D is to E, alternately, A is to D as B is to E (VII-13)

And since B is to C as E is to F, alternately, B is to E as C is to F (VII-13)

The ratio of B to E is the same as A to D and C to F, consequently A to D equals C to F

A:D = C:F

If there be as many numbers as we please, and others equal to them in multitude, which taken two and two are in the same ratio, they will also be in the same ratio ex aequali

				<u> </u>	
A			D		
					
B			Ė		
			- :		
C		† F			

A:B = D:E					
B:C = E:F					
D.C - L.I					
A:D = B:E					
B:E = C:F					

A:D = C:F					
A:C = D:F					

In other words

Given as many numbers as we please A,B,C

And an equal number of others, D,E,F

And let them be, two by two, in the same ratio (A is to B as D is to E, B is to C as E is to F)

Then, as A is to C, so is D to F

Proof

Since A is to B as D is to E, alternately, A is to D as B is to E (VII-13)

And since B is to C as E is to F, alternately, B is to E as C is to F (VII-13)

The ratio of B to E is the same as A to D and C to F, consequently A to D equals C to F

Alternately, A to C is equal to D to F (VII-13)

If there be as many numbers as we please, and others equal to them in multitude, which taken two and two are in the same ratio, they will also be in the same ratio ex aequali

i i i i i i i i i i i i i i i i i i i			<u> </u>	
A			שׁיִּ	
· · · · · · · · · · · · · · · · · · ·			<u> </u>	
В			‡ ‡ † † † † † † † † † † † † † † † † † †	
	<u> </u>			
			<u> </u>	
			F	
A:B =	D:E			
B:C =	E.E.			
A:D =	B:E			
B:E =	C·F			
A:D =	C:F			
A:D =	C:F			

In other words

Given as many numbers as we please A,B,C

And an equal number of others, D,E,F

And let them be, two by two, in the same ratio (A is to B as D is to E, B is to C as E is to F)

Then, as A is to C, so is D to F

Proof

Since A is to B as D is to E, alternately, A is to D as B is to E (VII-13)

And since B is to C as E is to F, alternately, B is to E as C is to F (VII-13)

The ratio of B to E is the same as A to D and C to F, consequently A to D equals C to F

Alternately, A to C is equal to D to F (VII-13)

A:C = D:F

Youtube Videos

https://www.youtube.com/c/SandyBultena











Except where otherwise noted, this work is licensed under http://creativecommons.org/licenses/by-nc/3.0