

# Euclid's Elements

## Book VII

### Definitions:

- 1 A unit is that by virtue of which each of the things that exist is called one
- 2 A number is a multitude composed of units. (not one)
- 3 A number is part of a number, the less of the greater, when it measures the greater
- 11 A prime number is that which is measured by a unit alone.
- 12 Numbers prime to one another are those which are measured by a unit alone as a common measure
- 15 A number is said to multiply a number when that which is multiplied is added to itself as many times as there are units in the other, and thus some number is produced.
- 20 Numbers are proportional when the first is the same multiple, or the same part, or the same parts, of the second that the third is of the fourth.

*As long as algebra and geometry have been separated, their progress have been slow and their uses limited; but when these two sciences have been united, they have lent each mutual forces, and have marched together towards perfection.*

**Joseph-Louis Lagrange**  
**(1736 to 1813)**



# Table of Contents, Chapter 7

1	Determine if two numbers are relatively prime	10	If $B = (p/q) \cdot A$ and $D = (p/q) \cdot C$ , and If $B = (r/s) \cdot D$ , then $A = (r/s) \cdot C$	21	If A,B are relatively prime, then A,B are the smallest whole numbers that can be used to describe the ratio A:B
2	Find the greatest common divisor for two numbers	<b>11</b>	<b>If <math>A:B = C:D</math>, then <math>(A-C):(B-D) = A:B</math></b>	22	If A,B are the smallest whole numbers that can be used to describe the ratio A:B, then A,B are relatively prime
3	Find the largest common divisor for three numbers	12	If $A:B = C:D$ , then $(A+C):(B+C) = A:B$	23	If A,B are relatively prime and if $A = n \cdot C$ , then B,C are relatively prime
4	Given two natural numbers, A and B, either B is part of A, or there exists a natural number (a part) that can measure both A and B	13	If $A:B = C:D$ , then $A:C = B:D$	24	If A,C are relatively prime and B,C are relatively prime then the $A \times B$ is relatively prime to C
5	If $B = (1/q) \cdot A$ and $D = (1/q) \cdot C$ , then $(B+D) = (1/q) \cdot (A+C)$	14	If $A:B = D:E$ and $B:C = E:F$ , then $A:C = D:F$	25	If A,B are relatively prime then $A^2, B$ are relatively prime
6	If $B = (p/q) \cdot A$ and $D = (p/q) \cdot C$ , then $(B+D) = (p/q) \cdot (A+C)$	15	If $B = i \cdot 1$ and $E = i \cdot D$ , and if $D = j \cdot 1$ then $E = j \cdot B$	26	If A is relatively prime to C and D, and if B is also relatively prime to C and D, then $A \times B$ is relatively prime to $C \times D$
7	If $B = A/q$ and $D = C/q$ , $B > D$ , then $(B-D) = (A-C)/q$	16	$A \times B = B \times A$	27	If A,B are relatively prime, then $A^2, B^2$ are relatively prime, and $A^3, B^3$ are relatively prime, and so on
8	If $B = (p/q) \cdot A$ and $D = (p/q) \cdot C$ , $B > D$ , then $(B-D) = (p/q) \cdot (A-C)$	17	If $D = A \times B$ and $E = A \times C$ then $D:E = B:C$		
9	If $B = (1/q) \cdot A$ and $D = (1/q) \cdot C$ , and If $B = (r/s) \cdot D$ , then $A = (r/s) \cdot C$	18	If $D = B \times A$ and $E = C \times A$ then $D:E = B:C$		
		19	If $A:B = C:D$ then $A \times D = B \times C$ If $A \times D = B \times C$ then $A:B = C:D$		
		20	Given the ratio A:B and C,D are the smallest numbers such that $A:B = C:D$ then $A = n \cdot C$ and $B = n \cdot D$		



# Table of Contents, Chapter 7

- |    |  |    |   |
|----|--|----|---|
| 28 | If A,B are relatively prime, then A,(A+B) are relatively prime   | 37 | If $A = p \cdot B$ , then $A = q \cdot C$ where $C = p \cdot 1$ |
| 29 | If A is prime, and $B \neq n \cdot A$ , then A,B are relatively prime                                    | 38 | If $A = (1/c) \cdot B$ and $C = c \cdot 1$ then $A = n \cdot C$ |
| 30 | If $C = A \times B$ and $C = i \cdot D$ where D is prime, then either $A = j \cdot D$ or $B = j \cdot D$ | 39 | Find the smallest number that has the fractions $1/a, 1/b, 1/c$ |
| 31 | If $A = B \times C$ , then $A = j \cdot D$ where D is prime  |    |   |
| 32 | If A is a number then it is either prime, or $A = j \cdot D$ where D is prime                            |    |   |
| 33 | Find the smallest numbers X,Y,Z where the ratio X:Y:Z is equal to the given ratio A:B:C                  |    |   |
| 34 | Find the lowest common denominator of 2 numbers  |    |   |
| 35 | If E is the lowest common denominator of A,B, and if $C = n \cdot A = m \cdot B$ , then $C = i \cdot E$  |    |   |
| 36 | Find the least common multiple of 3 numbers  |    |   |



# Proposition 11 of Book VII

If, as whole is to whole, so is a number subtracted to a number subtracted, the remainder will also be to the remainder as whole to whole.



## Proposition 11 of Book VII

If, as whole is to whole, so is a number subtracted to a number subtracted, the remainder will also be to the remainder as whole to whole.

## Definition 20


Numbers are proportional when the first is the same multiple, or the same part, or the same parts, of the second that the third is of the fourth





# Proposition 11 of Book VII

If, as whole is to whole, so is a number subtracted to a number subtracted, the remainder will also be to the remainder as whole to whole.



$$AB:CD = AE:CF$$

## Definition 20

Numbers are proportional when the first is the same multiple, or the same part, or the same parts, of the second that the third is of the fourth

## In other words


Given AB and CD are in the same proportion as AE and CF

Then EB and FD are in the same proportion as AB and CD



# Proposition 11 of Book VII

If, as whole is to whole, so is a number subtracted to a number subtracted, the remainder will also be to the remainder as whole to whole.



$$AB:CD = AE:CF$$
$$(AB-AE):(CD-CF) = AB:CD$$

## Definition 20

Numbers are proportional when the first is the same multiple, or the same part, or the same parts, of the second that the third is of the fourth


## In other words

Given AB and CD are in the same proportion as AE and CF  
Then EB and FD are in the same proportion as AB and CD



# Proposition 11 of Book VII

If, as whole is to whole, so is a number subtracted to a number subtracted, the remainder will also be to the remainder as whole to whole.



$$AB:CD = AE:CF$$

## Definition 20

Numbers are proportional when the first is the same multiple, or the same part, or the same parts, of the second that the third is of the fourth

## In other words

Given AB and CD are in the same proportion as AE and CF  
Then EB and FD are in the same proportion as AB and CD


## Proof





# Proposition 11 of Book VII

If, as whole is to whole, so is a number subtracted to a number subtracted, the remainder will also be to the remainder as whole to whole.



$$\begin{aligned} AB:CD &= AE:CF \\ AB &= (p/q)CD \\ AE &= (p/q)CF \end{aligned}$$

## Definition 20

Numbers are proportional when the first is the same multiple, or the same part, or the same parts, of the second that the third is of the fourth

## In other words

Given AB and CD are in the same proportion as AE and CF  
Then EB and FD are in the same proportion as AB and CD


## Proof

By definition of proportional, whatever part or parts AB is of CD, the same part or the same parts is AE of CF (VII Def 20).



# Proposition 11 of Book VII

If, as whole is to whole, so is a number subtracted to a number subtracted, the remainder will also be to the remainder as whole to whole.



$$AB:CD = AE:CF$$

$$AB = (p/q)CD$$

$$AE = (p/q)CF$$

$$(AB-AE) = (p/q)(CD-CF)$$

$$EB = (p/q)FD$$

## Definition 20

Numbers are proportional when the first is the same multiple, or the same part, or the same parts, of the second that the third is of the fourth

## In other words

Given AB and CD are in the same proportion as AE and CF

Then EB and FD are in the same proportion as AB and CD

## Proof


By definition of proportional, whatever part or parts AB is of CD, the same part or the same parts is AE of CF (VII Def 20).

Therefore the remainder EB is the same part or parts of FD that AB is of CD (VII·7,8)



# Proposition 11 of Book VII

If, as whole is to whole, so is a number subtracted to a number subtracted, the remainder will also be to the remainder as whole to whole.



$$AB:CD = AE:CF$$

$$AB = (p/q)CD$$

$$AE = (p/q)CF$$

$$(AB - AE) = (p/q)(CD - CF)$$

$$EB = (p/q)FD$$

$$EB:FD = AB:CD$$

## Definition 20

Numbers are proportional when the first is the same multiple, or the same part, or the same parts, of the second that the third is of the fourth

## In other words

Given AB and CD are in the same proportion as AE and CF

Then EB and FD are in the same proportion as AB and CD

## Proof

By definition of proportional, whatever part or parts AB is of CD, the same part or the same parts is AE of CF (VII Def 20).


Therefore the remainder EB is the same part or parts of FD that AB is of CD (VII·7,8)

Therefore, as EB is to FD, so is AB to CD (VII Def 20)



# Proposition 11 of Book VII

If, as whole is to whole, so is a number subtracted to a number subtracted, the remainder will also be to the remainder as whole to whole.



$$AB:CD = AE:CF$$

$$AB = (p/q)CD$$

$$AE = (p/q)CF$$

$$(AB - AE) = (p/q)(CD - CF)$$

$$EB = (p/q)FD$$

$$EB:FD = AB:CD$$

## Definition 20

Numbers are proportional when the first is the same multiple, or the same part, or the same parts, of the second that the third is of the fourth

## In other words

Given AB and CD are in the same proportion as AE and CF

Then EB and FD are in the same proportion as AB and CD

## Proof

By definition of proportional, whatever part or parts AB is of CD, the same part or the same parts is AE of CF (VII Def 20).

Therefore the remainder EB is the same part or parts of FD that AB is of CD (VII·7,8)

Therefore, as EB is to FD, so is AB to CD (VII Def 20)



# Youtube Videos

<https://www.youtube.com/c/SandyBultena>

*Copyright © 2019 by Sandy Bultena.*



Except where otherwise noted, this work is licensed under  
<http://creativecommons.org/licenses/by-nc/3.0>