

Discrete Mathematics (ITPC-309)

Ordered Sets and Lattices – Part II



Dr. Sanga Chaki

Department of Information Technology

Dr. B. R. Ambedkar National Institute of Technology, Jalandhar



Contents

1. Lattices
2. Duality
3. Idempotent Law
4. Lattices and Order
5. Bounded Lattices
6. Distributive Lattices
7. Complements
8. Complemented Lattices

Lattices

1. There are two ways to define a lattice L .
 - In terms of Posets
 - In terms of axioms
2. A lattice L may be defined as a partially ordered set in which $\inf(a,b)$ and $\sup(a, b)$ exist for any pair of elements $a, b \in L$.

Lattices

1. Let L be a nonempty set closed under two binary operations called meet and join, denoted respectively by \wedge and \vee .
2. Then (L, \wedge, \vee) is called lattice if the following axioms hold where a, b, c are elements in L :

[L₁] Commutative law:

$$(1a) \quad a \wedge b = b \wedge a$$

$$(1b) \quad a \vee b = b \vee a$$

[L₂] Associative law:

$$(2a) \quad (a \wedge b) \wedge c = a \wedge (b \wedge c)$$

$$(2b) \quad (a \vee b) \vee c = a \vee (b \vee c)$$

[L₃] Absorption law:

$$(3a) \quad a \wedge (a \vee b) = a$$

$$(3b) \quad a \vee (a \wedge b) = a$$

Duality

1. The dual of any statement in a lattice (L, \wedge, \vee) is defined to be the statement that is obtained by interchanging \wedge and \vee .
2. For example, the dual of $a \wedge (b \vee a) = a \vee a$ is $a \vee (b \wedge a) = a \wedge a$
3. Notice that the dual of each axiom of a lattice is also an axiom.
4. Accordingly, the principle of duality holds: **The dual of any theorem in a lattice is also a theorem**

Idempotent Law

1. Important property of lattices following directly from the absorption laws:
the idempotent laws:

$$(i) a \wedge a = a; \quad (ii) a \vee a = a.$$

2. Proof:

$$\begin{aligned} a \wedge a &= a \wedge (a \vee (a \wedge b)) && \text{(using (3b))} \\ &= a && \text{(using (3a))} \end{aligned}$$

$$(3b) \quad a \vee (a \wedge b) = a$$

$$(3a) \quad a \wedge (a \vee b) = a$$

Lattices and Order

1. Given a lattice L , we can define a partial order on L as follows:

$$a \lesssim b \quad \text{if} \quad a \wedge b = a$$


Analogously, we could define

$$a \lesssim b \quad \text{if} \quad a \vee b = b$$

2. Now that we have a partial order on any lattice L , we can picture L by a diagram as was done for partially ordered sets in general – Hasse Diagrams

Lattices and Order - Example

1. Let C be a collection of sets closed under intersection and union.
2. Then (C, \cap, \cup) is a lattice.
3. In this lattice, the partial order relation is the same as the set inclusion relation.
4. Below figure shows the diagram of the lattice L of all subsets of $\{a, b, c\}$.



Bounded Lattices

1. A lattice L is said to have a lower bound 0 if for any element x in L we have $0 \preceq x$.
2. Analogously, L is said to have an upper bound I if for any x in L we have $x \preceq I$.
3. We say L is bounded if L has both a lower bound 0 and an upper bound I .
4. In such a lattice we have the following identities for any element a in L .

$$a \vee I = I, \quad a \wedge I = a, \quad a \vee 0 = a, \quad a \wedge 0 = 0$$

Bounded Lattices - Examples

1. Example 1: The nonnegative integers with the usual ordering, $0 < 1 < 2 < 3 < 4 < \dots$ have 0 as a lower bound but have no upper bound.
2. Example 2: The lattice $P(U)$ of all subsets of any universal set U is a bounded lattice with U as an upper bound and the empty set Φ as a lower bound.
3. Example 3: Suppose $L = \{a_1, a_2, \dots, a_n\}$ is a finite lattice. Then $a_1 \vee a_2 \vee \dots \vee a_n$ and $a_1 \wedge a_2 \wedge \dots \wedge a_n$ are upper and lower bounds for L , respectively.
4. Note: Every finite lattice L is bounded

Distributive Lattices

1. A lattice L is said to be distributive if for any elements a, b, c in L we have the following:

[L₄] Distributive law:

$$(4a) \ a \wedge (b \vee c) = (a \wedge b) \vee (a \wedge c) \qquad (4b) \ a \vee (b \wedge c) = (a \vee b) \wedge (a \vee c)$$

2. Otherwise, L is said to be non-distributive.

Complements

1. Let L be a bounded lattice with lower bound 0 and upper bound I .
2. Let a be an element of L .
3. An element x in L is called a complement of a if

$$a \vee x = I \quad \text{and} \quad a \wedge x = 0$$

4. Complements need not exist and need not be unique

Complements

1. Let L be a bounded distributive lattice.
2. Then complements are unique if they exist.

Proof: Suppose x and y are complements of any element a in L . Then

$$a \vee x = I, \quad a \vee y = I, \quad a \wedge x = 0, \quad a \wedge y = 0$$

Using distributivity,

$$x = x \vee 0 = x \vee (a \wedge y) = (x \vee a) \wedge (x \vee y) = I \wedge (x \vee y) = x \vee y$$

Similarly,

$$y = y \vee 0 = y \vee (a \wedge x) = (y \vee a) \wedge (y \vee x) = I \wedge (y \vee x) = y \vee x$$

Thus

$$x = x \vee y = y \vee x = y$$

Complemented Lattices

1. A lattice L is said to be complemented if L is bounded and every element in L has a complement.
2. The figure shows a complemented lattice where complements are not unique:
3. Find the complements of each of the elements of the below lattice

