STATISTICS WORKSHEET-3

Q1 to Q9 have only one correct answer. Choose the correct option to answer your question.

- 1. Which of the following is the correct formula for total variation?
- b) Total Variation = Residual Variation + Regression Variation
- 2. Collection of exchangeable binary outcomes for the same covariate data are called outcomes.
- c) binomial
- 3. How many outcomes are possible with Bernoulli trial?
- a) 2
- 4. If Ho is true and we reject it is called
- a) Type-I error
- 5. Level of significance is also called:
- a) Power of the test
- 6. The chance of rejecting a true hypothesis decreases when sample size is:
- b) Increase
- 7. Which of the following testing is concerned with making decisions using data?
- b) Hypothesis
- 8. What is the purpose of multiple testing in statistical inference?
- d) All of the mentioned
- 9. Normalized data are centred at and have units equal to standard deviations of the original data
- a) 0

Q10and Q15 are subjective answer type questions, Answer them in your own words briefly

10. What Is Bayes' Theorem?

Bayes, is a mathematical formula for determining conditional Probability is the likelihood of an outcome occurring, based on a previous outcome having occurred in similar circumstances. Bayes' theorem provides a way to revise existing predictions or theories (update probabilities) given new or additional evidence.

P(A/B)=P(A/B)P(A)/P(B)

It is used in Machine Learning.

11. What is z-score?

Z-score is a statistical measurement that describes a value's relationship to the mean of a group of values. Z-score is measured in terms of standard deviations from the mean. If a Z-score is 0, it indicates that the data point's score is identical to the mean score

$$z = (x - \mu) / \sigma$$

- z = Z-score
- x = the value being evaluated
- μ = the mean
- σ = the standard deviation

12. What is t-test?

a statistical test that is used to compare the means of two groups. It is often used in hypothesis testing to determine whether a process or treatment actually has an effect on the population of interest, or whether two groups are different from one another.

13. What is percentile?

A percentile is a comparison score between a particular score and the scores of the rest of a group. It shows the percentage of scores that a particular score surpassed.

14. What is ANOVA?

Analysis of variance, or ANOVA, is a statistical method that separates observed variance data into different components to use for additional tests. A one-way ANOVA is used for three or more groups of data, to gain information about the relationship between the dependent and independent variables.

15. How can ANOVA help?

ANOVA is helpful for **testing three or more variables**. It is similar to multiple two-sample t-tests. However, it results in fewer type I errors and is appropriate for a range of issues. ANOVA groups differences by comparing the means of each group and includes spreading out the variance into diverse sources