

Table S5. Binding affinities of CRB-701 and CRB-701-mAb to human Fc receptors and C1q determined by BLI

	CRB-701 mAb binding affinity, K_d (M)		CRB-701 binding affinity, K_d (M)	
	Mean	SD	Mean	SD
FcRn	2.7×10^{-8}	0.29×10^{-8}	3.0×10^{-8}	0.22×10^{-8}
FcγRIa	4.1×10^{-9}	0.78×10^{-6}	3.8×10^{-9}	0.69×10^{-6}
FcγRIIa	1.4×10^{-6}	0.21×10^{-6}	3.1×10^{-6}	0.53×10^{-6}
FcγRIIb	2.9×10^{-6}	0.20×10^{-6}	5.6×10^{-6}	0.46×10^{-6}
FcγRIIIa F176	3.8×10^{-6}	0.36×10^{-6}	3.4×10^{-6}	1.41×10^{-6}
FcγRIIIa V176	1.1×10^{-6}	0.16×10^{-6}	3.9×10^{-6}	1.18×10^{-6}
C1q	6.3×10^{-9}	0.11×10^{-9}	6.5×10^{-9}	0.12×10^{-9}

Mean and SD values are derived from three manufactured lots of CRB-701-mAb and four manufactured lots of CRB-701.

BLI, biolayer interferometry; K_d , binding affinity (k_{off}/k_{on}); mAb, monoclonal antibody; SD, standard deviation.