

Table S7. Stability of CRB-701-mAb (0.02, 0.2, or 2 μ M) over time following incubation with human, cynomolgus monkey, or Sprague Dawley rat plasma, or PBST control

	Remaining CRB-701-mAb (%) over time (hours)						
	0 h	24 h	48 h	72 h	96 h	168 h	336 h
Human							
CRB-701, 0.02 μ M	100	93.45	86.96	87.69	86.15	85.59	73.53
CRB-701, 0.2 μ M	100	94.18	90.06	93.26	80.37	84.85	75.53
CRB-701, 2 μ M	100	93.42	87.86	83.64	82.05	82.69	66.05
Cynomolgus monkey							
CRB-701, 0.02 μ M	100	98.69	100.19	98.88	100.29	100.13	89.88
CRB-701, 0.2 μ M	100	99.86	98.21	99.53	96.93	94.86	81.08
CRB-701, 2 μ M	100	95.40	97.84	100.66	100.06	101.18	88.89
Sprague Dawley rat							
CRB-701, 0.02 μ M	100	96.77	91.05	90.98	89.48	87.35	81.33
CRB-701, 0.2 μ M	100	91.22	88.65	87.60	85.05	83.59	75.91
CRB-701, 2 μ M	100	95.47	88.10	88.10	87.13	86.14	80.34
PBST (control)							
CRB-701, 0.02 μ M	100	98.92	98.46	99.41	160.6	98.11	112.16
CRB-701, 0.2 μ M	100	105.36	103.78	103.38	106.91	97.27	115.36
CRB-701, 2 μ M	100	103.64	100.99	103.03	101.42	88.00	101.87

CRB-701-mAb levels were determined by ELISA. Remaining CRB-701-mAb levels are shown as the proportion of the total concentration at time point 0. Mean values are shown for two replicates.

ELISA, enzyme-linked immunosorbent assay; mAb, monoclonal antibody; PBST, phosphate-buffered saline with 0.1% Tween-20.