

Table S8. Mean MMAE release (%) over time following incubation of CRB-701 (0.02, 0.2, or 2 μ M) with human, cynomolgus monkey, or Sprague Dawley rat plasma, or PBST control

	MMAE release (%) over time (hours)						
	0 h	24 h	48 h	72 h	96 h	168 h	336 h
Human							
CRB-701, 0.02 μ M	0.00	0.00	0.10	0.13	0.21	0.47	0.82
CRB-701, 0.2 μ M	0.00	0.05	0.10	0.16	0.22	0.43	0.93
CRB-701, 2 μ M	0.00	0.05	0.10	0.17	0.22	0.48	1.00
Cynomolgus monkey							
CRB-701, 0.02 μ M	0.00	0.00	0.11	0.19	0.24	0.44	0.72
CRB-701, 0.2 μ M	0.00	0.06	0.11	0.19	0.26	0.47	0.72
CRB-701, 2 μ M	0.00	0.05	0.12	0.19	0.26	0.52	0.75
Sprague Dawley rat							
CRB-701, 0.02 μ M	0.00	0.06	0.14	0.22	0.29	0.45	0.97
CRB-701, 0.2 μ M	0.00	0.07	0.15	0.24	0.30	0.55	1.20
CRB-701, 2 μ M	0.00	0.07	0.14	0.23	0.40	0.74	1.04
PBST (control)							
CRB-701, 0.02 μ M	0.00	0.00	0.00	0.14	0.10	0.18	0.20
CRB-701, 0.2 μ M	0.00	0.03	0.05	0.08	0.10	0.18	0.24
CRB-701, 2 μ M	0.00	0.02	0.05	0.08	0.11	0.18	0.25

MMAE release is determined by LC-MS/MS. The measured MMAE concentration at each time point is shown as a percentage of the theoretically expected molality. Mean values are shown for two replicates.

LC-MS/MS, liquid chromatography–tandem mass spectrometry; MMAE, monomethyl auristatin E; PBST, phosphate-buffered saline with 0.1% Tween-20.