
Caveat lector. One day I will type up the first few lectures.

8 1/30/23

Relevant reading: Weintraub pp. 11–13, Hatcher pp. 70–76.

8.1 Deck Transformations

We will first begin with an example to motivate our definition:

Example 8.1. Let $S^1 = \{z \in \mathbb{C} \mid |z| = 1\} \subseteq \mathbb{C}$ be the circle regarded as a subspace of \mathbb{C} . Then we saw that $p : \mathbb{R} \rightarrow S^1$ via $p(t) = e^{2\pi it}$ was a covering map. Then for any $z_0 \in S^1$ with $p(t_0) = z_0$, we have that $p^{-1}(\{z_0\}) = t_0 + \mathbb{Z}$. Equivalently, $p(t_0 + m) = p(t_0)$ for all $m \in \mathbb{Z}$. Define, for $m \in \mathbb{Z}$, $T_m : \mathbb{R} \rightarrow \mathbb{R}$ via $T_m(t) = t + m$ translation by m . Then by our discussion, $p \circ T_m = p$. We say that T_m is an example of a **deck transformation**.

Definition 8.2. Let $p : E \rightarrow B$ be a covering projection. Then the **group of deck transformations** is the set

$$\Gamma_p := \{T \in \text{Homeo}(E) \mid p \circ T = p\},$$

where the endowed operation is function composition. That is, it is the set of all homeomorphisms such that for any $T \in \Gamma_p$, the following diagram commutes:

$$\begin{array}{ccc} E & \xrightarrow{T} & E \\ & \searrow p & \downarrow p \\ & & B \end{array}$$

Now given a covering projection p , we may define an equivalence relation in the following manner: for $x, y \in E$, we say $x \sim y$ if and only if there exists some $T \in \Gamma_p$ such that $T(x) = y$. Now we may consider the quotient space E/Γ_p , i.e., the topology that makes the projection $\hat{p} : E \rightarrow E/\Gamma_p$ continuous.

Example 8.3. Returning to our example of S^1 and p defined in Example 8.1, we now ask the question, what is \mathbb{R}/Γ_p with this equivalence relation?

Claim 8.4. $\Gamma_p = \{T_m \mid m \in \mathbb{Z}\}$.

To see this, proceed in the manner as we did when proving that p was a covering map. If $e^{2\pi i T(t)} = e^{2\pi i t}$ for all $t \in \mathbb{R}$, then rearranging, we see that $e^{2\pi i (T(t) - t)} = 1$ for all t . Hence $T(t) - t \in \mathbb{Z}$ for all t , but since $T(t) - t$ is continuous, we conclude that $T(t) - t$ is constant, and so there exists some $m \in \mathbb{Z}$ such that $T(t) = t + m$ for all t .

We may now also further say that $\Gamma_p \simeq \mathbb{Z}$. Hence we may identify \mathbb{R}/Γ_p with \mathbb{R}/\mathbb{Z} , or with $[0, 1)$.

Notation. We will denote $\Gamma_p(x) := \text{orb}_{\Gamma_p}(x) = \{T(x) \mid T \in \Gamma_p\}$. In general, if we want to make some sort of identification for E/Γ_p with some set S , like we did in the previous example, we need $\#(\Gamma_p(x) \cap S) = 1$ for all x . Indeed, this is the case for $[0, 1)$.

8.2 Discontinuous Actions

Definition 8.5. Let E be a topological space, and let $\Gamma \leq \text{Homeo}(E)$. We say that Γ **acts discontinuously** if for all $x \in E$, there exists some open neighborhood U_x of x such that if $T \in \Gamma$ and $T(U_x) \cap U_x \neq \emptyset$, then $T = \text{id}$.

Remark 8.6. Some texts, like Hatcher, calls a discontinuous action as a covering space action.

One consequence of our definition is the following claim:

Claim 8.7. If Γ acts discontinuously on E and $S_1, S_2 \in \Gamma$, and $S_1(U) \cap S_2(U) \neq \emptyset$ for some nonempty U , then $S_1 = S_2$.

Proof of claim. Observe that, since S_1 and S_2 are homeomorphisms, $\emptyset \neq S_1(U) \cap S_2(U) = S_1(U \cap S_1^{-1} \circ S_2(U))$. In particular, this implies that $U \cap S_1^{-1} \circ S_2(U) \neq \emptyset$. Since $S_1^{-1} \circ S_2 \in \Gamma$ and Γ acts discontinuously, we conclude $S_1^{-1} \circ S_2 = \text{id}$. ■

Lemma 8.8. Suppose Γ acts discontinuously on E . Then $p : E \rightarrow E/\Gamma$ is a covering projection, where the quotient is defined by $x \sim_\Gamma y$ if and only if there is some $T \in \Gamma$ such that $Tx = y$.

Proof. Given $y \in E/\Gamma$, take $x \in E$ such that $p(x) = y$, and let U_x be the neighborhood that is granted by Definition 8.5. Then we claim that $p(U_x)$ is open. To see this, notice that $p^{-1}(p(U_x)) = \bigsqcup_{S \in \Gamma} S(U_x)$, and since each S is a homeomorphism, $S(U_x)$ is open, which implies that $p^{-1}(p(U_x))$ is open, as desired. Moreover, $p|_{S(U_x)} : S(U_x) \rightarrow p(U_x)$ is a homeomorphism, and thus p must be a covering map. ■

8.3 Universal Covering Spaces

We will state two key theorems, but we will not prove them.

Theorem 8.9. Let E be a simply connected space, and let $p : E \rightarrow B$ be a covering projection. Assume further that B is semilocally simply connected. Then if Γ_p is the group of deck transformations, then Γ acts discontinuously and B is homeomorphic to E/Γ_p . In particular, the following diagram commutes:

$$\begin{array}{ccc} E & \xrightarrow{\text{id}} & E \\ \downarrow p & & \downarrow \hat{p} \\ B & \xrightarrow{h} & E/\Gamma_p \end{array}$$

where $h : B \rightarrow E/\Gamma_p$ denotes the homeomorphism and \hat{p} is the projection map from E to E/Γ_p .

Theorem 8.10 (Existence and Universal Property of Universal Covers). Let B be a semilocally simply connected, locally path connected, connected space. Then there exists a simply connected and connected space E such that there is a covering projection $p : E \rightarrow B$. Moreover, if $q : X \rightarrow B$ is any other covering projection, with X connected, then there exists a unique continuous map $r : E \rightarrow X$ such that the following diagram commutes:

$$\begin{array}{ccc} E & \xrightarrow{r} & X \\ \downarrow p & \swarrow q & \\ B & & \end{array}$$

The space E is unique up to homeomorphism.

Definition 8.11. The space E in the previous theorem is called a **universal cover**, and we will denote a universal covering space of B by \tilde{B} .

Interpretation. We can interpret the previous two theorems in the following way: by Theorem 8.10 we know that for any semilocally simply connected space B there is a universal cover \tilde{B} , and Theorem 8.9 tells us that $B \approx \tilde{B}/\Gamma_p$. Moreover, in a sense, Γ_p is the fundamental group.

Proposition 8.12. Let $p : E \rightarrow B$ be a covering projection and assume further that E is simply connected and path connected. Suppose $b_0 \in B$, and $e_0 \in E$ such that $p(e_0) = b_0$. Then $\pi_1(B, b_0) \simeq \Gamma_p$.

Proof. We will show that there is a one-to-one correspondence between the two groups. Let $T \in \Gamma_p$. Let $\tilde{\alpha}$ be a curve in E connecting e_0 and $T(e_0)$, and let $\alpha := p \circ \tilde{\alpha}$. Then observe that $\alpha(0) = p(e_0) = b_0 = \alpha(1) = p(T(e_0))$. So $\alpha \in \pi_1(B, b_0)$. Thus this gives us a way to assign a loop in $\pi_1(B, b_0)$ for every $T \in \Gamma_p$. To see that this does not depend on our choice of curve $\tilde{\alpha}$, suppose $\tilde{\beta}$ is another curve connecting e_0 and $T(e_0)$, and set $\beta = p(\tilde{\beta})$. Then the concatenation $\tilde{\alpha} \cdot \tilde{\beta}^{-1}$ is a loop with basepoint e_0 , and since E is simply connected must be homotopically trivial. Hence there is a homotopy $\tilde{H} : I \times I \rightarrow E$ such that $h_0 = \tilde{\alpha} \cdot \tilde{\beta}^{-1}$, and $h_1 \equiv e_0$. Now consider $H : I \times I \rightarrow B$ given by $H(s, t) = p(\tilde{H}(s, t))$. Then this is a homotopy taking the concatenation $\alpha \cdot \beta^{-1}$ to the constant path b_0 , which shows that $\alpha = \beta$ in $\pi_1(B, b_0)$. Hence this map is well-defined.

On the other hand, let $\gamma \in \pi_1(B, b_0)$. Then by the path-lifting property, there exists some $\tilde{\gamma} : [0, 1] \rightarrow E$ such that $\tilde{\gamma}(0) = e_0$ and $p(\tilde{\gamma}(1)) = \gamma(1) = b_0$. Moreover, there exists some $T \in \Gamma_p$ such that $T(e_0) = \tilde{\gamma}(1)$; this defines the inverse map. By the homotopy lifting property, it is easy to check that this map does not depend on the representative of the homotopy class. Hence it is well-defined. The two maps are clearly bijective, and it is straightforward to check that they are homomorphisms. Hence this proves the isomorphism. ■


9 2/1/23

Today we will introduce the Seifert-van Kampen theorem. Relevant reading: Hatcher Chapter 1.2, Weintraub Section 2.3.

9.1 Free Group Products

Definition 9.1 (Free Group Products). Given two groups G_1 and G_2 , we denote $G_1 * G_2$ to be the **free product** of G_1 and G_2 , which is the coproduct of the groups G_1 and G_2 in the category of groups. That is, there are injective homomorphisms $\iota_1 : G_1 \hookrightarrow G_1 * G_2$ and $\iota_2 : G_2 \hookrightarrow G_1 * G_2$ and it satisfies the following universal property:

If G is any group and $f_1 : G_1 \rightarrow G$ and $f_2 : G_2 \rightarrow G$ are homomorphisms, then there exists a unique homomorphism $f : G_1 * G_2 \rightarrow G$ such that the following diagram commutes:



Example 9.2. $\mathbb{Z} * \mathbb{Z} = F_2$ the free group on two generators: alternatively, we can write F_2 to be the set of all finite words on two letters a, b .

Remark 9.3. In general, if $a_i \in G, b_i \in G$, we can write any element of $G_1 * G_2$ as $a_1 b_1 a_2 b_2 \cdots a_k b_k$.

Definition 9.4. Given a group G , and $A \subseteq G$ (not necessarily a subgroup), the **normal subgroup generated by A** is defined by $N(A) = \bigcap N$, where the intersection runs over all normal subgroups containing A : that is, it is the smallest normal subgroup of G containing A .


9.2 The Seifert-van Kampen Theorem and Applications

Theorem 9.5 (Seifert-van Kampen). *Let X be a path-connected space, and assume $X = U_1 \cup U_2$, where both U_1 and U_2 are open and path-connected. Let $x_0 \in U_1 \cap U_2$ and assume that $U_1 \cap U_2$ is also path-connected. Then $\pi(X, x_0) \simeq (\pi_1(U_1, x_0) * \pi_1(U_2, x_0)) / N(A)$, where if $(\iota_1)_*$ and $(\iota_2)_*$ are the homomorphisms induced by the inclusion map $\iota_i : U_i \rightarrow X$, we have*

$$A = \{(\iota_1)_*(g^{-1}) * (\iota_2)_*(g) \mid g \in \pi_1(U_1 \cap U_2, x_0)\}.$$


We will not prove the Seifert-van Kampen theorem today, but we will see some applications of it.

Example 9.6 (An Incorrect Application). Consider S^1 as the union of two open intervals I_1 and I_2 as in the figure. But $\pi_1(S^1)$ cannot be a quotient of the free product $\pi_1(I_1) * \pi_1(I_2)$ because the two factors are both trivial, but we already know that $\pi_1(S^1) \simeq \mathbb{Z}$. The error was in that the hypothesis $U_1 \cap U_2$ is not path-connected.



Proposition 9.7 (Fundamental Group of S^n). *For $n \geq 2$, S^n is simply connected.*


Proof. We will use the ideas from Example 9.6. Let $x_0 \in S^n$; by rotating the sphere, we may assume that x_0 is on the equator. Write $S^n = D_1 \cup D_2$, where D_1 and D_2 are the open sets in the figure below.



Note that D_1 and D_2 are both contractible, and so must have trivial fundamental group. Moreover, $D_1 \cap D_2 \approx S^{n-1} \times I$, which is also path-connected. Then applying the Seifert-van Kampen theorem, $\pi_1(S^n, x_0)$ must be a quotient of $\pi_1(D_1, x_0) * \pi_1(D_2, x_0) = \{0\}$. Hence S^n is simply connected. ■

Remark 9.8. We could have used the stereographic projection to map the sphere with the poles removed onto \mathbb{R}^n in the previous proof.

Example 9.9 (The Figure 8). Consider $E := S^1 \vee S^1$, or the “figure 8,” joined together at the point x_0 . Let U_1 and U_2 be as in the figure below, so that $U_1 \cup U_2 = E$, and $U_1 \cap U_2$ is the cross in the middle.




First observe that all our spaces are path-connected and so the hypotheses of the Seifert-van Kampen theorem are satisfied. Next, $U_1 \cap U_2$ is contractible, which implies that $\pi_1(U_1 \cap U_2, x_0) = \{0\}$.

Finally, observe that $U_1 \approx U_2 \approx S^1$, which implies that $\pi_1(U_1, x_0) \simeq \pi_1(U_2, x_0) \simeq \mathbb{Z}$. Appealing to the Seifert-van Kampen theorem, we conclude that $\pi(E, x_0) = \mathbb{Z} * \mathbb{Z}$.


Exercise 9.10. Apply induction to the previous example to conclude that the fundamental group of the n -petal rose is F_n , the free group on n elements.

Exercise 9.11. Let X and Y be topological spaces, and suppose $X \vee Y$ be locally contractible and/or semilocally simply connected at the attaching point x_0 . Show that $\pi_1(X \vee Y, x_0) \simeq \pi_1(X, x_0) * \pi_1(Y, x_0)$.


Example 9.12 (The Torus). Let \mathbb{T}^2 denote the torus $\mathbb{T}^2 = S^1 \times S^1$. We have already noted that $\pi_1(\mathbb{T}^2) \simeq \pi_1(S^1) \times \pi_1(S^1) \simeq \mathbb{Z} \times \mathbb{Z}$. Now we will use the Seifert-van Kampen's theorem to prove this. We have shown that the torus may be considered as the quotient space of the square where the opposite edges are identified. Now let U_1 and U_2 be as in the diagram, where U_1 is the “outer” part of the square, and U_2 the “inner” part.




To see that U_1 is open, note that on the edges, any ball would “bleed over” to the opposite edge, as in the following figure:



It is now easy to see that U_1 and U_2 are both open, $U_1 \cup U_2 = \mathbb{T}^2$, and U_1, U_2 , and $U_1 \cap U_2$ are all path-connected. The hypotheses of the Seifert-van Kampen theorem are now satisfied. Fix $x_0 \in U_1 \cap U_2$. First observe that U_2 is contractible, and so $\pi_1(U_2, x_0) = \{0\}$. On the other hand, we see that U_1 deformation retracts onto the boundary of the square, and then identified with the figure 8 in the following manner:



Since deformation retracts induce an isomorphism of fundamental groups, we have from Example 9.9 $\pi_1(U_1, x_0) \simeq \pi_1(S^1 \vee S^1) \simeq \mathbb{Z} * \mathbb{Z}$. Now $U_1 \cap U_2$ is the annulus, which deformation retracts onto the circle S^1 , so its fundamental group is the free group on one generator, the loop going around the annulus once counterclockwise. The following figure shows its image under $(\iota_1)_*$:



Now after the deformation retract, we see that in the image this loop is exactly the commutator $aba^{-1}b^{-1}$. But this was the image of the generator, and so we conclude that $N(A)$ (in the statement of the theorem) must be the commutator subgroup inside $\pi_1(U_1, x_0)$. Therefore $\pi(\mathbb{T}^2, x_0) \simeq \mathbb{Z} * \mathbb{Z} / \langle aba^{-1}b^{-1} \rangle = \mathbb{Z} \times \mathbb{Z}$.

Remark 9.13. The above proof can be adapted to compute $\pi_1(\mathbb{T}^n, x_0)$ with induction.


10 2/3/23

Today we will continue with examples of van Kampen's Theorem.


10.1 The Genus 2 Surface

Recall when we computed the fundamental group of the torus via Seifert-van Kampen theorem, we used the quotient of a square that is homeomorphic to the torus.


Method 1. For the genus 2 surface S , we will consider the quotient of an octagon as follows:



Then just as we did for the torus, decompose the octagon into following pieces:



First note that all these sets are path-connected, so the hypotheses of the van Kampen theorem are satisfied. Then notice that U_1 deformation retracts onto the boundary, which is homeomorphic to the 4-petal rose; thus $\pi_1(U_1) \simeq F_4$, the free group on four elements. Moreover, U_2 is contractible and so has trivial fundamental group. Finally, $U_1 \cap U_2$ is the annulus, which deformation retracts onto S^1 , so has fundamental group the free group on one generator. Then by the Seifert-van Kampen theorem, we have that $\pi_1(S) \simeq \pi_1(U_1)/N(\iota_1(g) \mid g \in \pi_1(U_1 \cap U_2))$. Let g be a loop in $U_1 \cap U_2$, like in the diagram below. Then considered as a loop in U_1 and its image in the deformation retract, its image is $abcd a^{-1} b^{-1} c^{-1} d^{-1}$.




Therefore $\pi_1(S) \simeq F_4 / \langle abcd a^{-1} b^{-1} c^{-1} d^{-1} \rangle \simeq \langle a, b, c, d \mid abcd = dcba \rangle$.


Method 2. The idea will be to decompose the surface into two parts, just like below:




Consider the octagon again, but this time as a subspace of the square below after cutting and pasting.



Just as above, consider U_1 and U_2 defined as in the figure below.



Then both U_1 and U_2 deformation retract onto the figure 8, and $U_1 \cap U_2$ is an annulus which deformation retract onto S^1 . Hence $\pi_1(U_1) \simeq \pi_1(U_2) \simeq F_2$ the free group on two generators, and $\pi_1(U_1 \cap U_2) \simeq \mathbb{Z}$. Now consider the single loop $g \in \pi_1(U_1 \cap U_2)$ given by the generator: that is, the loop that goes around once in the annulus. Then considered as a loop in U_1 and U_2 respectively, the diagram below shows that in U_1 it deformation retracts onto the loop $aba^{-1}b^{-1}$, and in U_2 it deformation retracts onto the loop $cdc^{-1}d^{-1}$.




Therefore $(\iota_1)_*(g) = aba^{-1}b^{-1}$ and $(\iota_2)_*(g) = cdc^{-1}d^{-1}$. Thus applying van Kampen's theorem, we conclude that $\pi_1(S) \simeq F_2 * F_2 / N \left(aba^{-1}b^{-1} (cdc^{-1}d^{-1})^{-1} \right) \simeq \langle a, b, c, d \mid [a, b] = [c, d] \rangle$.

Remark 10.1. We can compute the fundamental group of a genus g surface by induction.

Corollary 10.2. *The fundamental group of a genus 2 surface with a point deleted is the free group on 4 elements. In general, the fundamental group of a genus g surface, with $g \geq 2$, is the free group on $2g$ elements.*

Proof. The genus 2 surface with a point deleted can be identified with the quotient space of the octagon in Method 1 with a neighborhood deleted in the interior, as in the diagram. Then this deformation retracts onto the boundary, which is homeomorphic to the 4-petal rose.




The general case is handled by induction. ■

10.2 Higher Homotopy Groups

Recall that the fundamental group $\pi_1(X, x_0)$ was all about maps of the form $\gamma : (S^1, 1) \rightarrow (X, x_0)$, or equivalently maps of the form $([0, 1], \{0, 1\}) \rightarrow (X, x_0)$. Now similarly, we define the **higher homotopy groups** in the following manner: $\pi_n(X, x_0) := [(S^n, 1); (x, x_0)]$ where the bracket denotes the homotopy classes of maps. Equivalently, we may define $\pi_n(X, x_0)$ as $[(I_n, \partial I_n), (X, x_0)]$ where $I_n = [0, 1]^n$.

The group operation is defined as follows: as an example, we will use π_2 , and analogize.



Because we stipulate that the boundary gets mapped to x_0 , the multiplication is well-defined by the pasting lemma. The identity element is the constant map mapping to x_0 . Another way of writing the multiplication is as follows: if $\gamma_1, \gamma_2 : (t_1, \dots, t_n) \rightarrow X$, then we may write their product to be

$$(\gamma_1 * \gamma_2)(t_1, \dots, t_n) = \begin{cases} \gamma_1(2t_1, t_2, \dots, t_n), & t \in [0, 1/2] \\ \gamma_2(2t_1 - 1, t_2, \dots, t_n), & t \in [1/2, 1]. \end{cases}$$

Next, the following figure from Hatcher illustrates the following lemma:


$$\begin{array}{|c|c|} \hline f & g \\ \hline \end{array} \simeq \begin{array}{|c|c|} \hline \begin{array}{|c|c|} \hline f & g \\ \hline \end{array} & \\ \hline \end{array} \simeq \begin{array}{|c|} \hline \begin{array}{|c|} \hline f \\ \hline g \\ \hline \end{array} \\ \hline \end{array} \simeq \begin{array}{|c|c|} \hline \begin{array}{|c|} \hline g \\ \hline f \\ \hline \end{array} & \\ \hline \end{array} \simeq \begin{array}{|c|c|} \hline g & f \\ \hline \end{array}$$

Lemma 10.3. $\pi_n(X, x_0)$ is abelian for $n \geq 2$.

11 2/6/23


11.1 Relative Homotopy Groups

Definition 11.1. As we did last time, define $I_n = [0, 1]^n$, ∂I_n the boundary of I_n , and let $J_n := \partial I_n \setminus (I_{n-1} \times \{0\})$, as in the following diagram:




Then we define the **relative homotopy groups** as following: $\pi_n(X, A, x_0) := [(I_n, \partial I_n, J_n); (X, A, x_0)]$.

Because the elements of the homotopy groups are equivalence classes, we will write what it means for two elements to be equivalent. We say that for $\gamma, \eta \in \pi_n(X, A, x_0)$, $\gamma \sim \eta$ if and only if there exists $F : (I_n, \partial I_n, J_n) \times [0, 1] \rightarrow (X, A, x_0)$ such that $f_0 = F(\cdot, 0) = \gamma$ and $f_1 = F(\cdot, 1) = \eta$. Similar as was done in homotopy groups, the product $\gamma \cdot \eta$ is defined in the following way:




Lemma 11.2. This product makes $\pi_n(X, A, x_0)$ into a group, and for $n \geq 3$, is abelian.

Proof. To see that $\pi_n(X, A, x_0)$ is obvious. To see that it is obvious, consider the following figure:



It is important that the two cubes inside the big cube in the figure above do not have their bases taken off of $\partial I_n \setminus J_n$. The fact that this is not possible in dimension two illustrates why $\pi_n(X, A, x_0)$ is not abelian:



■

11.2 Exact Sequences

Definition 11.3. Given groups G_1, G_2, \dots and homomorphisms $L_n : G_n \rightarrow G_{n+1}$, we say that the sequence

$$G_1 \xrightarrow{L_1} G_2 \xrightarrow{L_2} \cdots \rightarrow G_n \xrightarrow{L_n} \cdots \rightarrow 0$$

is **exact** if $\ker L_{n+1} = \operatorname{im} L_n$ for each n .

Observe that if we have a sequence of groups, $L_{n+1} \circ L_n \equiv 0$ if and only if $\ker L_{n+1} \supseteq \operatorname{im} L_n$.

Example 11.4. Consider the sequence $0 \xrightarrow{L_1} G \xrightarrow{L_2} 0$. Certainly $L_2 \circ L_1 \equiv 0$. But $\operatorname{im} L_1 = \{0\}$ since L_1 is a homomorphism. Thus this sequence must be exact if and only if G is trivial.

Example 11.5. Consider the sequence $0 \xrightarrow{L_1} G \xrightarrow{L_2} H \xrightarrow{L_3} 0$, and suppose that it is exact. Then $\operatorname{im} L_1 = \{0\} = \ker L_2$, which implies that L_2 is injective. On the other hand, $H = \ker L_3 = \operatorname{im} L_2$, and so L_2 is surjective. Thus L_2 is a group isomorphism.

Example 11.6. Consider the exact sequence $0 \xrightarrow{L_1} N \xrightarrow{\iota} G \xrightarrow{\pi} H \xrightarrow{L_3} 0$. Since $H = \ker L_3 = \text{im } \pi$, we have that π is surjective. On the other hand, $\text{im } L_1 = \ker \iota = \{0\}$ and so ι is injective. Thus by the first isomorphism theorem, $G/\ker \pi = G/\iota(N) \simeq H$. Identifying N with its image under ι , we conclude that $G/N \simeq H$.


12 2/8/23

12.1 Long Exact Sequences of Relative Homotopy Groups


Theorem 12.1 (Long Exact Sequences of Relative Homotopy Groups). *Let $J : (X, x_0, x_0) \hookrightarrow (X, A, x_0)$ be the inclusion. Then there is a long exact sequence*

$$\rightarrow \pi_n(A, x_0) \xrightarrow{\iota_{\#}} \pi_n(X, x_0) \xrightarrow{J_{\#}} \pi_n(X, A, x_0) \xrightarrow{\partial} \pi_{n-1}(A, x_0) \rightarrow \cdots \rightarrow \pi_0(X, A, x_0) \rightarrow 0.$$

The boundary map $\partial : \pi_n(X, A, x_0) \rightarrow \pi_{n-1}(A, x_0)$ is defined in the following manner: for $\gamma \in \pi_n(X, A, x_0)$, say $\gamma : (I_n, \partial I_n, J_n) \rightarrow (X, A, x_0)$, the restriction $\gamma|_{I_{n-1} \times \{0\}}$ can be regarded as a map $(I_{n-1}, \partial I) \rightarrow (A, x_0)$. Then $\partial\gamma$ is precisely this restriction.




Proof. First we check that ∂ is well-defined. To this end suppose $\gamma_0 \sim \gamma_1$. Then there exists some $F : (I_n, \partial I_n, J_n) \times [0, 1] \rightarrow (X, A, x_0)$ such that $F(\cdot, 0) = f_0 = \gamma_0$ and $F(\cdot, 1) = f_1 = \gamma_1$ (a 2-dimensional schematic diagram is below).




Then restricting F to $I_{n-1} \times \{0\} \approx I_{n-1}$ gives a homotopy $F|_{I_{n-1} \times \{0\} \times [0,1]} : I_{n-1} \times [0,1] \rightarrow A$. Then this gives a homotopy $\partial(\gamma_1) = \gamma_1|_{I_{n-1} \times \{0\}}$ to $\partial(\gamma_2) = \gamma_2|_{I_{n-1} \times \{0\}}$. Thus $\partial(\gamma_1) \sim \partial(\gamma_2)$ in homotopy classes.

Now we check exactness. First, we will show that $\ker \iota_{\#} = \text{im } \partial$. To prove one direction, suppose that $\eta \in \text{im } \partial$. Then there exists some $\gamma \in \pi_{n+1}(X, A, x_0)$, $\gamma : (I_{n+1}, \partial I_{n+1}, J_{n+1}) \rightarrow (X, A, x_0)$ such that $\partial\gamma = \gamma|_{I_n \times \{0\}} = \eta$. In order to prove that $\eta \in \ker \iota_{\#}$, we must show that $\iota_{\#}(\eta) \sim 0$ in $\pi_n(X, x_0)$, that is, there exists some homotopy $F : I_n \times [0,1] \rightarrow X$ such that $f_0 = \eta$ and $f_1 = x_0$. Define $F(x, t) = \gamma(x, t)$, regarded as a map from $I_{n+1} = I_n \times [0,1]$ to X . Indeed, $F(x, 0) = \gamma(x, 0) = \gamma|_{I_n \times \{0\}}(x) = \eta(x)$. On the other hand, $F(x, 1) = \gamma(x, 1) = x_0$ since $\gamma \in \pi_{n+1}(X, A, x_0)$ (see the figure below). Hence $\iota_{\#}(\eta)$ is homotopic to the constant map in $\pi_n(X, x_0)$, so $\eta \in \ker \iota_{\#}$.



Conversely, suppose that $\eta \in \ker \iota_{\#}$. Then $\iota_{\#}(\eta)$ is homotopic to the constant map in $\pi_n(X, x_0)$, that is, there exists some $F : I_n \times [0,1] \rightarrow X$ such that $f_0(x) = \iota_{\#}(\eta)$ and $f_1(x) \equiv x_0$. Then proceeding as in the other direction, defining $\gamma : (I_{n+1}, \partial I_{n+1}, J_{n+1}) \rightarrow (X, A, x_0)$, with the identification $I_{n+1} = I_n \times [0,1] \rightarrow X$ via $\gamma(x, t) = F(x, t)$ we have that clearly $\partial\gamma = \eta$.

For the next part, we will show that $\text{im } \iota_{\#} = \ker J_{\#}$. Let $\eta \in \text{im } \iota_{\#}$. Then there exists some $\tilde{\eta} \in \pi_n(A, x_0)$, $\tilde{\eta} : (I_n, \partial I_n) \rightarrow (A, x_0)$ such that $\eta = \iota \circ \tilde{\eta}$. Now consider the map $J \circ \eta : (I_n, \partial I_n, J_n) \rightarrow (X, A, x_0)$, which obtained by changing the domain and codomain: note that $\eta = \iota \circ \tilde{\eta}$, and so the image of η is completely contained in A , and moreover for any $x \in \partial I_n$, we have $\eta(x) = \tilde{\eta}(x) = x_0$ since $\tilde{\eta} \in \pi_n(A, x_0)$. Thus $J \circ \eta$ as a map from $(I_n, \partial I_n, J_n) \rightarrow (X, A, x_0)$ makes sense. To show that $\eta \in \ker J_{\#}$, we will show that there is a homotopy $F : (I_n, \partial I_n, J_n) \times [0, 1] \rightarrow (X, A, x_0)$ such that $f_0 = J \circ \eta$ and $f_1 \equiv x_0$. Next, consider $F(x_1, \dots, x_{n-1}, x_n, t) := (J \circ \eta)(x_1, \dots, x_{n-1}, (1-t)x_n)$.



Indeed, $f_0(x_1, \dots, x_n) = F(x_1, \dots, x_n, 0) = J \circ \eta$, and $f_1(x_1, \dots, x_n) = F(x_1, \dots, x_n, 1) = (J \circ \eta)(x_1, \dots, x_{n-1}, 0) = \tilde{\eta}(x_1, \dots, x_{n-1}, 0) = x_0$. The schematic figure above shows that F indeed is the map of the desired form. Hence F is the desired homotopy.

Conversely, suppose $\eta \in \ker J_{\#}$. Then $\eta : (I_n, \partial I_n) \rightarrow (X, x_0)$ and $J \circ \eta : (I_n, \partial I_n, J_n) \rightarrow (X, A, x_0)$ is homotopically trivial. Thus there exists some $F : (I_n, \partial I_n, J_n) \times [0, 1] \rightarrow (X, A, x_0)$ such that $f_0 = J \circ \eta$ and $f_1 \equiv x_0$. Define $\tilde{\eta} : (I_n, \partial I_n) \rightarrow (A, x_0)$ by $\tilde{\eta}(x_1, \dots, x_n) = F(x_1, \dots, x_{n-1}, 0, x_n)$. Clearly by definition of F , $\tilde{\eta}$ takes image in A , and its boundary takes value in x_0 (see figure below for an illustration). We claim that $\iota \circ \tilde{\eta} \sim \eta$ in $\pi_n(X, x_0)$. Indeed, let $G : (I_n, \partial I_n) \times [0, 1] \rightarrow (X, x_0)$ via $G(x_1, \dots, x_n, t) = F(x_1, \dots, x_{n-1}, (1-t)x_n, tx_n)$. Then $g_0(x_1, \dots, x_n) = F(x_1, \dots, x_n, 0) = (J \circ \eta)(x_1, \dots, x_n) = \eta(x_1, \dots, x_n)$ and $g_1(x_1, \dots, x_n) = F(x_1, \dots, 0, x_n) = \tilde{\eta}(x_1, \dots, x_n)$. Pictorially, the green slanted rectangle depicts g_t during a time between 0 and 1, in the middle of the homotopy. Thus $\iota \circ \tilde{\eta} \sim \eta$ in $\pi_n(X, x_0)$, as desired.

13.1 Serre Fibrations and Hurewicz Fibrations

$$\begin{array}{ccc} X & \xrightarrow{h} & E \\ \downarrow \iota & \nearrow \tilde{F} & \downarrow p \\ X \times I & \xrightarrow{F} & B \end{array}$$

Lemma 13.2. *Let $p : E \rightarrow B$ be a continuous map, $\mathcal{U} = \{U_i\}_{i \in I}$ be an open cover of B , and let $p_i := p|_{U_i}$, that is, $p_i : p^{-1}(U_i) \rightarrow U_i$. If p_i has the homotopy lifting property for each i , then p has the homotopy lifting property for E .*

$$\begin{array}{ccc} F \times U_i & \xrightarrow{H_i} & p^{-1}(U_i) \\ & \searrow p_2 & \downarrow p \\ & & U_i \end{array}$$

17

Theorem 13.5 (Long Exact Sequences of Fibrations). *Given a Serre fibration $p : E \rightarrow B$, $p(e_0) = x_0$, $p^{-1}(x_0) = F$ and $e_0 \in F$, there is a long exact sequence*

$$\cdots \rightarrow \pi_n(F, e_0) \xrightarrow{\iota_{\#}} \pi_n(E, e_0) \xrightarrow{p_{\#}} \pi_n(B, b_0) \xrightarrow{\partial} \pi_{n-1}(F, e_0) \rightarrow \cdots,$$


where the boundary map is defined in the following manner: if $\gamma : (I_n, \partial I_n) \rightarrow (B, b_0) \in \pi_n(B, b_0)$, then define $H : I_{n-1} \times I \rightarrow B$ by viewing γ as a homotopy. That is, $H(x_1, \dots, x_{n-1}, t) = \gamma(x_1, \dots, x_{n-1}, t)$. Then notice that $h_1 \equiv b_0$. Then since p is a Serre fibration, p satisfies the homotopy lifting property, so H lifts to a unique homotopy $\tilde{H} : X \times I \rightarrow E$ such that $p \circ \tilde{H} = H$ and $\tilde{h}_1 \equiv e_0$. Then we define $\partial\gamma = \tilde{H}|_{I_{n-1} \times \{0\}} : I_{n-1} \rightarrow E$. Then $p(\partial\gamma(x)) = p(\tilde{H}(x, 0)) = H(x_1, \dots, x_{n-1}, 0) = b_0$. Hence $\partial\gamma(x) \in p^{-1}(b_0) = F$, so $\partial\gamma \in \pi_{n-1}(F, e_0)$.

Proof Idea. We will show that there is a natural isomorphism between $\pi_n(B, x_0)$ and $\pi_n(X, A, x_0)$, and then appeal to Theorem 12.1, so that we can fit this long exact sequence into the previous one.

Proof. Consider $\pi_n(E, F, e_0) \xrightarrow{p_{\#}} \pi_n(B, b_0, b_0) \simeq \pi_n(B, b_0)$. We claim that $p_{\#}$ is an isomorphism, which will allow $\pi_n(B, b_0)$ to naturally fit into the long exact sequence for relative homotopy groups. To see this we will construct an inverse for $p_{\#}$. Take $\gamma : (I_n, \partial I_n, J_n) \rightarrow (E, F, e_0) \in \pi_n(E, F, e_0)$. Then this fits into the diagram

$$\begin{array}{ccc} (I_n, \partial I_n, J_n) & \xrightarrow{\gamma} & (E, F, e_0) \\ & \searrow p \circ \gamma & \downarrow p \\ & & (B, b_0, b_0) \end{array}$$

Take H and \tilde{H} as in the statement of Theorem 13.5. Then \tilde{H} is a homotopy $I_{n-1} \times [0, 1] \rightarrow E$ such that $\tilde{h}_1 = \tilde{H}(\cdot, 1) \equiv e_0$, and moreover $\tilde{H}(\partial I_n) \subseteq F = p^{-1}(b_0)$. Now consider the following deformation:



This gives us a new map $\hat{H} : I_{n-1} \times [0, 1] \rightarrow E$ homotopic to \tilde{H} . Then this induces a map $\text{hat} : \pi_n(B, b_0, b_0) \rightarrow \pi_n(E, F, e_0)$, $\gamma \mapsto \hat{H}$. It is (presumably) straightforward to check that hat is the inverse of $p_\#$, which concludes the proof. ■

Example 13.6. Let $p : E \rightarrow B$ be a covering projection. Then $F = p^{-1}(b_0)$ is discrete; hence $\pi_n(F, e_0) = \{0\}$ for all $n \neq 0$. Then we have the long exact sequence

$$\cdots \rightarrow \underbrace{\pi_n(F, e_0)}_{=0} \rightarrow \pi_n(E, e_0) \rightarrow \pi_n(B, b_0) \rightarrow \underbrace{\pi_{n-1}(F, e_0)}_{=0}.$$

for $n - 1 \geq 1$, that is, $n \geq 2$. Thus we have proven the following:

Corollary 13.7. *For $n \geq 2$, and $p : E \rightarrow B$ a covering projection, then $p_\# : \pi_n(E, e_0) \rightarrow \pi_n(B, b_0)$ is an isomorphism.*

Corollary 13.8. $\pi_n(S^1, 1) = 0$ for all $n \geq 2$.

Proof. Contractible spaces have trivial homotopy groups. ■

Definition 13.9. Let G be a given group and $n \in \mathbb{Z}$ an integer. A space (X, x_0) is called **Eilenberg-MacLane Space**, and we write $K(G, n)$, if $\pi_n(X, x_0) = G$ and $\pi_\ell(X, x_0) = 0$ for all $\ell \neq n$.

One observation to make is that for $\ell \geq 2$, we need the group G to be abelian, for $\pi_n(X, x_0)$ is abelian for $n \geq 2$. Moreover, Corollary 13.8 shows that $K(\mathbb{Z}, 1) = S^1$.

Now one might be wondering what $K(\mathbb{Z}, 2)$ might be. Continuing the above discussion, with the facts that $\pi_1(S^2) = 0$ and $\pi_2(S^2) \simeq \mathbb{Z}$, one might wonder if $S^2 = K(\mathbb{Z}, 2)$, but this is not the case, for $\pi_3(S^2) = \mathbb{Z}$; one way to see this is to use something called the Hopf fibration.

14 2/13/23

14.1 Homotopy Groups of S^n

One of our goals today will be to give a partial answer about computing a subset of all homotopy groups of S^n . Computing *all* of the homotopy groups of S^n , however, is still an open question!

Theorem 14.1. *For all $n \geq 1$ and $0 \leq k \leq n - 1$, $\pi_k(S^n) = 0$.*

But before we move to the proof of this theorem, let's begin with a warm-up.

Proposition 14.2. *For all $n \geq 1$, S^n is path-connected.*

Proof. Fix $x, y \in S^n$, and let $\bar{\gamma}_{x,y} : [0, 1] \rightarrow D^{n+1}$ via $\bar{\gamma}_{x,y}(t) = tx + (1 - t)y$, which is the straight line through the $n + 1$ -dimensional ball connecting x and y . Now consider $\gamma_{x,y} : [0, 1] \rightarrow S^n$ via $\gamma_{x,y}(t) := \bar{\gamma}_{x,y}(t)/|\bar{\gamma}_{x,y}(t)|$. Now this path $\gamma_{x,y}$ is well-defined and connects x and y , as long as x and y are not antipodal: that is, $x \neq -y$. In the case that x and y are antipodal, choose $z \in S^n$ such that z is not antipodal to x and y . Then $\gamma_{x,z}$ and $\gamma_{z,y}$ is well-defined, and their concatenation is a path connecting x and y . ■

Lemma 14.3. *If $f : M \rightarrow S^n$ is a continuous map that is not surjective, then it is homotopically trivial.*

Proof. Suppose f is not onto, say $p \notin f(M)$. Now by stereographic projection $h : S^n \setminus \{p\} \rightarrow \mathbb{R}^n$, we have the homeomorphism $S^n \setminus \{p\} \approx \mathbb{R}^n$. Note that \mathbb{R}^n is contractible: the map $c : \mathbb{R}^n \times I \rightarrow \mathbb{R}^n$ defined by $c(x, t) = (1 - t)x$ is the homotopy that contracts the identity map to a constant map. Now consider $F : M \rightarrow S^n$ defined by $F(u, t) = h^{-1}(c(t, h(f(u))))$: clearly this is a composition of continuous functions and is continuous. Moreover, $f_0(u) = F(u, 0) = h^{-1}(c(0, h(f(u)))) = h^{-1}(h(f(u))) = f(u)$, and $f_1(u) = h^{-1}(0)$ which is constant. Hence F is the desired homotopy. ■

Lemma 14.4. *If M is a manifold and $f : M \rightarrow S^n$ and $g : M \rightarrow S^n$ satisfy $|f(x) - g(x)| < 2$ for all $x \in M$, then f is homotopic to g .*

Proof. Consider $F : M \times [0, 1] \rightarrow S^n$ given by

$$F(x, t) := \frac{tf(x) + (1 - t)g(x)}{|tf(x) + (1 - t)g(x)|}.$$

Since $|f(x) - g(x)| < 2$ for all x , it follows that $f(x)$ can never be antipodal to $g(x)$. Hence F is well defined for all x, t , and so this is a homotopy. ■

Next, we will need two results (actually, corollaries) from analysis and smooth manifolds, which we will take as given.

Lemma 14.5 (Stone-Weierstrass). *Given $f : S^k \rightarrow S^n$ and an $\epsilon > 0$, there exists a polynomial $p : \mathbb{R}^k \rightarrow \mathbb{R}^n$ such that $|p(x) - f(x)| < \epsilon$ for all $x \in S^k$.*

Lemma 14.6 (Sard's Theorem). *If $f : M \rightarrow N$ and $f \in C^\infty$, and $\dim M < \dim N$, then f is not onto.*

Exercise 14.7. Construct a continuous function $\gamma : [0, 1] \rightarrow [0, 1]^m$ that is surjective.

Lemma 14.8. *Suppose $f : S^k \rightarrow S^n$ is continuous and $k < n$. Then f is homotopic to a map that is not surjective.*

Proof. Suppose f is as prescribed. Then by Lemma 14.5, there exists a polynomial $p : \mathbb{R}^k \rightarrow \mathbb{R}^n$ such that $|p(x) - f(x)| < 2$. But applying Lemma 14.4, f is homotopic to p . But p is a polynomial and hence smooth; appealing to Lemma 14.6 yields the result. ■

Now we have enough machinery to accomplish what we set out to do in the beginning of the section.

Proof of Theorem 14.1. Suppose $\gamma : (I_k, \partial I_k) \rightarrow (S^n, x_0)$. Note that $\gamma(\partial I_k) = x_0$, and $I_k/\partial I_k \approx S^k$. Now define $\pi : I_k \rightarrow S^k$ via the natural projection from the homeomorphism, $\pi(\partial I_k) = b_0 \in S^k$. Now define $f : S^k \rightarrow S^n$ by $f(x) = \gamma(\pi^{-1}(x))$. We need to check that f is well-defined: the only problematic point is when $x \neq b_0$, which has ∂I_k as the preimage. But indeed, γ maps ∂I_k to one point, so f is well-defined. Further, we have that f is continuous. Now Lemma 14.8 implies that there exists a homotopy $F : S^k \times [0, 1] \rightarrow S^n$ such that $f_0 = f$ and $f_1 \equiv \text{const}$.

Consider the path $\eta(t) := F(b_0, t)$. Then $\eta(0) = f_0(b_0) = f(b_0) = \gamma(\pi^{-1}(b_0)) = x_0$. We want to modify F to another homotopy \tilde{F} such that $\tilde{F}(b_0, t) = x_0$ for all t . Since $\text{SO}(n, \mathbb{R})$ acts transitively on the sphere, i.e., for all $P, Q \in S^n$, there exists some $O \in \text{SO}(n, \mathbb{R})$ such that $OP = Q$, and $O(S^n) = S^n$. Moreover, the choice of O varies continuously with respect to P and Q . Therefore we can choose O_t such that $O_t(\eta(t)) = x_0$ for all t continuously and $O_0 = \text{id}$, and define $\tilde{F}(x, t) = O_t(F(x, t))$. Then it is easy to verify that $\tilde{F}(\cdot, 0) = F(\cdot, 0) = f_0 = f$, $\tilde{F}(\cdot, 1) = O_1(F(\cdot, 1)) =$

$O_1(x_0) = x_0$, and $\tilde{F}(b_0, t) = x_0$ for all t by construction. Now set $H : (I_k, \partial I_k) \times I \rightarrow (S, x_0)$ by $H(x, t) := \tilde{F}(\pi(x), t)$; then with the identification made earlier, this is the desired homotopy, and so γ is trivial in $\pi_k(S^n, x_0)$. ■

Remark 14.9. A similar argument will work for any connected manifold.

14.2 Fiber Bundles and Lie Groups

Note that $\mathrm{SO}(n)$ acts transitively on S^n , and consider $\mathbf{1} = (1, 0, \dots, 0)^t$. Then we consider $\mathrm{stab}(\mathbf{1}) = \{O \in \mathrm{SO}(n) \mid O\mathbf{1} = \mathbf{1}\}$. Note that if $O \in \mathrm{stab}(\mathbf{1})$, then O takes $\mathbf{1}$ to $\mathbf{1}$, so must have the first

column be $\begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix}$: that is,

$$O = \left(\begin{array}{c|ccc} 1 & 0 & \cdots & 0 \\ \hline 0 & & & \\ \vdots & & \mathrm{SO}(n-1) & \\ 0 & & & \end{array} \right)$$

We get a map $p : \mathrm{SO}(n) \rightarrow S^n$, $O \mapsto O \cdot \mathbf{1}$. Then $p^{-1}(\mathbf{1}) \simeq \mathrm{SO}(n-1)$.

Lemma 14.10. *This map p defines a fiber bundle.*

This would imply that p satisfies the path and homotopy lifting property. To see the lemma we will appeal to the following theorem:

Theorem 14.11. *Let G be a Lie group. If H is a closed subgroup, then $p : G \rightarrow G/H$ defines a fiber bundle with fiber is H .*

We will prove this theorem in the next two lectures.

We will now consider an application of the above theorem. To do this we define a new type of space. Take $\lambda \in \mathbb{C} \setminus \{0\}$. Then the map $M_\lambda : z \mapsto \lambda z$ acts on $\mathbb{C}^n \setminus \{0\}$. Note that $M_\lambda(S^{2n+1}) = S^{2n+1}$ if $|\lambda| = 1$. Next, consider the relation $z \sim w$ if and only if there exists some $\lambda \in \mathbb{C} \setminus \{0\}$ such that $\lambda z = w$ on $\mathbb{C}^n \setminus \{0\}$. Then the **complex projective space** is defined as $\mathbb{C}P^n := (\mathbb{C}^n \setminus \{0\}) / \sim = S^{2n+1} / \sim$. Then this gets us a projection map $p : S^{2n+1} \rightarrow \mathbb{C}P^n$, where each fiber is S^1 , for the same reason as in the theorem.

Corollary 14.12. *We have a long exact sequence*

$$\cdots \rightarrow \pi_k(S^1) \rightarrow \pi_k(S^{2n+1}) \rightarrow \pi_k(\mathbb{C}P^n) \rightarrow \pi_{k-1}(S^1) \rightarrow \cdots$$

Thus if $k-1 \geq 2$, that is, $k \geq 3$, $\pi_k(S^{2n+1}) \simeq \pi_k(\mathbb{C}P^n)$. For $k=2$, we have $\pi_2(\mathbb{C}P^n) \simeq \mathbb{Z}$.

In general, $\pi_{2n+1}(\mathbb{C}P^n) \simeq \pi_{2n+1}(S^{2n+1}) \simeq \mathbb{Z}$. To answer the question posed in the previous lecture, we note that $\mathbb{C}P^\infty$ is the Eilenberg-MacLane space needed: it is the sought for $K(\mathbb{Z}, 2)$.

15 2/15/23

Today we will start the proof of a theorem that was stated and unproved in the last lecture. We will state it again for a reminder.

Theorem 15.1. *Let G be a Lie group and H a closed subgroup of G . Then the projection $p : G \rightarrow G/H$ defines a fiber bundle with fiber H . That is, for every $x \in G/H$, there exists an open neighborhood U of x and a homeomorphism $Q : U \times H \rightarrow p^{-1}(U)$ such that $p(Q(u, h)) = u$ for all $u \in U$.*

15.1 A Short Course in Lie Groups

Let G be a Lie group. Then we will denote $\text{Lie}(G) := \mathfrak{g}$ to be the tangent space at the identity. Then the **exponential map** $\exp : \text{Lie}(G) \rightarrow G$ is defined in the following manner: if $v \in T_e G$ and X is a left-invariant vector field of G such that $X(0) = v$, $\alpha : \mathbb{R} \rightarrow G$ the associated one-parameter subgroup with $\alpha'(0) = X(0)$, then $v \mapsto \alpha(1)$.

Example 15.2. As an example $\text{GL}(n, \mathbb{R})$ is a Lie group, and $\text{Lie}(\text{GL}(n, \mathbb{R})) = \text{Mat}_{n \times n}(\mathbb{R})$. Then the exponential map coincides with the usual matrix exponential: $\exp(A) = \sum_{n=0}^{\infty} \frac{A^n}{n!}$. Then $\exp(0) = \text{id}$ and $d(\exp)_0 : \text{Mat}_{n \times n}(\mathbb{R}) \rightarrow T_e \text{GL}(n, \mathbb{R}) = \text{Mat}_{n \times n}(\mathbb{R})$, where $d(\exp)_0 \equiv \text{id}_{\text{Lie}(G)}$.

Now by the inverse function theorem, \exp is a local diffeomorphism; hence we can use the exponential map for charts for G . Moreover, since G is a Lie group, left multiplication $L_g : G \rightarrow G$ and $x \mapsto gx$ is a diffeomorphism, so once we know open sets near the identity, we know it everywhere. Thus studying the Lie algebra lets us study Lie groups.

Exercise 15.3. Let G be a path-connected topological group. Then $\pi_1(G, e_0)$ is abelian.

Suppose $\text{Lie}(G) = E \oplus F$ a direct sum of vector spaces. Then one can use $\exp|_E$ and $\exp|_F$ to define a new map $r_{E,F} : \text{Lie}(G) \rightarrow G$, given by

$$v = v_E + v_F \mapsto \exp(v_E) \exp(v_F).$$

Warning. $\exp(x + y) \neq \exp(x) \exp(y)$ in general, unless G is abelian.

However, $d(r_{E,F})_0 = \text{id}$, so the inverse function theorem tells us that $r_{E,F}$ is a local diffeomorphism.

Theorem 15.4 (Closed Subgroup Theorem). *Every closed subgroup of a Lie group is an embedded submanifold, and hence a Lie group.*

Note that if H is a closed subgroup of G , then $\text{Lie}(H)$ is a vector subspace of $\text{Lie}(G)$, and we say that $\text{Lie}(H)$ is a Lie subalgebra. Moreover, $\exp(\text{Lie}(H))$ is a subgroup of H , and is the connected component of H at the identity, sometimes also called the **analytic subgroup** associated to $\text{Lie}(H)$.

Now let $T \subseteq \text{Lie}(G)$ be a vector subspace such that $T \oplus \text{Lie}(H) = \text{Lie}(G)$. Then from linear algebra, naturally we have that $\text{Lie}(G)/\text{Lie}(H) \simeq T$. Hence $(\text{Lie}(G)/\text{Lie}(H)) \times \text{Lie}(H) \simeq \text{Lie}(G)$. This sketches out the proof of a special case of Theorem 15.1, when G is a vector space. In particular, it proves the theorem for $\text{Lie}(G)$ the tangent space at the identity.

Next we will state a theorem from Lie theory, which we take as given:

Theorem 15.5. *There exists a neighborhood W of the identity $e \in G$ and a neighborhood $V \subseteq \text{Lie}(G)$ of 0 such that the following hold:*

- (i) $\exp : V \rightarrow W$ is a diffeomorphism.
- (ii) if $h \in H \cap W$, then $(\exp)^{-1}(h) \in \text{Lie}(H)$; equivalently, $H \cap W \subseteq H^0$, where H^0 denotes the connected component of H with the identity.

Example 15.6. Let $G = \mathbb{R}^2$, and $H = \{(t, n_1 + n_2\sqrt{2}) \mid t \in \mathbb{R}, n_1, n_2 \in \mathbb{Z}\}$. Then $H^0 = \{(t, 0) \mid t \in \mathbb{R}\}$. But for any open $W \subseteq G$ with $0 \in W$, $H \cap W$ is not a subset of H^0 , since H is not a closed subgroup of G . In fact, H is dense in G !

Exercise 15.7. Show that, in the above example, H^0 is a normal subgroup of H .

16 2/17/23

16.1 The Proof of the Main Theorem

Today we will prove Theorem 15.1. Our goal is to show that the projection $p : G \rightarrow G/H$ defines a fiber bundle. That is, we want to show that for all $x \in G/H$, there exists some U an open neighborhood of x and a homeomorphism $Q : U \times H \rightarrow p^{-1}(U)$ such that $p(Q(u, h)) = u$. That is, the following diagram commutes:

$$\begin{array}{ccccc} U \times H & \xrightarrow{Q} & p^{-1}(U) & \xhookrightarrow{\iota} & G \\ \downarrow p_1 & & & \swarrow p & \\ U & \xhookrightarrow{\iota} & G/H & & \end{array}$$

Exercise 16.1. Let S be the genus 2 surface. Then there is no nontrivial $p : S \rightarrow B$, where B is a topological space, such that S together with p defines a fiber bundle. That is, B cannot be the singleton set and B cannot be equal to S .

Let G be a Lie group and H a closed subgroup of G ; consider $\text{Lie}(H)$, and $T \subseteq \text{Lie}(G)$ such that $\text{Lie}(G) = T \oplus \text{Lie}(H)$. Take $\Sigma \subseteq T$ and $B_H \subseteq \text{Lie}(H)$ both open sets such that $\Sigma \oplus B_H \sim \Sigma \times B_H$ are open sets. After shrinking if necessary, WLOG assume that $\Sigma = -\Sigma$ and the map

$$\begin{aligned} r : \Sigma \times B_H &\rightarrow G \\ (s, v) &\mapsto \exp(s) \exp(v) \end{aligned}$$

is a local diffeomorphism (this is a consequence of the fact that \exp is a local diffeomorphism); let $W := r(\Sigma \times B_H) \subseteq G$, and note that this is a neighborhood of the origin.

Now define

$$\begin{aligned} \widehat{Q} : \exp(\Sigma) \times H &\rightarrow G \\ (\sigma, h) &\mapsto \sigma h. \end{aligned}$$

Lemma 16.2. Consider \widehat{Q} as above. Then:

- (a) \widehat{Q} is one-to-one.
- (b) \widehat{Q} is continuous.
- (c) \widehat{Q} is an open map.

Proof. To prove (a), suppose $\sigma_1 h_1 = \sigma_2 h_2$. Then rearranging, $\sigma_2^{-1} \sigma_1 = h_2 h_1^{-1}$. Clearly $h_2 h_1^{-1} \in H$. On the other hand, $\exp(0) = e$, so $e \in \exp(\Sigma)$; hence $h_2 h_1^{-1} = e \cdot (h_2 h_1^{-1}) \in W$. Thus applying Theorem 15.5, there exists some $v \in \text{Lie}(H)$ such that $h_2 h_1^{-1} = \exp(v)$. Let $s_1, s_2 \in \Sigma$ be such

that $\exp(s_1) = \sigma_1$ and $\exp(s_2) = \sigma_2$. Then we have $\sigma_1 = \sigma_2 \exp(v_2)$, and looking the definition of r again, $r(s_1, e) = r(s_2, v)$. Now using the fact that r and \exp are local diffeomorphisms, we conclude that $0 = v$ and $s_1 = s_2$, and hence $\sigma_1 = \sigma_2$ and $h_1 = h_2$. This shows that \widehat{Q} is one-to-one.

Now to prove (b), obviously \widehat{Q} is continuous since multiplication is a continuous operator, coming from the fact that G is a Lie group.

Finally, to prove (c), let $U_1 \subseteq \exp(\Sigma)$ be open and $U_2 \subseteq H$ be open, and further assume that $e \in U_2$. Then defining $\widehat{U}_1 := (\exp|_{\Sigma})^{-1}(U_1)$ and $\widehat{U}_2 := (\exp|_{B_H})^{-1}(U_2)$, we see that $Q(U_1, U_2) = r(\widehat{U}_1, \widehat{U}_2)$. Since \exp is a local diffeomorphism, we have \widehat{U}_1 and \widehat{U}_2 are both open; similarly, since r is a local diffeomorphism, $r(\widehat{U}_1, \widehat{U}_2)$ must also be open. This proves the claim for the single case where U_2 is a neighborhood around the identity.

Now to handle the general case, take U_1 open in $\exp(\Sigma)$ and U_2 open in H without the assumption that $e \in U_2$. Let h and \widetilde{U}_2 be defined as $U_2 = \widetilde{U}_2 \cdot h$, and \widetilde{U}_2 is a neighborhood of e ; that is, $\widetilde{U}_2 = R_{h^{-1}}(U_2)$. Then

$$\widehat{Q}(U_1, U_2) = U_1 \cdot U_2 = U_1 \cdot (\widetilde{U}_2 \cdot h) = \widehat{Q}(U_1, \widetilde{U}_2) \cdot h,$$

and since by the special case $\widehat{Q}(U_1, \widetilde{U}_2)$ is open and right multiplication is a diffeomorphism, we have that $\widehat{Q}(U_1, U_2)$ is open. ■

Corollary 16.3. *The image $\widehat{Q}(\exp(\Sigma) \times H)$ is open in G and \widehat{Q} is a homeomorphism onto its image.*

Now we return to studying p and the quotient.

Lemma 16.4. *Consider the restriction of the map p to $\exp(\Sigma)$, that is, $p : \exp(\Sigma) \rightarrow G/H$. Then:*

- (a) p is one-to-one,
- (b) p is continuous,
- (c) p is an open map.

Proof. To prove (a), suppose that $p(\sigma_1) = p(\sigma_2)$ for some $\sigma_1, \sigma_2 \in \exp(\Sigma)$. Then by definition, $\sigma_1 H = \sigma_2 H$. Then there exist $h_1, h_2 \in H$ such that $\sigma_1 h_1 = \sigma_2 h_2$, that is, $\widehat{Q}(\sigma_1, h_1) = \widehat{Q}(\sigma_2, h_2)$. But by Lemma 16.2 we see that $\sigma_1 = \sigma_2$.

Statement (b) is trivial: this is the restriction of a continuous map.

Finally, to prove (c), take any open set $U \subseteq \exp(\Sigma)$. Note that $p(U)$ is open in G/H if and only if $p^{-1}(p(U))$ is open in G , by the definition of quotient topology. But $p^{-1}(p(U)) = UH = \widehat{Q}(U \times H)$, which is open again by Lemma 16.2. ■

Corollary 16.5. *The image $p(\exp(\Sigma))$ is open in G/H and $p|_{\exp(\Sigma)}$ is a homeomorphism onto its image.*

Now we finish the proof of Theorem 15.1. Let $A := p(\exp(\Sigma))$. Then A is open by the previous corollary. Define

$$Q : A \times H \rightarrow p^{-1}(A),$$

$$(a, h) \mapsto (p|_{\exp(\Sigma)})^{-1}(a) \cdot h = \widehat{Q}\left((p|_{\exp(\Sigma)})^{-1}(a), h\right).$$

Then now defining

$$f : A \times H \rightarrow \exp(\Sigma) \times H,$$

$$(a, h) \mapsto \left((p|_{\exp(\Sigma)})^{-1}(a), h\right),$$

we have that $Q = \widehat{Q} \circ f$; if you like commutative diagrams, this means that the following diagram commutes:

$$\begin{array}{ccc} A \times H & \xrightarrow{f} & \exp(\Sigma) \times H \\ & \searrow Q & \downarrow \widehat{Q} \\ & & p^{-1}(A) \end{array}$$

By Corollary 16.5 f is a homeomorphism, and hence Q is a homeomorphism. And indeed, for $(a, h) \in A \times H$,

$$\begin{aligned} (p \circ Q)(a, h) &= p\left(\widehat{Q}\left((p|_{\exp(\Sigma)})^{-1}(a), h\right)\right) \\ &= p\left((p|_{\exp(\Sigma)})^{-1}(a) \cdot h\right) \\ &= a. \end{aligned}$$

Finally, noting that left multiplication is a diffeomorphism from G onto itself, we can translate any neighborhood about the identity and functions on it diffeomorphically. Hence this defines a fiber bundle, as desired. \blacksquare

Remark 16.6. We did not really use the full power of G being a Lie group, but we only used the local structure: that is, the existence of $W \subseteq G$ a neighborhood and a homeomorphism to $\Sigma \times B_H \rightarrow W$, where $e \in \Sigma \subseteq G$, and B_H also containing e . Then if G was any topological group with the above property, the above proof would have gone through.

In fact, we can prove the following theorem for topological groups using similar ideas as above:

Definition 16.7. A subgroup $H \leq G$ is **discrete** if $\{e\}$ is an open set in H .

Theorem 16.8. If H is discrete, then $p : G \rightarrow G/H$ is a covering map.

17 2/20/23

Today we start discussing homology. First we will approach it axiomatically.

17.1 The Eilenberg-Steenrod Axioms

To any topological pair (X, A) , we associate a sequence of abelian groups $\{H_i(X, A)\}_{i \in \mathbb{Z}}$ and a sequence of homomorphisms $\partial : H_i(X, A) \rightarrow H_{i-1}(A, \emptyset)$, called the **boundary map**.

Notation. We will denote $H_i(X, \emptyset) = H_i(X)$ for brevity.

To each continuous map $f : (X, A) \rightarrow (Y, B)$, we associate a family of homomorphisms $\{f_i : H_i(X, A) \rightarrow H_i(Y, B)\}_{i \in \mathbb{Z}}$. Then we have a **homology theory** if the following axioms are satisfied:

(H1) if $f : (X, A) \rightarrow (X, A)$ is the identity map, then $f_i \equiv \text{id}_{H_i(X, A)}$ for each i .

(H2) if $f : (X, A) \rightarrow (Y, B)$ and $g : (Y, B) \rightarrow (Z, C)$ then $(g \circ f)_i = g_i \circ f_i$ for each i .

(H3) If $f : (X, A) \rightarrow (Y, B)$ then the following diagram commutes:

$$\begin{array}{ccc} H_i(X, A) & \xrightarrow{f_i} & H_i(Y, B) \\ \downarrow \partial & & \downarrow \partial \\ H_{i-1}(A) & \xrightarrow{(f|_A)_{i-1}} & H_{i-1}(B) \end{array}$$

(H4) Letting $\iota : (A, \emptyset) \hookrightarrow (X, \emptyset)$ and $J : (X, \emptyset) \hookrightarrow (X, A)$ be the inclusion maps, the following sequence is exact:

$$\cdots \rightarrow H_i(A) \xrightarrow{\iota_i} H_i(X) \xrightarrow{J_i} H_i(X, A) \xrightarrow{\partial_i} H_{i-1}(A) \rightarrow \cdots$$

(H5) If $f : (X, A) \rightarrow (Y, B)$ and $g : (X, A) \rightarrow (Y, B)$ are homotopic, then $f_i = g_i$ for all i .

(H6) If $U \subseteq A$ with $\overline{U} \subseteq \text{int } A$, then $\iota_i : H_i(X \setminus U, A \setminus U) \rightarrow H_i(X, A)$ is an isomorphism for all i , where $\iota : (X \setminus U, A \setminus U) \hookrightarrow (X, A)$ is the inclusion map.

(Warning.) It is not true in general that $H_i(X, A) \simeq H_i(X \setminus A)$.

(H7) If $X = \{x_0\}$, then $H_i(X) = 0$ for all $i \neq 0$, and $H_0(X)$ is called the **coefficient group**.

Remark 17.1. Often times we would have $H_0(X) \simeq \mathbb{Z}$.

Lemma 17.2. If $f : (X, A) \rightarrow (Y, B)$ is a homotopy equivalence, then $f_i : H_i(X, A) \rightarrow H_i(Y, B)$ is an isomorphism for all i .

Proof. If f is a homotopy equivalence, there is a $g : (Y, B) \rightarrow (X, A)$ such that $g \circ f$ is homotopic to $\text{id}_{(X, A)}$. Then by Axiom (H5), $(g \circ f)_i : H_i(X, A) \rightarrow H_i(X, A)$ is equal to $\text{id}_{H_i(X, A)}$ for all i . But by Axiom (H2), $g_i \circ f_i = \text{id}_{H_i(X, A)}$. By the same argument, *mutatis mutandis*, we have $f_i \circ g_i = \text{id}_{H_i(Y, B)}$. Hence f_i is invertible for each i , and so must be an isomorphism. \blacksquare

Definition 17.3. Given a pair of subsets U and A such that $U \subseteq A \subseteq X$, we say that the inclusion $\iota : (X \setminus U, A \setminus U) \hookrightarrow (X, A)$ is **excisive** if it induces isomorphisms in homology.

Equivalently, given X and a subset $A \subseteq B \subseteq X$, and $A \subseteq C$, we say that the inclusion of pairs $\iota : (B, A) \hookrightarrow (X, C)$ is excisive if $\iota_i : H_i(B, A) \rightarrow H_i(X, C)$ is an isomorphism for all i . To see the equivalence, consider $B = U^c$ and set $C = A$. Using this new definition, we can reformulate Axiom (H6) in the following way: if $\overline{U} \subseteq \text{int } A$, the inclusion $J : (X \setminus U, A \setminus U) \hookrightarrow (X, A)$ is excisive.

Definition 17.4. If $A, B \subseteq X$, then we say that A and B are an **excisive pair** if the inclusion $\iota : (B, A \cap B) \hookrightarrow (A \cup B, A)$ is excisive.

Definition 17.5. A **generalized homology theory** is a theory satisfying Axioms (H1) through (H6).

Definition 17.6. Let

$$0 \rightarrow M_1 \rightarrow N \rightarrow M_2 \rightarrow 0$$

be a short exact sequence. We say that this short exact sequence **splits** if there is an isomorphism $\varphi : N \rightarrow M_1 \oplus M_2$ such that the following diagram commutes:

$$\begin{array}{ccccccc} 0 & \longrightarrow & M_1 & \longrightarrow & N & \longrightarrow & M_2 \longrightarrow 0 \\ & & \downarrow \text{id} & & \downarrow \varphi & & \downarrow \text{id} \\ 0 & \longrightarrow & M_1 & \xrightarrow{\iota} & M_1 \oplus M_2 & \xrightarrow{p} & M_2 \longrightarrow 0 \end{array}$$

where $\iota : M_1 \rightarrow M_1 \oplus M_2$ is natural injection and $p : M_1 \oplus M_2 \rightarrow M_2$ is the projection onto the second factor.

Lemma 17.7 (Splitting Lemma). *Suppose we have a short exact sequence*

$$0 \rightarrow M_1 \xrightarrow{r} N \xrightarrow{s} M_2 \rightarrow 0.$$

Then the following are equivalent:

- (a) r has a left-inverse.
- (b) s has a right-inverse.
- (c) This sequence splits.

Lemma 17.8. (a) $H_i(\emptyset) = 0$ for all i .

- (b) If $A \subseteq X$ is a retract of X , that is, there exists some continuous $r : X \rightarrow A$ such that $r \circ \iota = \text{id}_A$, then $\iota_i : H_i(A) \rightarrow H_i(X)$ is one-to-one and $r_i : H_i(X) \rightarrow H_i(A)$ is onto for all i . Moreover, $H_i(X) \simeq H_i(A) \oplus H_i(X, A)$ and $H_i(X, A) \simeq \ker r_i$.
- (c) Let $X = X_1 \sqcup X_2$ be a disjoint union of X_1 and X_2 , with X_1 and X_2 open. Then $J^l : H_i(X_l) \rightarrow H_i(X)$ are one-to-one for each l and $(J^1 + J^2)_i : H_i(X_1) \oplus H_i(X_2) \rightarrow H_i(X)$ is an isomorphism.

Proof. Given in class as an exercise, but we will prove it.

To see part (a), note that $\emptyset \subseteq \{x_0\}$, and \emptyset is open. So by Axiom (H4) and (H7) we have the following long exact sequence:

$$\cdots \rightarrow H_i(\emptyset) \xrightarrow{\iota_i} \underbrace{H_i(\{x_0\})}_{=0} \xrightarrow{J_i} \underbrace{H_i(\{x_0\}, \emptyset)}_{=0} \xrightarrow{\partial} H_{i-1}(\emptyset) \rightarrow \cdots,$$

from which it is immediate.

To see part (b), suppose $A \subseteq X$ is as prescribed. Then as a consequence of Axiom (H1) and (H3), $r_i \circ \iota_i = \text{id}_{H_i(A)}$ for each i . This proves the first portion. Next, consider the exact sequence given by Axiom (H4). Since ι_i is injective for each i , by exactness $0 = \ker \iota_{i-1} = \text{im } \partial_i$. Thus ∂_i is the trivial

map, and so $H_i(X, A) = \ker \partial_i = \operatorname{im} J_i$, and so J_i is surjective for each i . Then this implies that we have the following short exact sequence:

$$0 \rightarrow H_i(A) \xrightarrow{\iota_i} H_i(X) \xrightarrow{J_i} H_i(X, A) \rightarrow 0.$$

Then appealing to the Splitting Lemma we deduce that $H_i(X) \simeq H_i(A) \oplus H_i(X, A)$. On the other hand, again by the Splitting Lemma the following diagram commutes:

$$\begin{array}{ccc} H_i(A) & \xrightarrow{\iota_i} & H_i(X) \\ \downarrow \text{id} & & \downarrow \varphi \\ H_i(A) & \xrightarrow{\iota} & H_i(A) \oplus H_i(X, A) \end{array}$$

where $\varphi : H_i(X) \rightarrow H_i(A) \oplus H_i(X, A)$ is an isomorphism. Then defining $p_1 : H_i(A) \oplus H_i(X, A) \rightarrow H_i(A)$ to be the projection onto the first factor, we see that $p_1 = r_i \circ \varphi^{-1}$, and since φ^{-1} is an isomorphism $\ker p_1 \simeq \ker r_i$, but $\ker p_1 \simeq H_i(X, A)$. This proves (b).

To prove (c), fix $x_0 \in X_1$ and consider the map $r : X \rightarrow X_1$ given by

$$r(x) = \begin{cases} x, & x \in X_1 \\ x_0, & x \in X_2. \end{cases}$$

Then r is continuous since any open set $U \subseteq X_1$ not containing x_0 has preimage U , and if $x_0 \in U$ then $r^{-1}(U) = U \cup X_2$ which is open. Moreover, if $J^1 : X_1 \hookrightarrow X$ is the inclusion then $r \circ J^1 = \operatorname{id}_A$, so part (b) we have $J^1 : H_i(X_1) \rightarrow H_i(X_2)$ is one-to-one for each i . Furthermore, $H_i(X) \simeq H_i(X_1) \oplus H_i(X, X_1)$. But by Axiom (H6), we have the inclusion $J^2 : (X_2, \emptyset) = (X \setminus X_1, X_1 \setminus X_1) \hookrightarrow (X, X_1)$ is excisive so $H_i(X_2) \simeq H_i(X, X_1)$ for each i via this isomorphism. Hence $J^1 + J^2$ induces the desired isomorphism. ■

17.2 Some Foreshadowing

Our goal is to show next Monday that $H_n(S^n) \simeq H_0(S^n) \simeq \mathbb{Z}$ for all n . This would imply that by Lemma 17.2 that S^n is not contractible. Therefore, $\pi_n(S^n, e_0) \neq 0$, for $\operatorname{id}_{S^n} \in \pi_n(S^n, e_0)$ but is nontrivial. We are also going to prove Invariance of Domain: that if $\mathbb{R}^n \approx \mathbb{R}^k$, then $n = k$, and if $f : U \rightarrow \mathbb{R}^n$ is continuous and one-to-one with $U \subseteq \mathbb{R}^n$ open, then $f(U)$ is open in \mathbb{R}^n .

18 2/22/22

18.1 Reduced Homology Groups

Definition 18.1. Denote $\tilde{H}_i(X) := \ker(f_i : H_i(X) \rightarrow H_i(x_0))$ where $f(X) = x_0$. Then $\tilde{H}_i(X)$ are called the **reduced homology groups**.

Observation. Note that by Axiom (H7), $H_i(x_0) = 0$ for all $i \neq 0$, so by definition $\tilde{H}_i(X) = H_i(X)$ for all $i \neq 0$. Thus $\tilde{H}_0(X)$ is the only group that could be different from H_0 .

Theorem 18.2. (a) For all $x_0 \in X$, $\tilde{H}_i(X) \simeq H_i(X, x_0)$.

$$(b) \ H_i(X) \simeq H_i(x_0) \oplus \tilde{H}_i(X).$$

Proof. Note that $f : X \rightarrow x_0$ given by $f(x) = x_0$ is a retract of X . Then applying Lemma 17.6 part (b), we conclude that $H_i(X) \simeq H_i(x_0) \oplus H_i(X, x_0)$ with $H_i(X, x_0) \simeq \ker f_i = \tilde{H}_i(X)$. ■

Note that in part (b) of this theorem, $H_0(X) = H_0(x_0) \oplus \tilde{H}_0(X)$ is the only interesting part.

Corollary 18.3. *If $f : X \rightarrow Y$, then there exists some $f_i : \tilde{H}_i(X) \rightarrow \tilde{H}_i(Y)$.*

18.2 Mayer-Vietoris Sequence

Theorem 18.4 (Mayer-Vietoris). *Let X be a space, and assume that $X = X_1 \cup X_2$, with $X_1, X_2 \subseteq X$, and let $A := X_1 \cap X_2$. Assume further that $(X_1, A) \hookrightarrow (X, X_2)$ is excisive. Then we have the following long exact sequence:*

$$\cdots \rightarrow H_i(A) \xrightarrow{\alpha} H_i(X_1) \oplus H_i(X_2) \xrightarrow{\beta} H_i(X) \xrightarrow{\Delta} H_{i-1}(A) \rightarrow \cdots. \quad (18.5)$$

Remark 18.6. Leopold Vietoris, one of the people for whom this theorem is named, lived until he was 110 years old.

We will finish the proof in the next lecture, and just outline the main ideas for the proof. Let $\alpha_j : A \hookrightarrow X_j$, $\beta_j : X_j \hookrightarrow X$ be the inclusion maps, and also let $\gamma_1 : (X_1, \emptyset) \hookrightarrow (X, A)$ and $\gamma_2 : (X, \emptyset) \hookrightarrow (X, X_2)$ also be the inclusion maps. Then Axiom (H4) gives us exact sequences

$$\cdots \rightarrow H_i(A) \xrightarrow{\alpha_1} H_i(X_1) \xrightarrow{\gamma_1} H_i(X_1, A) \xrightarrow{\partial_1} H_{i-1}(A) \rightarrow \cdots$$

and

$$\cdots \rightarrow H_i(X_2) \xrightarrow{\beta_2} H_i(X) \xrightarrow{\gamma_2} H_i(X, X_2) \xrightarrow{\partial_2} H_{i-1}(X_2) \rightarrow \cdots.$$

By the excision axiom (H6) we have the existence of an isomorphism $\varepsilon : H_i(X_1, A) \rightarrow H_i(X, X_2)$. Then we can link them up so that the following diagram commutes, by Axiom (H3) and the observation that $\alpha_j, \beta_j, \gamma_j$, and ε are all induced by inclusion maps:

$$\begin{array}{ccccccc} \cdots & \longrightarrow & H_i(A) & \xrightarrow{\alpha_1} & H_i(X_1) & \xrightarrow{\gamma_1} & H_i(X_1, A) & \xrightarrow{\partial_1} & H_{i-1}(A) & \longrightarrow & \cdots \\ & & \downarrow \alpha_2 & & \downarrow \beta_1 & & \downarrow \varepsilon & & \downarrow \alpha_2 & & \\ \cdots & \longrightarrow & H_i(X_2) & \xrightarrow{\beta_2} & H_i(X) & \xrightarrow{\gamma_2} & H_i(X, X_2) & \xrightarrow{\partial_2} & H_{i-1}(X_2) & \longrightarrow & \cdots \end{array}$$

Now we want to make the sequence in (18.5) exact using the α_j and β_j . First, take the obvious map $\beta : H_i(X_1) \oplus H_i(X_2) \rightarrow H_i(X)$ via $\beta(v_1 + v_2) = \beta_1(v_1) + \beta_2(v_2)$. Similarly, the temptation is to define α as $\alpha_1 + \alpha_2$, but this will not guarantee exactness. Thus we define $\alpha : H_i(A) \rightarrow H_i(X_1) \oplus H_i(X_2)$ via $\alpha(v) = \alpha_1(v) - \alpha_2(v)$, so that $\beta \circ \alpha = 0$. Indeed,

$$\beta(\alpha(v)) = \beta(\alpha_1(v) - \alpha_2(v)) = \beta_1(\alpha_1(v)) - \beta_2(\alpha_2(v)) = 0$$

where the last equality is by the commutativity of the diagram. Hence $\text{im } \alpha \subseteq \ker \beta$. Now the next step is to define the map Δ , which we will postpone until the next time.

19 2/24/23

Today we will finish up the proof of Theorem 18.4.

19.1 Mayer-Vietoris Sequence, Part Two

Now it is time to define Δ in (18.5), and to this end we will refer back to the diagram we drew in the last lecture. Notice how $H_i(X)$ fits in the bottom row of the diagram, and we can get to $H_{i-1}(A)$ by defining $\Delta : H_i(X) \rightarrow H_{i-1}(A)$ by $\Delta := \partial_1 \circ \varepsilon^{-1} \circ \gamma_2$, which is well-defined since ε is an isomorphism and so has a defined inverse.

We now verify that defining Δ this way gives us an exact sequence. First we will show that $\text{im } \beta = \ker \Delta$. First suppose that $v \in \text{im } \beta$, that is, $v = \beta(w_1 + w_2) = \beta_1(w_1) + \beta_2(w_2)$ for $w_1 + w_2 \in H_i(X_1) \oplus H_i(X_2)$. Then

$$\gamma_2(v) = \gamma_2(\beta_1(w_1)) + \underbrace{\gamma_2(\beta_2(w_2))}_{=0} = \varepsilon(\gamma_1(w_1)),$$

where the first equality is by the exactness of the sequence and the second equality is by the commutativity of the diagram. Thus rearranging, $\gamma_1(w_1) = (\varepsilon^{-1} \circ \gamma_2)(v)$. Then $\Delta(v) = (\partial_1 \circ \varepsilon^{-1} \circ \gamma_2)(v) = (\partial_1 \circ \gamma_1)(w_1) = 0$ by exactness. Hence $v \in \ker \Delta$.

Conversely, suppose that $v \in \ker \Delta$. Then $(\partial_1 \circ \varepsilon^{-1} \circ \gamma_2)(v) = 0$, and so $(\varepsilon^{-1} \circ \gamma_2)(v) \in \ker \partial_1 = \text{im } \gamma_1$ by exactness. So there exists some $v_1 \in H_i(X_1)$ such that

$$(\varepsilon^{-1} \circ \gamma_2)(v) = \gamma_1(v_1). \quad (19.1)$$

Define $\widehat{v}_2 := v - \beta_1(v_1)$. Then by construction we have $v = \beta_1(v_1) + \widehat{v}_2$. To show that $v \in \text{im } \beta$, we need to show that $\widehat{v}_2 = \beta(v_2)$ for some $v_2 \in H_i(X_2)$: that is, $\widehat{v}_2 \in \text{im } \beta = \ker \gamma_2$. So it suffices to show that $\gamma_2(\widehat{v}_2) = 0$. Indeed,

$$\gamma_2(\widehat{v}_2) = \gamma_2(v) - \gamma_2(\beta_1(v_1)) = \gamma_2(v) - \varepsilon(\gamma_1(v_1)) = 0,$$

where the second equality is by commutativity of the diagram and the last inequality is by (19.1).

Next, we will verify that $\text{im } \alpha = \ker \beta$. One direction has already been proven in the last section. Suppose that $v_1 + v_2 \in \ker \beta$. Then $\beta(v_1 + v_2) = \beta_1(v_1) + \beta_2(v_2) = 0$, and so $\beta_2(v_2) = -\beta_1(v_1)$. Then $\gamma_2(\beta_2(v_2)) = 0 = \gamma_2(\beta_1(v_1))$. But by commutativity of the diagram, $\gamma_2(\beta_1(v_1)) = \varepsilon(\gamma_1(v_1))$, but since ε is an isomorphism, this implies that $v_1 \in \ker \gamma_1 = \text{im } \alpha_1$. Hence there exists some $w_1 \in H_i(A)$ such that $v_1 = \alpha_1(w_1)$. Now by the commutativity of the diagram, $(\beta_2 \circ \alpha_2)(w_1) = (\beta_1 \circ \alpha_1)(w_1) = -\beta_2(v_2)$ and so $\beta_2(v_2 + \alpha_2(w_1)) = 0$, that is, $v_2 + \alpha_2(w_1) \in \ker \beta_2 = \text{im } \partial_2$. Then there exists some $u \in H_{i+1}(X, X_2)$ such that $\partial_2(u) = v_2 + \alpha_2(w_1)$. Now since ε is an isomorphism, consider $w_2 := (\partial_1 \circ \varepsilon^{-1})(u)$. Then

$$\alpha_2(w_2) = (\alpha_2 \circ \partial_1 \circ \varepsilon^{-1})(u) = (\partial_2 \circ \varepsilon \circ \varepsilon^{-1})(u) = \partial_2(u) = v_2 + \alpha_2(w_1).$$

Now, noting that $w_2 \in \text{im } \partial_1 = \ker \alpha_1$, we have

$$\alpha_1(w_1 - w_2) = \alpha_1(w_1) - \alpha_2(w_2) = v_1,$$

and

$$\alpha_2(w_1 - w_2) = \alpha_2(w_1) - (v_2 + \alpha_2(w_1)) = -v_2.$$

Thus we conclude that

$$\alpha(w_1 - w_2) = \alpha_1(w_1 - w_2) - \alpha_2(w_1 - w_2) = v_1 + v_2,$$

as needed.

The final step is to show that $\text{im } \Delta = \ker \alpha$. Suppose that $\alpha(r) = 0$ for some $r \in H_i(A)$. Then $\alpha_1(r) = \alpha_2(r) = 0$. But by exactness $\ker \alpha_1 = \text{im } \partial_1$ and so there exists some $s \in H_{i+1}(X_1, A)$ such that $\partial_1(s) = r$. Now by the commutativity of the diagram, we have

$$0 = \alpha_2(r) = (\alpha_2 \circ \partial_1)(s) = (\partial_2 \circ \varepsilon)(s).$$

Hence $\varepsilon(s) \in \ker \partial_2 = \text{im } \gamma_2$. Hence there exists some $q \in H_i(X)$ such that $\gamma_2(q) = \varepsilon(s)$. Now observe:

$$\Delta(q) = (\partial_1 \circ \varepsilon^{-1} \circ \gamma_2)(q) = (\partial_1 \circ \varepsilon^{-1})(\varepsilon(s)) = \partial_1(s) = r,$$

so $r \in \text{im } \Delta$.

Conversely, let $r \in \text{im } \Delta$, so $r = \Delta(q)$ for some $q \in H_{i+1}(X)$. Then $\alpha_1(r) = (\alpha_1 \circ \partial_1 \circ \varepsilon^{-1} \circ \gamma_2)(q) = 0$ since $\alpha_1 \circ \partial_1 = 0$ by exactness. On the other hand,

$$\alpha_2(r) = (\alpha_2 \circ \partial_1 \circ \varepsilon^{-1} \circ \gamma_2)(q) = (\partial_2 \circ \gamma_2)(q) = 0$$

by the commutativity of the diagram. Hence $\alpha(r) = \alpha_1(r) - \alpha_2(r) = 0$, concluding the proof of exactness. \blacksquare

19.2 Consequences of the Mayer-Vietoris Sequence


Corollary 19.2 (Mayer-Vietoris Sequence for Reduced Homology Groups). *Assume the same hypothesis as in Theorem 18.4 with the addition that $A = X_1 \cap X_2 \neq \emptyset$. Then we have the long exact sequence*

$$\cdots \rightarrow \tilde{H}_i(A) \xrightarrow{\alpha} \tilde{H}_i(X_1) \oplus \tilde{H}_i(X_2) \xrightarrow{\beta} \tilde{H}_i(X) \xrightarrow{\Delta} \tilde{H}_{i-1} \rightarrow \cdots$$

Remark 19.3. Recall that $\tilde{H}_i(X) \leq H_i(X)$ for each i , and moreover the Mayer-Vietoris sequence for reduced homology groups is identical to the usual Mayer-Vietoris sequence, except for H_0 .

Proposition 19.4. *Let A be a nonempty closed subset of X , and suppose that ∂A has an open neighborhood $C \subseteq X \setminus \text{int } A$ such that the inclusions $A \hookrightarrow C \cup A$ and $\partial A \hookrightarrow C$ are both homotopy equivalences. Then $(X \setminus \text{int } A, \partial A) \hookrightarrow (X, A)$ is excisive.*

Proof. Since A is closed subset of $A \cup C$, we see that by Axiom (H6), the inclusion $(X \setminus \text{int } A, (C \cup A) \setminus \text{int } A) \hookrightarrow (X, C \cup A)$ is excisive. Now $(C \cup A) \setminus \text{int } A = C$, so now we have $(X \setminus \text{int } A, C) \hookrightarrow (X, C \cup A)$ is excisive; applying the hypothesis, the homotopy equivalences induce isomorphisms of homology groups. Thus $(X \setminus \text{int } A, \partial A) \hookrightarrow (X, A)$ is excisive.





Also recall that the suspension of a space was defined in lecture 7; if X is a space, then its suspension ΣX was the space $X \times [-1, 1]$ quotiented by identifying $X \times \{-1\}$ with a point c_- and $X \times \{1\}$ with another point c_+ .

Theorem 19.5 (Suspension Theorem). *Let X be a space.*


- (a) *There exists an isomorphism $\Sigma : \tilde{H}_{i+1}(\Sigma X) \rightarrow \tilde{H}_i(X)$.*
- (b) *If $f : X \rightarrow Y$, then we can define $\Sigma f : \Sigma X \rightarrow \Sigma Y$, where $\Sigma[x, t] = [f(x), t]$ where $[x, t]$ is the equivalence class under the quotient defining the suspension, such that Σf is well-defined and the following diagram commutes:*

$$\begin{array}{ccc} \tilde{H}_{i+1}(\Sigma X) & \xrightarrow{(\Sigma f)_i} & \tilde{H}_{i+1}(\Sigma Y) \\ \downarrow \Sigma & & \downarrow \Sigma \\ \tilde{H}_i(X) & \xrightarrow{f_i} & \tilde{H}_i(Y) \end{array}$$

Proof. Let $\epsilon > 0$ be small, and define $X_1 := X \times [-1, 0]/(X \times \{-1\} \sim c_-)$, $X_2 := X \times [0, 1]/(X \times \{1\} \sim c_+)$. Then $A = X_1 \cap X_2 = X \times \{0\}$ which is homotopy equivalent to X in the obvious way. Now if the hypotheses for the Mayer-Vietoris sequence held, then we would have the following exact sequence:

$$\begin{array}{ccccccc} \cdots & \longrightarrow & \tilde{H}_{i+1}(X \times \{0\}) & \longrightarrow & \tilde{H}_{i+1}(X_1) \oplus \tilde{H}_{i+1}(X_2) & \longrightarrow & \tilde{H}_{i+1}(\Sigma X) \\ & & \searrow & & \searrow & & \searrow \\ & & \tilde{H}_i(X \times \{0\}) & \longrightarrow & \tilde{H}_i(X_1) \oplus \tilde{H}_i(X_2) & \longrightarrow & \tilde{H}_i(\Sigma X) \longrightarrow \cdots \end{array}$$

But since X_1 and X_2 are both contractible, by Lemma 17.2 we have that $\tilde{H}_i(X_1) = \tilde{H}_i(X_2) = 0$ for all i . Therefore this implies that $\tilde{H}_{i+1}(\Sigma X) \simeq \tilde{H}_i(X \times \{0\}) \simeq \tilde{H}_i(X)$. Thus it suffices to verify excision; to do this, we will appeal to Proposition 19.4: indeed, X_2 is a closed subspace of ΣX , $\partial X_2 = X \times \{0\} = X_1 \cap X_2$, and $X \setminus \text{int } X_2 = X_1$. Let $\epsilon > 0$, and let $C := X \times (-\epsilon, 0]$ as in the figure below.



Then clearly $C \cup X_2$ deformation retracts onto X_2 and C deformation retracts onto $X_1 \cap X_2$; hence the inclusions are homotopy equivalences. We conclude that $(X_1, X_1 \cap X_2) \hookrightarrow (X, X_2)$ is excisive, as desired. ■

20 2/27/23

20.1 Consequences of the Suspension Theorem

Theorem 20.1. *The homology groups of S^n are given as follows:*

$$H_i(S^n) = \begin{cases} H_0(x_0), & n \neq 0, i = n \text{ or } i = 0 \\ H_0(x_0) \oplus H_0(x_0), & i = n = 0 \\ 0 & \text{otherwise,} \end{cases}$$

and the reduced homology groups of S^n are

$$\tilde{H}_i(S^n) = \begin{cases} H_0(x_0), & i = n \\ 0 & \text{otherwise.} \end{cases}$$

Proof. Recall that $\Sigma S^n \approx S^{n+1}$. Then by Theorem 19.5 and using induction, we have $\tilde{H}_{i+k}(\Sigma^k X) \simeq \tilde{H}_i(X)$ for a space X , and hence $\tilde{H}_{i+k}(S^k) \simeq \tilde{H}_i(S^0)$. But $S^0 = \{-1, 1\}$, and so by Lemma 17.8 we have $H_i(S^0) \simeq H_i(x_0) \oplus H_i(x_0)$, which is 0 unless $i = 0$, and $H_0(S^0) \simeq H_0(x_0) \oplus H_0(x_0)$. Hence $\tilde{H}_i(S^0) = 0$ if $i \neq 0$. On the other hand, $\tilde{H}_0(S^0) = \ker(f_i : H_0(S^0) \rightarrow H_0(x_0)) \simeq H_0(x_0)$.

Now if $n \neq 0$, we observe that

$$\tilde{H}_{i+n}(S^n) \simeq \tilde{H}_i(S^0) = \begin{cases} 0, & i \neq 0 \\ H_0(x_0), & i = 0. \end{cases}$$

From here we easily deduce the statement of the theorem for reduced homology groups. Now in order to compute the nonreduced homology groups, we only need to worry about the case $i = 0$. We only consider $f : H_0(S^n) \rightarrow H_0(x_0)$ for $n \neq 0$ since the $n = 0$ has been computed above; this map is one-to-one since the reduced homology group $\tilde{H}_0(S^n) = 0$ and this is the kernel of f . On the other hand, the singleton point is a retract of every space: if $\iota : \{x_0\} \hookrightarrow S^n$ is the inclusion, then $f \circ \iota = \text{id}_{\{x_0\}}$. Hence by Lemma 17.8, $f : H_n(S^n) \rightarrow H_n(x_0)$ is onto. Hence f must be an isomorphism. ■

Theorem 20.2 (Brouwer Fixed Point Theorem). *If $f : \overline{B_n} \rightarrow \overline{B_n}$ is a continuous map, where $\overline{B_n} \subseteq \mathbb{R}^n$ is the closed n -dimensional ball, then there exists some $x_0 \in \overline{B_n}$ such that $f(x_0) = x_0$.*

Proof. The proof mirrors that of the strategy of what we did when we proved this for $n = 2$, *mutatis mutandis*. The only thing that changes is the following proposition. ■

Theorem 20.3 (Brouwer No-Retract Theorem). *There is no retraction $r : \overline{B_n} \rightarrow S^{n-1}$.*

Proof. Towards a contradiction suppose there is a retract $r : \overline{B_n} \rightarrow S^{n-1}$, so if $\iota : S^{n-1} \hookrightarrow \overline{B_n}$ is the inclusion, then $r \circ \iota = \text{id}_{S^{n-1}}$. Then by Lemma 17.8, $r_i : H_i(\overline{B_n}) \rightarrow H_i(S^{n-1})$ is onto. But if $i = n - 1$, then $\text{im } r_{n-1} = H_{n-1}(S^{n-1}) \simeq H_0(x_0) \neq 0$. But on the other hand $\overline{B_n}$ is contractible, so $H_{n-1}(\overline{B_n}) \simeq H_{n-1}(x_0)$, which is trivial unless $n = 1$, a contradiction. If $n = 1$, then $H_0(\overline{B_n}) \simeq H_0(x_0)$ and $H_0(S^n) \simeq H_0(x_0) \oplus H_0(x_0)$, which also yields a contradiction. ■

Theorem 20.4 (Invariance of Dimension). *If $h : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a homeomorphism, then $n = m$.*

Proof. We have that $\mathbb{R}^n \setminus \{0\}$ is homeomorphic to $\mathbb{R}^m \setminus \{0\}$, via the map $h|_{\mathbb{R}^n \setminus \{0\}}$. But S^{n-1} is homotopy equivalent to $\mathbb{R}^n \setminus \{0\}$ via a deformation retract. Then we have the following isomorphisms of homology groups for all i :

$$H_i(S^{n-1}) \simeq H_i(\mathbb{R}^n \setminus \{0\}) \simeq H_i(\mathbb{R}^m \setminus \{0\}) \simeq H_i(S^{m-1}).$$

Thus it follows that $n = m$ by Theorem 20.1. ■

Remark 20.5. Using excision, we can extend the above result to hold for any open sets: that is, if $U \subseteq \mathbb{R}^n$ and $V \subseteq \mathbb{R}^m$ and $f : U \rightarrow V$ is a homeomorphism, then $m = n$.

21 3/1/22

21.1 Invariance of Domain

Recall in the last lecture we proved the invariance of dimension theorem, Theorem 20.4. In this lecture we will go a step further and generalize the result to arbitrary topological manifolds.

Definition 21.1. A **topological manifold** is a topological space X such that for all $x \in X$, there is an open neighborhood U containing x such that U is homeomorphic to an open subset $V \subseteq \mathbb{R}^n$.

Theorem 21.2 (Invariance of Domain). *If M is a connected topological manifold, then its dimension is well-defined: it is the unique integer $n = \dim M$ such that $H_n(M, M \setminus \{x\}) \neq 0$ for all $x \in M$, and $H_k(M, M \setminus \{x\}) = 0$ for all $k \neq n$. Furthermore, if M is homeomorphic to N and both are connected, then $\dim M = \dim N$.*

Proof. Step 1. Using the excision axiom, for all k and any open $U \subseteq M$, $H_k(M, M \setminus \{x\}) \simeq H_k(U, U \setminus \{x\})$.

Step 2. Let $x \in M$ be arbitrary, and choose an open neighborhood of x $U \subseteq M$ that is homeomorphic to some $V \subseteq \mathbb{R}^{n(x)}$, where $n(x)$ may be dependent on x . Now using excision and writing φ to be the homeomorphism, $H_k(V, V \setminus \{\varphi(x)\}) \simeq H_k(B, B \setminus \{\varphi(x)\})$ for some open ball contained in V also containing $\varphi(x)$. But an open ball in $\mathbb{R}^{n(x)}$ is homeomorphic to $\mathbb{R}^{n(x)}$ itself. To conclude: $H_k(U, U \setminus \{x\}) \simeq H_k(\mathbb{R}^{n(x)}, \mathbb{R}^{n(x)} \setminus \{\varphi(x)\})$ where φ is the final homeomorphism, after abusing notation. But by Theorem 20.4, we conclude that

$$H_k(M, M \setminus \{x\}) \simeq H_k(\mathbb{R}^{n(x)}, \mathbb{R}^{n(x)} \setminus \{\varphi(x)\}) \simeq \begin{cases} 0, & k \neq n(x) \\ H_0(x_0), & k = n(x). \end{cases}$$

Step 3. Step 2 gives us a function $n : M \rightarrow \mathbb{N}$. Our aim is to now show that n is constant;. Indeed, if $y \in U$, then $n(y) = n(x)$, since

$$H_k(U, U \setminus \{y\}) \simeq H_k(\mathbb{R}^{n(x)}, \mathbb{R}^{n(x)} \setminus \{\varphi(y)\}) \simeq H_k(\mathbb{R}^{n(y)}, \mathbb{R}^{n(y)} \setminus \{\varphi(x)\}),$$

and hence n is a locally constant function on M . Now with the observation that M is connected, we conclude that $n(x) = n(y)$ for all $x, y \in M$.

Step 4. To prove the last statement, if $\psi : M \rightarrow N$ is a homeomorphism, then we have the isomorphisms $H_k(M, M \setminus \{x\}) \simeq H_k(N, N \setminus \{\psi(x)\})$ for all k . ■

We can generalize the definition of manifold a little further via the previous theorem.

Definition 21.3. A space X is called a **homological manifold** at x if there exists some n_x such that $H_k(X, X \setminus \{x\}) = 0$ for all $k \neq n_x$, $H_{n_x}(X, X \setminus \{x\}) \simeq H_0(x_0)$. We call n_x the **pointwise homological dimension**, and we write $\dim xX = n_x$.

21.2 Orientability

For this section, we will assume that $H_0(x_0) \simeq \mathbb{Z}$. Notice that if so, there are two choices of isomorphisms: either we can send the generator of $H_0(x_0)$ to 1 in \mathbb{Z} , or we can send it to -1 in \mathbb{Z} .

Remark 21.4. While two distinct choices for an isomorphisms exist for \mathbb{Z} , no two distinct choices for an isomorphism can exist for the case of \mathbb{Z}_2 : that is, if $H_0(x_0) \simeq \mathbb{Z}_2$, then there cannot be more than one isomorphism.

Now suppose that M is a connected manifold, with $\dim M = n$. Then as we saw in Theorem 21.2, $H_n(X, X \setminus \{x\}) \simeq H_0(x_0) \simeq \mathbb{Z}$. Then in order to define a notion of orientability, we want to be able to pick an element $1 \in H_n(X, X \setminus \{x\})$ to choose an isomorphism.

Exercise 21.5. Define $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ via

$$f(x_1, \dots, x_n) = (-x_1, x_2, \dots, x_n).$$

Then consider the induced maps on homology $f_n : H_n(\mathbb{R}^n, \mathbb{R}^n \setminus \{0\}) \rightarrow H_n(\mathbb{R}^n, \mathbb{R}^n \setminus \{0\})$. Show that $f_n = -\text{id}_{H_n(\mathbb{R}^n, \mathbb{R}^n \setminus \{0\})}$.

Definition 21.6. A manifold M is **orientable** if there exists a covering of M by open sets $\mathcal{U} = \{U_i\}_{i \in I}$ and a family of charts $\varphi_i : \mathbb{R}^n \rightarrow U_i$ homeomorphisms such that if $U_i \cap U_j \neq \emptyset$, then $\varphi_i^{-1} \circ \varphi_j : \varphi_j^{-1}(U_i \cap U_j) \rightarrow \varphi_i^{-1}(U_i \cap U_j)$ is **orientation-preserving**. That is, $\varphi_i^{-1} \circ \varphi_j$ induces a map $(\varphi_i^{-1} \circ \varphi_j)_n : H_n(\varphi_j^{-1}(U_i \cap U_j), \varphi_j^{-1}(U_i \cap U_j) \setminus \{a\}) \rightarrow H_n(\varphi_i^{-1}(U_i \cap U_j), \varphi_i^{-1}(U_i \cap U_j) \setminus \{\varphi_i^{-1}(\varphi_j(a))\})$; then the inclusion map $\iota : \varphi_j^{-1}(U_i \cap U_j) \hookrightarrow \mathbb{R}^n$ and similarly for $\varphi_i^{-1}(U_i \cap U_j)$ induces a map $\psi : H_n(\mathbb{R}^n, \mathbb{R}^n \setminus \{a\}) \rightarrow H_n(\mathbb{R}^n, \mathbb{R}^n \setminus \{\varphi_i^{-1}(\varphi_j(a))\})$. Now by translating $\varphi_i^{-1}(\varphi_j(a))$ to a in \mathbb{R}^n , we induce a map from $H_n(\mathbb{R}^n, \mathbb{R}^n \setminus \{a\})$ to itself. If this map is the identity, then we say that $\varphi_i^{-1} \circ \varphi_j$ is orientation preserving. To summarize, if the isomorphism on the bottom diagonal arrow in the following diagram is the identity map on $H_n(\mathbb{R}^n, \mathbb{R}^n \setminus \{a\})$, then we say $\varphi_i^{-1} \circ \varphi_j$ is orientation-preserving.


$$\begin{array}{ccc}
H_n(\varphi_j^{-1}(U_i \cap U_j), \varphi_j^{-1}(U_i \cap U_j) \setminus \{a\}) & \xrightarrow{(\varphi_i^{-1} \circ \varphi_j)^n} & H_n(\varphi_i^{-1}(U_i \cap U_j), \varphi_i^{-1}(U_i \cap U_j) \setminus \{\varphi_i^{-1}(\varphi_j(a))\}) \\
\downarrow \iota & & \downarrow \iota \\
H_n(\mathbb{R}^n, \mathbb{R}^n \setminus \{a\}) & \xrightarrow{\psi} & H_n(\mathbb{R}^n, \mathbb{R}^n \setminus \{\varphi_j^{-1}(\varphi_i(a))\}) \\
& \searrow \simeq & \downarrow \text{trans.} \\
& & H_n(\mathbb{R}^n, \mathbb{R}^n \setminus \{a\})
\end{array}$$

Remark 21.7. If M is orientable, then there is a coherent set of generators $1 \in H_n(M, M \setminus \{x\})$ for all x . That is, if x is contained in two different charts, then both charts induce the same generators in H_n .

Example 21.8. \mathbb{R}^n is orientable: \mathbb{R}^n itself is an atlas for \mathbb{R}^n .

Example 21.9. Any open, connected subset of \mathbb{R}^n is orientable.

Example 21.10. The Möbius strip is not orientable.



Exercise 21.11. Prove the above statement. [Hint: covering spaces.]

22 3/3/23

22.1 Axioms of Cohomology

Today we will discuss the axioms of cohomology and some of their consequences. Many of the results will mirror what we did in homology and the proofs are similar, so will be omitted.

For any pair of spaces (X, A) , we associate abelian groups $H^i(X, A)$ and boundary maps $\partial : H^i(A) \rightarrow H^{i+1}(X, A)$ and as before identify $H^i(X, \emptyset) = H^i(X)$, and for every map $f : (X, A) \rightarrow (Y, B)$ we have maps $f^i : H^i(Y, B) \rightarrow H^i(X, A)$, satisfying the following axioms:

(C1) if $f : (X, A) \rightarrow (X, A)$ is the identity map, then f^i is the identity map for each i .

(C2) If $(X, A) \xrightarrow{f} (Y, B) \xrightarrow{g} (Z, C)$ then the following diagram commutes:

$$\begin{array}{ccccc}
H^i(Z, C) & \xrightarrow{g^i} & H^i(Y, B) & \xrightarrow{f^i} & H^i(X, A) \\
& & \searrow (g \circ f)^i & \nearrow & \\
& & & &
\end{array}$$

That is, $(g \circ f)^i = f^i \circ g^i$ for each i .

(C3) The following diagram commutes:

$$\begin{array}{ccc} H^{i+1}(Y, B) & \xrightarrow{f^{i+1}} & H^{i+1}(X, A) \\ \partial \uparrow & & \partial \uparrow \\ H^i(B) & \xrightarrow{f^i} & H^i(A) \end{array}$$

(C4) We have the following long exact sequence:

$$\cdots \leftarrow H^i(A) \xleftarrow{J} H^i(X) \xleftarrow{\iota} H^i(X, A) \xleftarrow{\partial} H^{i-1}(A) \leftarrow \cdots,$$

where $\iota : (X, \emptyset) \hookrightarrow (X, A)$ and $J : A \hookrightarrow X$ are the inclusion maps.

(C5) If f is homotopic to g , then $f^i = g^i$ for all i .

(C6) If $U \subseteq A$ and $\bar{U} \subseteq \text{int } A$, then $\iota : (X \setminus U, A \setminus U) \hookrightarrow (X, A)$ induces isomorphisms of cohomology groups.

(C7) If $i \neq 0$, $H^i(x_0) = 0$.

Theorem 22.1 (Mayer-Vietoris). *Let $X = X_1 \cup X_2$ and $A = X_1 \cap X_2$, and assume that $\iota : (X, A) \hookrightarrow (X, X_2)$ is excisive, that is, $H^i(X, X_2) \rightarrow H^i(X, A)$ is an isomorphism. Then there is a long exact sequence*

$$\cdots \leftarrow H^i(A) \xleftarrow{\alpha} H^i(X_1) \oplus H^i(X_2) \xleftarrow{\beta} H^i(X) \xleftarrow{\Delta} H^{i-1}(A) \leftarrow \cdots.$$

Definition 22.2. If $f : X \rightarrow \{x_0\}$ a map into the one-point space, then this induces maps $f^i : H^i(x_0) \rightarrow H^i(X)$, and $\text{im } f^i \subseteq H^i(X)$. Then the **reduced cohomology groups** are the cokernels of these homomorphisms, $\tilde{H}^i(X) := \text{coker } f^i = H^i(X) / \text{im}(f^i)$.

Corollary 22.3. *The exact sequence in Theorem 22.1 holds for reduced cohomology groups.*

Theorem 22.4 (Suspension Theorem). *For any n , $\tilde{H}^n(\Sigma X) \simeq \tilde{H}^{n-1}(X)$.*

Theorem 22.5. *$\tilde{H}^n(S^n) \simeq H^0(x_0)$ and $\tilde{H}^k(S^n) = 0$ for $k \neq n$.*

22.2 Eilenberg-MacLane Cohomology (Optional)

Remember the definition of Eilenberg-MacLane spaces (Definition 13.9). Here we will concern ourselves with a special case: path connected spaces B_n such that $\pi_k(B_n, x_0) = 0$ for every $k \neq n$ and $\pi_n(B_n, 0) \simeq \mathbb{Z}$. Thus as seen before, $B_1 = S^1$ and $B_2 = \mathbb{C}P^\infty = \bigcup_{n \geq 0} \mathbb{C}P^n$.

Define a sort of cohomology using these spaces as follows: let $H^n(X) := [X, B_n]$, the set of homotopy classes of maps $X \rightarrow B_n$. Now if $f : X \rightarrow Y$, this induces maps $f^n : H^n(Y) \rightarrow H^n(X)$ via $f^n(v) = v \circ f$, i.e., if $v : Y \rightarrow B_n$, then $v \circ f : X \rightarrow B_n$. These maps are well-defined because we can compose homotopy. Moreover, it maps the identity to the identity, where the identity is the constant map. Finally, $H^0(x_0) = \mathbb{Z}$ by definition.

Now we ask ourselves, what is $H^1(X)$? If $v \in H^1(X)$, then we have $v : X \rightarrow S^1$. Take $x_0 \in X$. Then v induces a map $v_\# : \pi_1(X, x_0) \rightarrow \pi_1(S^1, v(x_0)) \simeq \mathbb{Z}$ in the obvious manner: if $\alpha : [0, 1] \rightarrow X$ is a loop based at x_0 , then $(v_\#(\alpha))(t) = v(\alpha(t))$. It can be checked that this map is well-defined, that is, it only depends on the homotopy class. Then we get a map

$$\begin{aligned} L : H^1(X) &= [X, S^1] \rightarrow \text{Hom}(\pi_1(X, x_0) \rightarrow \mathbb{Z}) \\ v &\mapsto v_\#. \end{aligned}$$

Now if we assume that X is path-connected and has a universal covering, then it can be shown that L is an isomorphism!