

1 buildH: Build hydrogen atoms from united-atom  
2 molecular dynamics of lipids and calculate the order  
3 parameters

4 **Hubert Santuz<sup>1, 2</sup>, Amélie Bacle<sup>3</sup>, Pierre Poulain<sup>4</sup>, and Patrick F.J.  
5 Fuchs<sup>\*5, 6</sup>**

6 **1** CNRS, Université de Paris, UPR 9080, Laboratoire de Biochimie Théorique, 13 Rue Pierre et  
7 Marie Curie, F-75005 Paris, France **2** Institut de Biologie Physico-Chimique-Fondation Edmond de  
8 Rothschild, PSL Research University, Paris, France **3** Laboratoire Coopératif "Lipotoxicity and  
9 Channelopathies - ConicMeds," Université de Poitiers, F-86000 Poitiers, France **4** Université de  
10 Paris, CNRS, Institut Jacques Monod, F-75006, Paris, France **5** Sorbonne Université, Ecole Normale  
11 Supérieure, PSL Research University, CNRS, Laboratoire des Biomolécules (LBM), F-75005 Paris,  
12 France **6** Université de Paris, UFR Sciences du Vivant, F-75013 Paris, France

DOI: [10.21105/joss.03521](https://doi.org/10.21105/joss.03521)

#### Software

- [Review](#) ↗
- [Repository](#) ↗
- [Archive](#) ↗

**Editor:** Richard Gowers ↗

#### Reviewers:

- [@lilyminium](#)
- [@ojeda-e](#)
- [@blakeaw](#)

**Submitted:** 08 July 2021

**Published:** 22 July 2021

#### License

Authors of papers retain  
copyright and release the work  
under a Creative Commons  
Attribution 4.0 International  
License ([CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

## 13 Background

14 Molecular dynamics (MD) simulations of lipids are widely used to understand the complex  
15 structure and dynamics of biological or model membranes ([Feller, 2000](#); [Lyubartsev & Rabi-  
16 novich, 2011](#); [Tieleman et al., 1997](#)). They are very complementary to biophysical experiments  
17 and have thus become important to get insights at the microscopic scale. Many of them are  
18 performed using all-atom (AA) or united-atom (UA) representations. In AA force fields (such  
19 as CHARMM36 ([Klauda et al., 2010](#))), all the atoms are considered whereas in UA force  
20 fields (such as Berger ([Berger et al., 1997](#))) the aliphatic hydrogen atoms (Hs) are merged  
21 to their parent carbon into a larger particle representing a CH, CH<sub>2</sub> or CH<sub>3</sub> (e.g. a methyl  
22 group is represented by a single CH<sub>3</sub> particle). In simulations of phospholipids, the use of UA  
23 representations allows to divide by almost 3 the number of atoms to simulate because these  
24 molecules contain many aliphatic Hs. It thus reduces the computational cost without losing  
25 important chemical details.

26 MD simulations of lipids are usually validated against experimental data ([Klauda et al., 2010](#))  
27 or used to help interpret experiments ([Feller, 2007](#)). One type of experiment which is often  
28 used for that is <sup>2</sup>H NMR. In this type of experiment, aliphatic Hs atoms are replaced by  
29 deuterons. <sup>2</sup>H NMR allows one to measure the order parameter of a given C-H bond (where  
30 the H is replaced by a deuteron):

$$S_{CH} = \frac{1}{2} \langle 3\cos^2(\theta) - 1 \rangle$$

31 where  $\theta$  is the angle between the C-H bond and the magnetic field. The symbol  $\langle \dots \rangle$  means  
32 averaging over time and molecules.

33 This order parameter is useful because it is directly related to the flexibility of the given C-H  
34 bond. It describes the amount of possible orientations visited by the C-H bond, varies between  
35 1 and -0.5 and is unitless. Values close to 0 mean high mobility, when it goes away from 0  
36 (towards negative or positive values), it means the C-H bond gets less mobile. Although it  
37 varies theoretically between 1 and -0.5, its absolute value  $|S_{CH}|$  is often reported because

\*corresponding author

its sign is usually difficult to measure experimentally (Ollila, 2014). In MD simulations,  $\theta$  is the angle between the C-H bond and the normal to the membrane (usually the  $z$  axis). For AA simulations,  $S_{CH}$  is trivial to calculate. However, it is more difficult for UA simulations since the aliphatic Hs are not present. There are two main strategies to compute  $S_{CH}$  from UA simulations (Piggot et al., 2017): i) expressing  $S_{CH}$  as a function of the coordinates of other atoms (Douliez et al., 1995), ii) reconstructing Hs coordinates and calculate  $S_{CH}$  as in AA simulations. The trend in the last years was more to use strategy ii), such as in the NMRLipids project (Antila et al., 2019; Bacle et al., 2021; Botan et al., 2015; Catte et al., 2016). NMRLipids is an open science project which uses MD simulations and experimental  $S_{CH}$  with the goal of improving lipid force fields or making fundamental research on the structure and dynamics of lipid membranes.

## Statement of Need

Reconstructing Hs from the heavy atom coordinates can be done using standard geometric rules respecting stereochemistry. In the first NMRLipids project (Botan et al., 2015), H reconstruction was performed using a program called `g_protonate` from the software GROMACS version 3.\* (Berendsen et al., 1995). However, `g_protonate` has been removed from GROMACS version 4.\* or higher. So, there was a need to find other solutions. Currently, there are many programs in the field of chemoinformatics that are able to build Hs such as OpenBabel (O'Boyle et al., 2011) or other proprietary softwares. It is also possible to use `pdb2gmx` from the GROMACS software (Abraham et al., 2015) to build Hs, but it is not initially intended for that since its main purpose is to build a topology. Many of these solutions remain workarounds where one uses a sophisticated software for just doing one basic task. Using these open or proprietary software usually has a slow learning curve. Moreover, it is not always easy to use them in the Unix command line when they have complicated graphical interfaces, thus complicating their use in automatic pipelines of trajectory analyses. Here, we propose the `buildH` software to meet this need. `buildH` is very light and usable in the Unix command line or as a Python module, making it a tool of choice to integrate in analyses pipelines or Jupyter notebooks. It can be easily extended or customized if needed. `buildH` is currently widely used in the NMRLipids project IVb dealing with the conformational plasticity of lipid headgroups in cellular membranes and protein-lipid complexes (Bacle et al., 2021). In addition, it is planned to use `buildH` in the next NMRLipids project dealing with a databank containing MD trajectories of lipids (Kiirikki & Ollila, 2021).

## Overview

`buildH` is a Python software that enables automatic analyses of order parameter calculations from UA trajectories of lipids. The software has the following features:

- It reads a single structure or a trajectory of lipids in a UA representation.
- It reconstructs the aliphatic Hs using standard geometric rules.
- From the reconstructed Hs, it calculates and outputs the order parameters on each requested C-H bond.
- Optionally, it outputs a structure (in `pdb` format) and a trajectory (in `xtc` format) with all reconstructed Hs.

Beyond order parameter calculations, the trajectory with Hs can be used for any further analyses (e.g. precise molecular volume calculation).

81 buildH has been natively developed for a use in the Unix command line. It possesses a  
82 minimum number of options, making it easy to use. It is also possible to utilize it as a Python  
83 module, which may be convenient in some cases.

84 To reconstruct H atoms, buildH uses standard geometric rules. These rules require so-  
85 called *helper* atoms. For example, the reconstruction of the two Hs of a CH<sub>2</sub> on carbon  $C_i$ ,  
86 requires two helpers which are  $C_{i-1}$  and  $C_{i+1}$ , that is, the two neighbors of  $C_i$  in the chain  
87 (note that helpers can also be other heavy atoms such as oxygen or nitrogen). The list of  
88 helpers used for the reconstruction of each H is written in a json file. Many json files are  
89 already present on the buildH repository representing the major lipids: Phosphatidylcholine  
90 (PC), Phosphatidylethanolamine (PE), Phosphatidylglycerol (PG) for the polar heads and  
91 palmitoyl, myristoyl, oleoyl for the aliphatic chains, as well as cholesterol. Major UA force  
92 fields are also represented (Berger (Berger et al., 1997), GROMOS-CPK (Piggot et al., 2012),  
93 CHARMM-UA (Lee et al., 2014)). In case a user wants to analyze a lipid which is not  
94 present in buildH, a step-by-step documentation guides the user in the process of creating  
95 and supplying his/her lipid description as a json file.

96 All structure and trajectory read / write operations are handled by the MDAnalysis module  
97 (Gowers et al., 2016; Michaud-Agrawal et al., 2011). Mathematical vector operations are  
98 performed by Numpy (Harris et al., 2020) and accelerated with Numba (Lam et al., 2015),  
99 leading to very decent performances. For example, the reconstruction of all Hs and order  
100 parameter calculation on a trajectory of 2500 frames with 128 POPC molecules can be handled  
101 in approximately 7 minutes using a single core Xeon processor at 3.60 GHz, whereas it was  
102 almost 30 minutes long without Numba.

103 buildH has been implemented with good practices of software development in mind (Jiménez  
104 et al., 2017; Taschuk & Wilson, 2017):

- 105     ▪ version controlled repository on GitHub <https://github.com/patrickfuchs/buildH>,
- 106     ▪ open-source license (BSD-3-Clause),
- 107     ▪ continuous integration through tests,
- 108     ▪ and documentation <https://buildh.readthedocs.io/>.

109 Some notebooks are provided in the GitHub repository to explain how buildH works and how  
110 to analyze the data produced. In case of trouble, any user can post an issue on GitHub.

111 buildH is available in the Python Package Index (PyPI) as well as in the Bio-  
112 conda repository. The current version 1.5.0 of buildH is archived in the Zenodo  
113 repository (<https://zenodo.org/record/5080126>) and in the Software Heritage archive  
114 ([sw.h1:dir:eb46a03f6e6188f93bd0b39ab78b4640e777ecd1](https://sw.h1:dir:eb46a03f6e6188f93bd0b39ab78b4640e777ecd1)).

## 115 Acknowledgements

116 The authors thank the community of [NMRLipids](#) for useful discussions, especially Samuli Ollila.

## 117 References

- 118 Abraham, M. J., Murtola, T., Schulz, R., Páll, S., Smith, J. C., Hess, B., & Lindahl, E. (2015).  
119 GROMACS: High performance molecular simulations through multi-level parallelism from  
120 laptops to supercomputers. *SoftwareX*, 1-2, 19–25. [https://doi.org/10.1016/j.softx.2015.](https://doi.org/10.1016/j.softx.2015.06.001)  
121 [06.001](https://doi.org/10.1016/j.softx.2015.06.001)
- 122 Antila, H., Buslaev, P., Favela-Rosales, F., Ferreira, T. M., Gushchin, I., Javanainen, M., Kav,  
123 B., Madsen, J. J., Melcr, J., Miettinen, M. S., Määttä, J., Nencini, R., Ollila, O. H. S., &

- Piggot, T. J. (2019). Headgroup structure and cation binding in phosphatidylserine lipid bilayers. *The Journal of Physical Chemistry B*, 123(43), 9066–9079. <https://doi.org/10.1021/acs.jpcc.9b06091>
- Bacle, A., Buslaev, P., Fandiño, R. G., Favela-Rosales, F., Ferreira, T. M., Fuchs, P. F. J., Gushchin, I., Javanainen, M., Kiirikki, A. M., Madsen, J. J., Melcr, J., Rodríguez, P. M., Miettinen, M. S., Ollila, O. H. S., Papadopoulos, C. G., Peón, A., Piggot, T. J., Piñero, Ángel, & Virtanen, S. I. (2021). <https://doi.org/10.5281/zenodo.4748646>
- Berendsen, H. J. C., van der Spoel, D., & van Drunen, R. (1995). GROMACS: A message-passing parallel molecular dynamics implementation. *Computer Physics Communications*, 91(1), 43–56. [https://doi.org/10.1016/0010-4655\(95\)00042-E](https://doi.org/10.1016/0010-4655(95)00042-E)
- Berger, O., Edholm, O., & Jähnig, F. (1997). Molecular dynamics simulations of a fluid bilayer of dipalmitoylphosphatidylcholine at full hydration, constant pressure, and constant temperature. *Biophysical Journal*, 72(5), 2002–2013. [https://doi.org/10.1016/S0006-3495\(97\)78845-3](https://doi.org/10.1016/S0006-3495(97)78845-3)
- Botan, A., Favela-Rosales, F., Fuchs, P. F. J., Javanainen, M., Kanduč, M., Kulig, W., Lamberg, A., Loison, C., Lyubartsev, A., Miettinen, M. S., Monticelli, L., Määttä, J., Ollila, O. H. S., Retegan, M., Róg, T., Santuz, H., & Tynkkynen, J. (2015). Toward atomistic resolution structure of phosphatidylcholine headgroup and glycerol backbone at different ambient conditions. *The Journal of Physical Chemistry B*, 119(49), 15075–15088. <https://doi.org/10.1021/acs.jpcc.5b04878>
- Catte, A., Grych, M., Javanainen, M., Loison, C., Melcr, J., Miettinen, M. S., Monticelli, L., Määttä, J., Oganessian, V. S., Ollila, O. H. S., Tynkkynen, J., & Vilov, S. (2016). Molecular electrometer and binding of cations to phospholipid bilayers. *Phys. Chem. Chem. Phys.*, 18, 32560–32569. <https://doi.org/10.1039/C6CP04883H>
- Douliez, J. P., Léonard, A., & Dufourc, E. J. (1995). Restatement of order parameters in biomembranes: Calculation of c-c bond order parameters from c-d quadrupolar splittings. *Biophysical Journal*, 68(5), 1727–1739. [https://doi.org/10.1016/S0006-3495\(95\)80350-4](https://doi.org/10.1016/S0006-3495(95)80350-4)
- Feller, S. E. (2000). Molecular dynamics simulations of lipid bilayers. *Current Opinion in Colloid & Interface Science*, 5(3), 217–223. [https://doi.org/10.1016/S1359-0294\(00\)00058-3](https://doi.org/10.1016/S1359-0294(00)00058-3)
- Feller, S. E. (2007). Molecular dynamics simulations as a complement to nuclear magnetic resonance and x-ray diffraction measurements. In A. M. Dopico (Ed.), *Methods in membrane lipids* (pp. 89–102). Humana Press. [https://doi.org/10.1007/978-1-59745-519-0\\_7](https://doi.org/10.1007/978-1-59745-519-0_7)
- Gowers, Richard J., Linke, Max, Barnoud, Jonathan, Reddy, Tyler J. E., Melo, Manuel N., Seyler, Sean L., Domański, Jan, Dotson, David L., Buchoux, Sébastien, Kenney, Ian M., & Beckstein, Oliver. (2016). MDAnalysis: A Python Package for the Rapid Analysis of Molecular Dynamics Simulations. In Sebastian Benthall & Scott Rostrup (Eds.), *Proceedings of the 15th Python in Science Conference* (pp. 98–105). <https://doi.org/10.25080/Majors-629e541a-00e>
- Harris, C. R., Millman, K. J., Walt, S. J. van der, Gommers, R., Virtanen, P., Cournapeau, D., Wieser, E., Taylor, J., Berg, S., Smith, N. J., Kern, R., Picus, M., Hoyer, S., Kerkwijk, M. H. van, Brett, M., Haldane, A., Río, J. F. del, Wiebe, M., Peterson, P., ... Oliphant, T. E. (2020). Array programming with NumPy. *Nature*, 585(7825), 357–362. <https://doi.org/10.1038/s41586-020-2649-2>
- Jiménez, R. C., Kuzak, M., Alhamdoosh, M., Barker, M., Batut, B., Borg, M., Capella-Gutierrez, S., Chue Hong, N., Cook, M., Corpas, M., Flannery, M., Garcia, L., Gelpí, J. Ll., Gladman, S., Goble, C., González Ferreira, M., Gonzalez-Beltran, A., Griffin, P. C., Grüning, B., ... Crouch, S. (2017). Four simple recommendations to encourage best practices in research software. *F1000Research*, 6, 876. <https://doi.org/10.12688/f1000research.11407.1>

- 174 Kiirikki, A., & Ollila, S. (2021). *NMRLipids databank*. [https://github.com/NMRLipids/](https://github.com/NMRLipids/Databank)  
175 [Databank](https://github.com/NMRLipids/Databank)
- 176 Klauda, J. B., Venable, R. M., Freites, J. A., O'Connor, J. W., Tobias, D. J., Mondragon-  
177 Ramirez, C., Vorobyov, I., MacKerell, A. D., & Pastor, R. W. (2010). Update of the  
178 CHARMM all-atom additive force field for lipids: Validation on six lipid types. *The Journal*  
179 *of Physical Chemistry B*, 114(23), 7830–7843. <https://doi.org/10.1021/jp101759q>
- 180 Lam, S. K., Pitrou, A., & Seibert, S. (2015). Numba: A LLVM-based python JIT compiler.  
181 *LLVM '15: Proceedings of the Second Workshop on the LLVM Compiler Infrastructure in*  
182 *HPC*, 1–6. <https://doi.org/10.1145/2833157.2833162>
- 183 Lee, S., Tran, A., Allsopp, M., Lim, J. B., Hénin, J., & Klauda, J. B. (2014). CHARMM36  
184 united atom chain model for lipids and surfactants. *The Journal of Physical Chemistry B*,  
185 118(2), 547–556. <https://doi.org/10.1021/jp410344g>
- 186 Lyubartsev, A. P., & Rabinovich, A. L. (2011). Recent development in computer simulations  
187 of lipid bilayers. *Soft Matter*, 7, 25–39. <https://doi.org/10.1039/C0SM00457J>
- 188 Michaud-Agrawal, N., Denning, E. J., Woolf, T. B., & Beckstein, O. (2011). MDAAnalysis:  
189 A toolkit for the analysis of molecular dynamics simulations. *Journal of Computational*  
190 *Chemistry*, 32(10), 2319–2327. <https://doi.org/10.1002/jcc.21787>
- 191 O'Boyle, N. M., Banck, M., James, C. A., Morley, C., Vandermeersch, T., & Hutchison, G.  
192 R. (2011). Open babel: An open chemical toolbox. *Journal of Cheminformatics*, 3(33).  
193 <https://doi.org/10.1186/1758-2946-3-33>
- 194 Ollila, S. (2014). *On the signs of the order parameters*. [http://nmrlipids.blogspot.com/2014/](http://nmrlipids.blogspot.com/2014/04/on-signs-of-order-parameters.html)  
195 [04/on-signs-of-order-parameters.html](http://nmrlipids.blogspot.com/2014/04/on-signs-of-order-parameters.html)
- 196 Piggot, T. J., Allison, J. R., Sessions, R. B., & Essex, J. W. (2017). On the calculation  
197 of acyl chain order parameters from lipid simulations. *Journal of Chemical Theory and*  
198 *Computation*, 13(11), 5683–5696. <https://doi.org/10.1021/acs.jctc.7b00643>
- 199 Piggot, T. J., Piñeiro, Á., & Khalid, S. (2012). Molecular dynamics simulations of phos-  
200 phatidylcholine membranes: A comparative force field study. *Journal of Chemical Theory*  
201 *and Computation*, 8(11), 4593–4609. <https://doi.org/10.1021/ct3003157>
- 202 Taschuk, M., & Wilson, G. (2017). Ten simple rules for making research software more robust.  
203 *PLOS Computational Biology*, 13(4), e1005412. [https://doi.org/10.1371/journal.pcbi.](https://doi.org/10.1371/journal.pcbi.1005412)  
204 [1005412](https://doi.org/10.1371/journal.pcbi.1005412)
- 205 Tieleman, D. P., Marrink, S. J., & Berendsen, H. J. C. (1997). A computer perspective of  
206 membranes: Molecular dynamics studies of lipid bilayer systems. *Biochimica Et Biophysica*  
207 *Acta (BBA) - Reviews on Biomembranes*, 1331(3), 235–270. [https://doi.org/10.1016/](https://doi.org/10.1016/S0304-4157(97)00008-7)  
208 [S0304-4157\(97\)00008-7](https://doi.org/10.1016/S0304-4157(97)00008-7)