

LDDS: Python package for computing and visualizing Lagrangian Descriptors for Dynamical Systems

Broncio Aguilar-Sanjuan¹, Víctor J. García-Garrido², Vladimír
Krajňák¹, Shibabrat Naik¹, and Stephen Wiggins¹

¹ School of Mathematics, University of Bristol, Fry Building, Woodland Road, Bristol BS8 1UG,
United Kingdom ² Departamento de Física y Matemáticas, Universidad de Alcalá, Madrid, 28871,
Spain

DOI: [10.21105/joss.03482](https://doi.org/10.21105/joss.03482)

Software

- [Review](#) ↗
- [Repository](#) ↗
- [Archive](#) ↗

Editor: Richard Gowers ↗

Reviewers:

- [@nicoguardo](#)
- [@jmbr](#)

Submitted: 19 May 2021

Published: 11 July 2021

License

Authors of papers retain
copyright and release the work
under a Creative Commons
Attribution 4.0 International
License ([CC BY 4.0](#)).

Statement of Need

Nonlinear dynamical systems are ubiquitous in natural and engineering sciences, such as fluid mechanics, theoretical chemistry, ship dynamics, rigid body dynamics, atomic physics, solid mechanics, condensed matter physics, mathematical biology, oceanography, meteorology and celestial mechanics ([Wiggins, 1994](#) and references therein). There have been many advances in understanding phenomena across these disciplines using the geometric viewpoint of the solutions and the underlying structures in the phase space; for example ([MacKay et al., 1984](#)), ([V. Rom-Kedar et al., 1990](#)), ([Ozorio de Almeida et al., 1990](#)), ([V. Rom-Kedar & Wiggins, 1990](#)), ([J. D. Meiss, 1992](#)), ([Koon et al., 2000](#)), ([Waalkens et al., 2005](#)), ([J. D. Meiss, 2015](#)), ([Wiggins, 2016](#)), ([Zhong et al., 2018](#)), ([Zhong & Ross, 2020](#)). Chief among these phase space structures are the invariant manifolds that form a barrier between dynamically distinct solutions. In most nonlinear systems, the invariant manifolds are computed using numerical techniques that rely on some form of linearization around equilibrium points followed by continuation and globalization. However, these methods become computationally expensive and challenging when applied to the high-dimensional phase space of vector fields defined analytically, from numerical simulations or experimental data. This points to the need for techniques that can be paired with trajectory calculations, without the excessive computational overhead and at the same time can allow visualization along with trajectory data. The Python package, LDDS, serves this need for analyzing deterministic and stochastic, continuous and discrete high-dimensional nonlinear dynamical systems described either by an analytical vector field or from data obtained from numerical simulations or experiments.

To the best of our knowledge, no other software for calculating Lagrangian descriptors exists. A variety of computational tools is available for competing approaches popular in fluid mechanics, such as the identification of Lagrangian coherent structures via finite-time Lyapunov exponents ([Briot & d'Ovidio, 2011](#)), ([Nelson & Jacobs, 2016](#)), ([Onu et al., 2015](#)), ([Finn & Apte, 2013](#)), ([Dabiri Lab, 2009](#)), ([Haller et al., 2020](#)) and finite-size Lyapunov exponents ([Briot & d'Ovidio, 2011](#)) or Eulerian coherent structures ([Katsanoulis & Haller, 2018](#)).

Summary and Functionalities

The LDDS software is a Python-based module that provides the user with the capability of analyzing the phase space structures of both continuous and discrete nonlinear dynamical systems in the deterministic and stochastic settings through the method of Lagrangian descriptors (LDs). The main idea behind this methodology is to define a Lagrangian descriptor as the integral of a non-negative function g that captures a dynamical property of the evolution of a trajectory. Different formulations of the Lagrangian descriptor exist in the literature where the non-negative function g is: the arclength of a trajectory in phase space ([Jiménez](#)

43 Madrid & Mancho, 2009), (Mancho et al., 2013), the arclength of a trajectory projected on
 44 the configuration space (Craven & Hernandez, 2015), the p -norm or p -quasinorm (Lopesino
 45 et al., 2017), and the Maupertuis' action of Hamiltonian mechanics (Montoya & Wiggins,
 46 2020). The approach provided by Lagrangian descriptors for revealing phase space structure
 47 has also been adapted to address discrete-time systems (maps) and stochastic systems.

48 Consider a continuous-time dynamical system:

$$\frac{d\mathbf{x}}{dt} = \mathbf{f}(\mathbf{x}(t), t) \quad (1)$$

49 where $\mathbf{x} \in \mathbb{R}^n$ and \mathbf{f} is the vector field. Starting from an initial condition $\mathbf{x}_0 = \mathbf{x}(t_0)$ at time
 50 $t = t_0$, $g(\mathbf{x}(t); \mathbf{x}_0)$ is integrated together with the trajectory forward and backward time over
 51 the interval $[t_0 - \tau, t_0 + \tau]$ to give the Lagrangian descriptor,

$$\mathcal{L}(\mathbf{x}_0, t_0, \tau) = \int_{t_0 - \tau}^{t_0 + \tau} g(\mathbf{x}(t); \mathbf{x}_0) dt. \quad (2)$$

52 Large differences in values obtained for initial conditions on a predefined grid indicate the
 53 presence of the phase space structures in the system and provide insight into their geometry.
 54 One of the main goals we pursue with this software is to make Lagrangian descriptors available
 55 to a large scientific audience and enable them to use this tool for reproducible research.

56 This open-source package incorporates the following features:

- 57 ■ Computation of LDs for two-dimensional maps.
- 58 ■ Study of the phase space structure of two-dimensional continuous dynamical systems
 59 with LDs.
- 60 ■ Computation of LDs for a system of two stochastic differential equations with additive
 61 noise.
- 62 ■ Computation of LDs on two-dimensional phase space planes for Hamiltonian systems
 63 with 2 or more degrees of freedom (DoF).
- 64 ■ Application of LDs to Hamiltonian systems with 2 DoF where the potential energy
 65 surface is known on a discrete spatial grid.
- 66 ■ Computation of LDs from a spatio-temporal discretization of a two-dimensional time-
 67 dependent vector field.
- 68 ■ Visual extraction of the invariant stable and unstable manifolds from the LD scalar field
 69 values.
- 70 ■ Addition to time-dependent external forcings for two-dimensional continuous dynamical
 71 systems.
- 72 ■ Different definitions for the Lagrangian descriptor function found in the literature.

73 All the different features of the module, and their usage across different settings, are illustrated
 74 through Jupyter-notebook tutorials. These tutorials would help users better understand how
 75 to set up a model dynamical system to which LDs is applied, and present them with different
 76 options for visualizing the results obtained from the analysis. We believe that these resources
 77 provide useful material for the development of an effective learning process that could motivate
 78 the integration of this tool into users' research/academic projects. Moreover, this will surely
 79 encourage future contributions from the scientific community to extend the features and
 80 applicability of this software package to other areas.

81 Example systems

82 The following dynamical systems are included in this software package as examples to illustrate
 83 the application of Lagrangian descriptors:

84 Maps:

85 ▪ Standard map

86 The standard map is a two-dimensional map used in dynamical systems to study a number
87 of physical systems such as the cyclotron particle accelerator or a kicked rotor (Chirikov,
88 1971),(J. D. Meiss, 1992),(J. D. Meiss, 2008). The equations of the discrete system are given
89 by the expressions:

$$\begin{cases} x_{n+1} = x_n + y_n - \frac{K}{2\pi} \sin(2\pi x_n) \\ y_{n+1} = y_n - \frac{K}{2\pi} \sin(2\pi x_n) \end{cases} \quad (3)$$

90 where K is the parameter that controls the forcing strength of the perturbation. The inverse
91 map is described by:

$$\begin{cases} x_n = x_{n+1} - y_{n+1} \\ y_n = y_{n+1} + \frac{K}{2\pi} \sin(2\pi(x_{n+1} - y_{n+1})) \end{cases} \quad (4)$$

92 In the following figure, we show the output produced by the LDDS software package for the
93 standard map using the model parameter value $K = 1.2$.




Figure 1: Lagrangian descriptor contour plot for the standard map, using $p = 0.5$ -quasinorm and integration time $\tau = 15$.

94 Flows:

95 ▪ Forced undamped Duffing oscillator

96 The Duffing oscillator is an example of a periodically driven oscillator with nonlinear elasticity
97 (Duffing, 1918), (Kovacic & Brennan, 2011). This can model the oscillations of a pendulum
98 whose stiffness does not obey Hooke's law or the motion of a particle in a double-well potential.
99 It is also known as a simple system that can exhibit chaos.

100 As a special case, the forced undamped Duffing oscillator is described by a time-dependent
101 Hamiltonian given by:

$$H(x, p_x, t) = \frac{1}{2}p_x^2 - \frac{\alpha}{2}x^2 + \frac{\beta}{4}x^4 - f(t)x \quad (5)$$

102 where α and β are the model parameters and $f(t)$ is the time-dependent forcing added to the
103 system. The non-autonomous vector field that defines the dynamical system is given by:

$$\begin{cases} \dot{x} = \frac{\partial H}{\partial p_x} = f_1(x, p_x) = p_x \\ \dot{p}_x = -\frac{\partial H}{\partial x} = f_2(x, p_x, t) = \alpha x - \beta x^3 + f(t) \end{cases} \quad (6)$$

104 In the following figure we show the output produced by the LDDS software package for the
105 forced Duffing oscillator using the model parameter value $\alpha = \beta = 1$. The initial time is
106 $t_0 = 0$ and the perturbation used is of the form $f(t) = A \sin(\omega t)$ where $A = 0.25$ and $\omega = \pi$.




Figure 2: Lagrangian descriptor contour plot for the Duffing oscillator, using $p = 0.5$ -quasinorm and integration time $\tau = 15$.

107 ■ A double gyre flow with stochastic forcing

108 The double gyre is a recurrent pattern occurring in geophysical flows (Coulliette & Wiggins,
109 2001). The stochastic dynamical system for a simplified model of this flow (Shadden et
110 al., 2005) with additive noise is described by the following stochastic differential equations
111 (Balibrea-Iniesta et al., 2016):

$$\begin{cases} dX_t = \left(-\pi A \sin\left(\frac{\pi f(X_t, t)}{s}\right) \cos\left(\frac{\pi Y_t}{s}\right) - \mu X_t \right) dt + \sigma_1 dW_t^1 \\ dY_t = \left(\pi A \cos\left(\frac{\pi f(X_t, t)}{s}\right) \sin\left(\frac{\pi Y_t}{s}\right) \frac{\partial f}{\partial x}(X_t, t) - \mu Y_t \right) dt + \sigma_2 dW_t^2 \end{cases} \quad (7)$$

112 where W^1 and W^2 are Wiener processes and we have that:

$$f(X_t, t) = \varepsilon \sin(\omega t + \psi) X_t^2 + (1 - 2\varepsilon \sin(\omega t + \psi)) X_t \quad (8)$$

113 In the following figure we show the output produced by the LDDS software package for the
114 stochastically forced double gyre using a noise amplitude of $\sigma_1 = \sigma_2 = 0.1$. The double gyre
115 model parameters are $A = 0.25$, $\omega = 2\pi$, $\psi = \mu = 0$, $s = 1$, $\varepsilon = 0.25$, and the initial time is
116 $t_0 = 0$.




Figure 3: Lagrangian descriptor contour plot for the Double-gyre with stochastic forcing, using $p = 0.5$ -quasinorm and integration time $\tau = 15$.

117 Four-dimensional phase space:

- 118 ■ Hénon-Heiles Hamiltonian.

119 The Hénon-Heiles system is a simplified model describing the restricted motion of a star around
120 the center of a galaxy (Henon & Heiles, 1964). This system is a paradigmatic example of a
121 time-independent Hamiltonian with two degrees of freedom, given by the function:

$$H(x, y, p_x, p_y) = \frac{1}{2}(p_x^2 + p_y^2) + \frac{1}{2}(x^2 + y^2) + x^2y - \frac{1}{3}y^3 \quad (9)$$

122 where the vector field is:

$$\begin{aligned} \dot{x} &= \frac{\partial H}{\partial p_x} = p_x \\ \dot{y} &= \frac{\partial H}{\partial p_y} = p_y \\ \dot{p}_x &= -\frac{\partial H}{\partial x} = -x - 2xy \\ \dot{p}_y &= -\frac{\partial H}{\partial y} = -x^2 - y + y^2 \end{aligned} \quad (10)$$

123 In the next figure, we show the computation of Lagrangian descriptors with the LDDS software
124 package on the phase space slice described by the condition $x = 0$, $p_x > 0$ for the energy of
125 the system $H_0 = 1/5$.




Figure 4: Lagrangian descriptor contour plot for the Hénon-Heiles Hamiltonian, using $p = 0.5$ -quasinorm and integration time $\tau = 15$.

Relation to ongoing research projects

Lagrangian descriptors form the basis of several past and present research projects (Cámara et al., 2012), (Cámara et al., 2013), (Lopesino et al., 2015), (Craven & Hernandez, 2015), (Craven & Hernandez, 2016), (García-Garrido et al., 2016), (Balibrea-Iniesta et al., 2016), (Demian & Wiggins, 2017), (Craven et al., 2017), (Feldmaier et al., 2017), (Junginger et al., 2017), (García-Garrido et al., 2018), (Ramos et al., 2018), (Patra & Keshavamurthy, 2018), (Naik et al., 2019), (Naik & Wiggins, 2019), (Curbelo et al., 2019a), (Curbelo et al., 2019b), (Revuelta et al., 2019), (García-Garrido, Naik, et al., 2020), (García-Garrido, Agaoglou, et al., 2020), (Krajňák et al., 2020), (Naik & Wiggins, 2020), (Montoya & Wiggins, 2020), (Katsanikas et al., 2020). The common theme of all these projects is the investigation of phase space structures that govern phase space transport in nonlinear dynamical systems. We have also co-authored an open-source book project using Jupyter book (Executable Books Community, 2020) on the theory and applications of Lagrangian descriptors (Agaoglou et al., 2020). This open-source package is the computational companion to that book.

Acknowledgements

We acknowledge the support of EPSRC Grant No. EP/P021123/1 (Champs project) and Office of Naval Research (Grant No. N00014-01-1-0769).

References

- Agaoglou, M., Aguilar-Sanjuan, B., García-Garrido, V. J., González-Montoya, F., Katsanikas, M., Krajňák, V., Naik, S., & Wiggins, S. (2020). *Lagrangian descriptors: Discovery and quantification of phase space structure and transport*. zenodo: 10.5281/zenodo.3958985. <https://doi.org/10.5281/zenodo.3958985>
- Balibrea-Iniesta, F., Lopesino, C., Wiggins, S., & Mancho, A. M. (2016). Lagrangian descriptors for stochastic differential equations: A tool for revealing the phase portrait of stochastic dynamical systems. *International Journal of Bifurcation and Chaos*, 26(13), 1630036. <https://doi.org/10.1142/S0218127416300366>

- 152 Briol, F., & d'Ovidio, F. (2011). *Lagrangian*. [https://bitbucket.org/cnes_aviso/lagrangian/
153 src/master/](https://bitbucket.org/cnes_aviso/lagrangian/src/master/)
- 154 Cámara, A. de la, Mancho, A. M., Ide, K., Serrano, E., & Mechoso, C. R. (2012). Routes of
155 transport across the Antarctic polar vortex in the southern spring. *J. Atmos. Sci.*, 69(2),
156 753–767. <https://doi.org/10.1175/JAS-D-11-0142.1>
- 157 Cámara, A. de la, Mechoso, C. R., Mancho, A. M., Serrano, E., & Ide, K. (2013). Isen-
158 tropic transport within the Antarctic polar night vortex: Rossby wave breaking evidence
159 and Lagrangian structures. *J. Atmos. Sci.*, 70, 2982–3001. [https://doi.org/10.1175/
160 JAS-D-12-0274.1](https://doi.org/10.1175/JAS-D-12-0274.1)
- 161 Chirikov, B. V. (1971). Research concerning the theory of non-linear resonance and stochas-
162 ticity [translated at CERN by A. T. Sanders]. *CERN Trans.*, 71(40). [https://cds.cern.ch/
163 record/325497](https://cds.cern.ch/record/325497)
- 164 Coulliette, C., & Wiggins, S. (2001). Nonlinear Processes in Geophysics Intergyre transport in
165 a wind-driven, quasigeostrophic double gyre: An application of lobe dynamics. *Nonlinear
166 Processes in Geophysics*, 8, 69–94.
- 167 Craven, G. T., & Hernandez, R. (2015). Lagrangian descriptors of thermalized transition
168 states on time-varying energy surfaces. *Physical Review Letters*, 115(14), 148301. [https://
169 doi.org/10.1103/physrevlett.115.148301](https://doi.org/10.1103/physrevlett.115.148301)
- 170 Craven, G. T., & Hernandez, R. (2016). Deconstructing field-induced ketene isomerization
171 through lagrangian descriptors. *Physical Chemistry Chemical Physics*, 18(5), 4008–4018.
172 <https://doi.org/10.1039/C5CP06624G>
- 173 Craven, G. T., Junginger, A., & Hernandez, R. (2017). Lagrangian descriptors of driven
174 chemical reaction manifolds. *Physical Review E*, 96(2), 022222. [https://doi.org/10.
175 1103/PhysRevE.96.022222](https://doi.org/10.1103/PhysRevE.96.022222)
- 176 Curbelo, J., Mechoso, C. R., Mancho, A. M., & Wiggins, S. (2019a). Lagrangian study of
177 the final warming in the southern stratosphere during 2002: Part i. The vortex split-
178 ting at upper levels. *Climate Dynamics*, 53(5), 2779–2792. [https://doi.org/10.1007/
179 s00382-019-04832-y](https://doi.org/10.1007/s00382-019-04832-y)
- 180 Curbelo, J., Mechoso, C. R., Mancho, A. M., & Wiggins, S. (2019b). Lagrangian study of
181 the final warming in the southern stratosphere during 2002: Part II. 3D structure. *Climate
182 Dynamics*, 53(3), 1277–1286. <https://doi.org/10.1007/s00382-019-04833-x>
- 183 Dabiri Lab. (2009). *LCS Matlab kit* (Version 2.3). <http://dabirilab.com/software>.
- 184 Demian, A. S., & Wiggins, S. (2017). Detection of periodic orbits in Hamiltonian systems
185 using Lagrangian descriptors. *International Journal of Bifurcation and Chaos*, 27(14),
186 1750225. <https://doi.org/10.1142/S021812741750225X>
- 187 Duffing, G. (1918). *Erzwungene schwingungen bei veränderlicher eigenfrequenz und ihre
188 technische bedeutung*. Sammlung Vieweg, No 41/42. Vieweg & Sohn, Braunschweig.
- 189 Executable Books Community. (2020). *Jupyter book* (Version v0.10) [Computer software].
190 Zenodo. <https://doi.org/10.5281/zenodo.4539666>
- 191 Feldmaier, M., Junginger, A., Main, G., J. and Wunner, & Hernandez, R. (2017). Obtaining
192 time-dependent multi-dimensional dividing surfaces using lagrangian descriptors. *Chemical
193 Physics Letters*, 687, 194–199. <https://doi.org/10.1016/j.cplett.2017.09.008>
- 194 Finn, J., & Apte, S. V. (2013). Integrated computation of finite time Lyapunov exponent
195 fields during direct numerical simulation of unsteady flows. *Chaos*, 23, 013145. [https://
196 doi.org/10.1063/1.4795749](https://doi.org/10.1063/1.4795749)
- 197 García-Garrido, V. J., Agaoglou, M., & Wiggins, S. (2020). Exploring isomerization dy-
198 namics on a potential energy surface with an index-2 saddle using lagrangian descrip-

- tors. *Communications in Nonlinear Science and Numerical Simulation*, 89, 105331. <https://doi.org/10.1016/j.cnsns.2020.105331>
- García-Garrido, V. J., Curbelo, J., Mancho, A. M., Wiggins, S., & Mechoso, C. R. (2018). The application of lagrangian descriptors to 3D vector fields. *Regular and Chaotic Dynamics*, 23(5), 551–568. <https://doi.org/10.1134/S1560354718050052>
- García-Garrido, V. J., Naik, S., & Wiggins, S. (2020). Tilting and squeezing: Phase space geometry of Hamiltonian saddle-node bifurcation and its influence on chemical reaction dynamics. *International Journal of Bifurcation and Chaos*, 30(04), 2030008. <https://doi.org/10.1142/S0218127420300086>
- García-Garrido, V. J., Ramos, A., Mancho, A. M., Coca, J., & Wiggins, S. (2016). A dynamical systems perspective for a real-time response to a marine oil spill. *Marine Pollution Bulletin.*, 1–10. <https://doi.org/10.1016/j.marpolbul.2016.08.018>
- Haller, G., Katsanoulis, S., Holzner, M., Frohnapfel, B., & Gatti, D. (2020). Objective barriers to the transport of dynamically active vector fields. *J. Fluid Mech.*, 905, A17. <https://doi.org/10.1017/jfm.2020.737>
- Henon, M., & Heiles, C. (1964). The Applicability of the Third Integral Of Motion: Some Numerical Experiments. *THE ASTRONOMICAL JOURNAL*, 69(1). <https://doi.org/10.1086/109234>
- Jiménez Madrid, J. A., & Mancho, A. M. (2009). Distinguished trajectories in time dependent vector fields. *Chaos*, 19. <http://dx.doi.org/10.1063/1.3056050>
- Junginger, A., Duvenbeck, L., Feldmaier, M., Main, G., J. and Wunner, & Hernandez, R. (2017). Chemical dynamics between wells across a time-dependent barrier: Self-similarity in the lagrangian descriptor and reactive basins. *The Journal of Chemical Physics*, 147(6), 064101. <https://doi.org/10.1063/1.4997379>
- Katsanikas, M., García-Garrido, V. J., & Wiggins, S. (2020). The dynamical matching mechanism in phase space for caldera-type potential energy surfaces. *Chemical Physics Letters*, 743, 137199. <https://doi.org/10.1016/j.cplett.2020.137199>
- Katsanoulis, S., & Haller, G. (2018). *BarrierTool*. <https://github.com/haller-group/BarrierTool>
- Koon, W. S., Lo, M. W., Marsden, J. E., & Ross, S. D. (2000). Heteroclinic connections between periodic orbits and resonance transitions in celestial mechanics. *Chaos: An Interdisciplinary Journal of Nonlinear Science*, 10(2), 427–469. <https://doi.org/10.1063/1.166509>
- Kovacic, I., & Brennan, M. J. (2011). *The Duffing Equation: Nonlinear Oscillators and their Behaviour*. John Wiley; Sons. <https://doi.org/10.1002/9780470977859>
- Krajňák, V., Ezra, G. S., & Wiggins, S. (2020). Using lagrangian descriptors to uncover invariant structures in chesnavich's isokinetic model with application to roaming. *International Journal of Bifurcation and Chaos*, 30(5), 2050076. <https://doi.org/10.1142/S0218127420500765>
- Lopesino, C., Balibrea-Iniesta, F., García-Garrido, V. J., Wiggins, S., & Mancho, A. M. (2017). A theoretical framework for Lagrangian descriptors. *International Journal of Bifurcation and Chaos*, 27(01), 1730001. <https://doi.org/10.1142/S0218127417300014>
- Lopesino, C., Balibrea-Iniesta, F., Wiggins, S., & Mancho, A. M. (2015). Lagrangian descriptors for two dimensional, area preserving autonomous and nonautonomous maps. *Communications in Nonlinear Science and Numerical Simulation*, 27(1-3), 40–51. <https://doi.org/10.1016/j.cnsns.2015.02.022>
- MacKay, R. S., Meiss, J. D., & Percival, I. C. (1984). Transport in hamiltonian systems. *Physica D: Nonlinear Phenomena*, 13(1-2), 55–81. [https://doi.org/10.1016/0167-2789\(84\)90270-7](https://doi.org/10.1016/0167-2789(84)90270-7)

- Mancho, A. M., Wiggins, S., Curbelo, J., & Mendoza, C. (2013). Lagrangian Descriptors: A method for revealing phase space structures of general time dependent dynamical systems. *Communications in Nonlinear Science and Numerical Simulation*, 18(12), 3530–3557. <https://doi.org/10.1016/j.cnsns.2013.05.002>
- Meiss, J. D. (1992). Symplectic maps, variational principles, and transport. *Rev. Mod. Phys.*, 64, 795–848. <https://doi.org/10.1103/RevModPhys.64.795>
- Meiss, J. D. (2008). *Visual explorations of dynamics: The standard map* (No. 6; Vol. 70, pp. 965–988). <https://doi.org/10.1007/s12043-008-0103-3>
- Meiss, J. D. (2015). Thirty years of turnstiles and transport. *Chaos*, 25(9). <https://doi.org/10.1063/1.4915831>
- Montoya, F. G., & Wiggins, S. (2020). Revealing roaming on the double morse potential energy surface with lagrangian descriptors. *Journal of Physics A: Mathematical and Theoretical*, 53(23), 235702. <https://doi.org/10.1088/1751-8121/ab8b75>
- Montoya, F. G., & Wiggins, S. (2020). Revealing roaming on the double morse potential energy surface with lagrangian descriptors. *Journal of Physics A: Mathematical and Theoretical*, 53(23), 235702. <https://doi.org/10.1088/1751-8121/ab8b75>
- Naik, S., García-Garrido, V. J., & Wiggins, S. (2019). Finding NHIM: Identifying high dimensional phase space structures in reaction dynamics using Lagrangian descriptors. *Communications in Nonlinear Science and Numerical Simulation*, 79, 104907. <https://doi.org/10.1016/j.cnsns.2019.104907>
- Naik, S., & Wiggins, S. (2019). Finding normally hyperbolic invariant manifolds in two and three degrees of freedom with Hénon-Heiles type potential. *Phys. Rev. E*, 100, 022204. <https://doi.org/10.1103/PhysRevE.100.022204>
- Naik, S., & Wiggins, S. (2020). Detecting reactive islands in a system-bath model of isomerization. *Phys. Chem. Chem. Phys.* <https://doi.org/10.1039/D0CP01362E>
- Nelson, D. A., & Jacobs, G. B. (2016). High-order visualization of three-dimensional Lagrangian coherent structures with DG-FTLE. *Computers & Fluids*, 139, 197. <https://doi.org/10.1016/j.compfluid.2016.07.007>
- Onu, K., Huhn, F., & Haller, G. (2015). LCS Tool: A computational platform for Lagrangian coherent structures. *J. Comput. Sci.*, 7, 26. <https://doi.org/10.1016/j.jocs.2014.12.002>
- Ozorio de Almeida, A. M., De Leon, N., Mehta, M. A., & Marston, C. C. (1990). Geometry and dynamics of stable and unstable cylinders in hamiltonian systems. *Physica D: Nonlinear Phenomena*, 46(2), 265–285. [https://doi.org/10.1016/0167-2789\(90\)90040-V](https://doi.org/10.1016/0167-2789(90)90040-V)
- Patra, S., & Keshavamurthy, S. (2018). Detecting reactive islands using lagrangian descriptors and the relevance to transition path sampling. *Physical Chemistry Chemical Physics*, 20(7), 4970–4981. <https://doi.org/10.1039/C7CP05912D>
- Ramos, A. G., García-Garrido, V. J., Mancho, A. M., Wiggins, S., Coca, J., Glenn, S., Schofield, O., Kohut, J., Aragon, D., Kerfoot, J., Haskins, T., Miles, T., Haldeman, C., Strandskov, N., Allsup, B., Jones, C., & Shapiro, J. (2018). Lagrangian coherent structure assisted path planning for transoceanic autonomous underwater vehicle missions. *Scientific Reports*, 8, 4575. <https://doi.org/10.1038/s41598-018-23028-8>
- Revuelta, F., Benito, R. M., & Borondo, F. (2019). Unveiling the chaotic structure in phase space of molecular systems using lagrangian descriptors. *Physical Review E*, 99(3), 032221. <https://doi.org/10.1103/PhysRevE.99.032221>
- Rom-Kedar, V., Leonard, A., & Wiggins, S. (1990). An analytical study of transport, mixing and chaos in an unsteady vortical flow. *Journal of Fluid Mechanics*, 214, 347–394. <https://doi.org/10.1017/S0022112090000167>

- 294 Rom-Kedar, V., & Wiggins, S. (1990). Transport in two-dimensional maps. *Arch. Ration.*
295 *Mech. A.I.*, 109(3). <https://doi.org/10.1007/BF00375090>
- 296 Shadden, S. C., Lekien, F., & Marsden, J. E. (2005). Definition and properties of Lagrangian
297 coherent structures from finite-time Lyapunov exponents in two-dimensional aperiodic
298 flows. *Physica D: Nonlinear Phenomena*, 212(3-4), 271–304. [https://doi.org/10.1016/j.](https://doi.org/10.1016/j.physd.2005.10.007)
299 [physd.2005.10.007](https://doi.org/10.1016/j.physd.2005.10.007)
- 300 Waalkens, H., Burbanks, A., & Wiggins, S. (2005). Escape from planetary neighbourhoods.
301 *Monthly Notices of the Royal Astronomical Society*, 361(3), 763–775. [https://doi.org/](https://doi.org/10.1111/j.1365-2966.2005.09237.x)
302 [10.1111/j.1365-2966.2005.09237.x](https://doi.org/10.1111/j.1365-2966.2005.09237.x)
- 303 Wiggins, S. (2016). The role of normally hyperbolic invariant manifolds (NHIMs) in the
304 context of the phase space setting for chemical reaction dynamics. *Regular and Chaotic*
305 *Dynamics*, 21(6), 621–638. <https://doi.org/10.1134/S1560354716060034>
- 306 Wiggins, S. (1994). *Normally hyperbolic invariant manifolds in dynamical systems* (Vol. 105).
307 Springer Science & Business Media. <https://doi.org/10.1007/978-1-4612-4312-0>
- 308 Zhong, J., & Ross, S. D. (2020). Geometry of escape and transition dynamics in the presence
309 of dissipative and gyroscopic forces in two degree of freedom systems. *Communications*
310 *in Nonlinear Science and Numerical Simulation*, 82, 105033. [https://doi.org/10.1016/j.](https://doi.org/10.1016/j.cnsns.2019.105033)
311 [cnsns.2019.105033](https://doi.org/10.1016/j.cnsns.2019.105033)
- 312 Zhong, J., Virgin, L. N., & Ross, S. D. (2018). A tube dynamics perspective governing
313 stability transitions: An example based on snap-through buckling. *International Journal*
314 *of Mechanical Sciences*, 149, 413–428. <https://doi.org/10.1016/j.ijmecsci.2017.10.040>

DRAFT