

# DBSP\_DRP: A Python package for automated spectroscopic data reduction of DBSP data

Milan S. Roberson<sup>1,2</sup>, Christoffer Fremling<sup>2</sup>, and Mansi M. Kasliwal<sup>2</sup>

<sup>1</sup> Schmidt Academy of Software Engineering, California Institute of Technology <sup>2</sup> Division of Physics, Mathematics and Astronomy, California Institute of Technology <sup>3</sup> Caltech Optical Observatories, California Institute of Technology

DOI: [10.21105/joss.03511](https://doi.org/10.21105/joss.03511)

## Software

- [Review](#) ↗
- [Repository](#) ↗
- [Archive](#) ↗

Editor: [Pending Editor](#) ↗

Submitted: 19 July 2021

Published: 19 July 2021

## License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

## Summary

In astronomy, the spectrum of light emitted from astrophysical sources is of great use, allowing astronomers to classify objects and measure their properties. To measure the spectrum, astronomers use spectrographs, which use dispersive elements to split the incoming light into its constituent wavelengths, and then image this dispersed light with a detector, most commonly a CCD. But to do science with the spectrum, the 2D image in pixel coordinates taken by the CCD must be converted into a 1D spectrum of flux vs. wavelength. To increase the signal-to-noise ratio, astronomers can take multiple exposures of the same object and coadd their 1D spectra to reveal faint absorption lines or increase the precision with which an important emission line can be measured. Many spectrographs have multiple paths that light can go through, and multiple detectors, each measuring a particular part of the spectrum, to increase the wavelength range that can be captured in a single exposure, or to allow the high resolution observation of distinct wavelength ranges. If two detectors cover an overlapping region, caused by partial reflectance of a dichroic (wavelength-dependent beam splitter), then the spectra from each detector need to be spliced together, combining the light collected by each detector. This process of converting 2D CCD images into 1D spectra is called data reduction.

DBSP\_DRP is a python package that provides fully automated data reduction of data taken by the Double Spectrograph (DBSP) at the 200-inch Hale Telescope at Palomar Observatory (Oke & Gunn, 1982). The underlying data reduction functionality to extract 1D spectra, perform flux calibration and correction for atmospheric absorption, and coadd spectra together is provided by Pypelt (Prochaska et al., 2020). The new functionality that DBSP\_DRP brings is in orchestrating the complex data reduction process by making smart decisions so that no user input is required after verifying the correctness of the metadata in the raw FITS files in a table-like GUI. Though the primary function of DBSP\_DRP is to automatically reduce an entire night of data without user input, it has the flexibility for astronomers to fine-tune the data reduction with GUIs for manually identifying the faintest objects, as well as exposing the full set of Pypelt parameters to be tweaked for users with particular science needs. DBSP\_DRP also handles some of the occasional quirks specific to DBSP, such as swapping FITS header cards, adding (an) extra null byte/s to FITS files making them not conform to the FITS specification, and not writing the coordinates of the observation to file. Additionally, DBSP\_DRP contains a quicklook script for making real-time decisions during an observing run, and can open a GUI displaying a minimally reduced exposure in under 15 seconds. Docker containers are available for ease of deploying DBSP\_DRP in its quicklook configuration (without some large atmospheric model files) or in its full configuration.

## Statement of Need

Palomar Observatory, located near San Diego, CA, is a multinational observatory with a broad user base. Users come from large and small institutions, and their observing experience ranges from novice to expert. One responsibility for serving such a diverse user base is to provide software data reduction pipelines for the most frequently used instruments, such as the Palomar Double Spectrograph (DBSP). Although DBSP was commissioned in 1982, it remains the workhorse instrument of the 200" Hale Telescope. It is used on 42% of the nights in a year, comprising nearly all of the valuable "dark" (moonless) time. In previous years, standard astronomical practice left the data reduction up to the user. However, attitudes in instrument building have shifted since DBSP was built. The pipeline is now considered an indispensable component of the astronomical instrument. In fact, the difference between a good pipeline and a great pipeline means the difference between counting some of the photons vs. counting all of the photons.

Spectroscopy is a severe bottleneck in time-domain astronomy; currently less than 10% of discoveries are spectroscopically classified. Without a pipeline, data reduction is a difficult process and the standard method without a pipeline is to use IRAF, a 35 year old program on which development and maintenance was discontinued in 2013 and whose use is discouraged by many in the field e.g. [Ogaz & Tollerud \(2018\)](#). Needless to say, data reduction sans pipeline is extremely time-consuming. There is a clear need for a modern and stable automated data reduction pipeline for DBSP.

During observing runs, one would like to be able to quickly inspect data as it is taken, in order to ensure that it is of sufficient quality to do the desired science with. For objects whose brightness may have changed between a previous observation and the observing run, the observer may have uncertainties regarding how long of an exposure is needed to produce quality data. For very faint objects or objects in crowded fields, the observer may not even be sure that the telescope is pointed at the right object! A quicklook functionality, that can do a rudimentary reduction to correct for instrumental signatures and subtract light from the sky, revealing the spectra of the objects observed, can answer questions of exposure time and whether the object observed is the right one.

DBSP\_DRP is currently being used by the ZTF Bright Transient Survey ([Fremming et al., 2020](#); [Perley et al., 2020](#)), the ZTF Census of the Local Universe ([De et al., 2020](#)), and a program investigating ZTF Superluminous Supernovae ([Lunnan et al., 2020](#); [Chen et al., in preparation](#)). [Ravi et al. \(2021\)](#) is the first (known) publication that used DBSP\_DRP for data reduction. The development of DBSP\_DRP also lays the groundwork towards a fully automated pipeline for the Next Generation Palomar Spectrograph that is planned to be deployed on the Palomar 200-inch Hale Telescope in 2022.

## Acknowledgements

MSR acknowledges funding from the Schmidt Academy of Software Engineering, which is supported by the generosity of Eric and Wendy Schmidt by recommendation of the Schmidt Futures program.

We thank the following members of the time domain astronomy group at Caltech for beta-testing and providing valuable feedback during the development of this pipeline: Andy Tzanidakis, Lin Yan, Aishwarya Dahiwalé, Yuhang Yao, Yashvi Sharma, and Igor Andreoni.

MSR is extremely grateful to the welcoming, friendly, and helpful team of developers on the Pypelt team, without whom this package would not exist.

## References

- De, K., Kasliwal, M. M., Tzanidakis, A., Fremling, U. C., Adams, S., Aloisi, R., Andreoni, I., Bagdasaryan, A., Bellm, E. C., Bildsten, L., Cannella, C., Cook, D. O., Delacroix, A., Drake, A., Duev, D., Dugas, A., Frederick, S., Gal-Yam, A., Goldstein, D., ... Yao, Y. (2020). The Zwicky Transient Facility Census of the Local Universe. I. Systematic Search for Calcium-rich Gap Transients Reveals Three Related Spectroscopic Subclasses. *905*(1), 58. <https://doi.org/10.3847/1538-4357/abb45c>
- Fremling, C., Miller, A. A., Sharma, Y., Dugas, A., Perley, D. A., Taggart, K., Sollerman, J., Goobar, A., Graham, M. L., Neill, J. D., Nordin, J., Rigault, M., Walters, R., Andreoni, I., Bagdasaryan, A., Belicki, J., Cannella, C., Bellm, E. C., Cenko, S. B., ... Kulkarni, S. R. (2020). The Zwicky Transient Facility Bright Transient Survey. I. Spectroscopic Classification and the Redshift Completeness of Local Galaxy Catalogs. *The Astrophysical Journal*, *895*(1), 32. <https://doi.org/10.3847/1538-4357/ab8943>
- Lunnan, R., Yan, L., Perley, D. A., Schulze, S., Taggart, K., Gal-Yam, A., Fremling, C., Soumagnac, M. T., Ofek, E., Adams, S. M., Barbarino, C., Bellm, E. C., De, K., Fransson, C., Frederick, S., Golkhou, V. Z., Graham, M. J., Hallakoun, N., Ho, A. Y. Q., ... Yao, Y. (2020). Four (Super)luminous Supernovae from the First Months of the ZTF Survey. *The Astrophysical Journal*, *901*(1), 61. <https://doi.org/10.3847/1538-4357/abaec>
- Ogaz, S., & Tollerud, E. (2018). Removing the Institute's Dependence on IRAF (You can do it too!). *STScI Newsletter*, *35*(03).
- Oke, J. B., & Gunn, J. E. (1982). An Efficient Low Resolution and Moderate Resolution Spectrograph for the Hale Telescope. *Publications of the Astronomical Society of the Pacific*, *94*, 586. <https://doi.org/10.1086/131027>
- Perley, D. A., Fremling, C., Sollerman, J., Miller, A. A., Dahiwal, A. S., Sharma, Y., Bellm, E. C., Biswas, R., Brink, T. G., Bruch, R. J., De, K., Dekany, R., Drake, A. J., Duev, D. A., Filippenko, A. V., Gal-Yam, A., Goobar, A., Graham, M. J., Graham, M. L., ... Yan, L. (2020). The Zwicky Transient Facility Bright Transient Survey. II. A Public Statistical Sample for Exploring Supernova Demographics. *The Astrophysical Journal*, *904*(1), 35. <https://doi.org/10.3847/1538-4357/abbd98>
- Prochaska, J. X., Hennawi, J. F., Westfall, K. B., Cooke, R. J., Wang, F., Hsyu, T., Davies, F. B., Farina, E. P., & Pelliccia, D. (2020). Pypelt: The python spectroscopic data reduction pipeline. *Journal of Open Source Software*, *5*(56), 2308. <https://doi.org/10.21105/joss.02308>
- Ravi, V., Law, C. J., Li, D., Aggarwal, K., Burke-Spolaor, S., Connor, L., Lazio, T. J. W., Simard, D., Somalwar, J., & Tendulkar, S. P. (2021). The host galaxy and persistent radio counterpart of FRB 20201124A. *arXiv e-Prints*, arXiv:2106.09710. <http://arxiv.org/abs/2106.09710>