

Madym: A C++ toolkit for quantitative DCE-MRI analysis

Michael Berks^{*1}, Geoff J M Parker^{2, 3}, Ross Little¹, and Sue Cheung¹

¹ Quantitative Biomedical Imaging Laboratory, Division of Cancer Sciences, Manchester, UK ² Centre for Medical Image Computing, University College London, London, UK ³ Bioxydyn Ltd, Manchester, UK

DOI: [10.21105/joss.03523](https://doi.org/10.21105/joss.03523)

Software

- [Review](#) ↗
- [Repository](#) ↗
- [Archive](#) ↗

Editor: [Elizabeth DuPre](#) ↗

Reviewers:

- [@agahkarakuzu](#)
- [@matteomancini](#)

Submitted: 24 June 2021

Published: 22 July 2021

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Summary

In dynamic contrast-enhanced magnetic resonance imaging (DCE-MRI) a sequence of MRI images are acquired to measure the passage of a contrast-agent within a tissue of interest. Quantitative DCE-MRI (DCE-MRI), in which one or more tracer-kinetic models are fitted to the contrast-agent concentration time-series, enables the estimation of clinically useful parameters of tissue microvasculature.

Madym is a C++ toolkit for quantitative DCE-MRI analysis developed at the University of Manchester. It comprises a set of command line tools and a graphical user-interface based on an extendable C++ library. It is cross-platform, and requires few external libraries to build from source. Pre-built binaries for Windows, MacOS and Linux are available on request. We have also developed complementary interfaces in Matlab and python (available in separate open-source repositories ([M. Berks, 2021b](#)), ([M. Berks, 2021c](#))), that allow the flexibility of developing in those scripting languages, while allowing C++ to do the heavy-duty computational work of tracer-kinetic model fitting.

Statement of need

Madym has been designed with the following principles:

- **Ease-of-use:** Madym supports many advanced features for DCE-MRI analysis, however the tools have been designed to be usable by anyone, including clinical scientists with no software/programming knowledge. Extensive documentation is provided on the project wiki ([M. Berks, 2021a](#)), and an example test set is included with the toolkit, including walk through instructions of how to perform a standard analysis on these data.
- **Reproducible research:** even the simplest DCE-MRI analysis pipeline requires configuring many parameters (*ie* typically more than 20), which in some packages may be implicitly encoded in sub-methods, and may therefore differ in non-transparent ways between different implementations of the same analysis pipeline. Wherever possible, Madym exposes all parameters as configurable options, with a single source file specifying their default values used throughout the toolkit. A consistent interface is provided for configuring individual options, either via input config files, setting options directly at the command-line or adjusting interactively in the GUI. Whenever an analysis is run, the complete configuration - including the final set of parameter option values, the version of Madym used and the machine ID on which the analysis was run is saved with the output results.

^{*}corresponding author

Thus Madym provides both flexibility in configuring analyses to individual datasets, while supporting reproducibility with a complete record of how results were obtained. In doing so we support the aims of the ISMRM (International Society for Magnetic Resonance in Medicine) Reproducible Research Study Group (Stikov et al., 2019), and have listed Madym on the ISMRM MR-hub (ISMRM, 2021).

- Extensibility: Madym includes several of the most commonly used tracer-kinetic models as standard, including the Patlak (Patlak et al., 1983), extend-Tofts (Tofts, 1997) and two compartment exchange models (Sourbron et al., 2009), as well as more complex models for fitting contrast-agents that are actively metabolised by tissue and/or require dual vascular supply functions (Michael Berks et al., 2021). However these are by no means an exhaustive list and, by decoupling model optimisation from the model definitions, the toolkit has been designed to make adding new models very easy, simply by sub-classing the main abstract model class. Instructions for doing so are provided in the project wiki. Extending T_1 fitting methods (currently variable flip-angle and inversion recovery methods are supported), or even adding a new command-line tool, are designed and documented in the same way.
- Performance: Madym is designed with the aim of voxel-wise model fitting (where a model is fitted to individual tissue voxels rather than spatially averaged regions-of-interest). 3D MRI images have many hundreds of thousands of voxels (eg a typical image may have dimensions $128 \times 128 \times 40 = 655,360$ voxels). By using C++ and externally developed open-source optimisation library (ALGLIB, (Bochkanov, 2019)), on a standard desktop Madym requires $\approx 10\mu s$ per voxel to estimate baseline T_1 (allowing T_1 mapping of whole volumes in a few seconds) and $< 30ms$ per voxel to fit the extended-Tofts model (so a typical tumour of 500-1,000 voxels can be analysed in 20-30 seconds, while whole organs can be fitted in a few hours).

Madym has been developed over approximately 20 years and has been used to perform DCE-MRI analyses in more than 20 research papers and many more conference abstracts (landmark papers include (Jayson et al., 2018) and (O'Connor et al., 2012), see the project wiki for a more complete list). Until this year, these used previous non-open source versions, however the first paper using Madym as an open-source toolkit has just been published (Michael Berks et al., 2021), and we hope will be the first of many in the future.

Acknowledgements

Code updates to the open source version of Madym have been made entirely by Michael Berks, with significant input on the design and features of the new version from the remaining authors. We would also like to thank Gio Buonnacorsi, Anita Banerji and Angela Caunce for their work in developing previous non-public versions, following Geoff Parker's original design. While working on Madym, Michael Berks was funded by the following grants: Cancer Research UK (CRUK) Clinician Scientist award (grant C19221/A22746); CRUK and EPSRC Cancer Imaging Centre in Cambridge and Manchester funding to The University of Manchester (grant C8742/A18097).

References

- Berks, M. (2021a). Madym: A c++ DCE-MRI toolkit - project wiki. In *GitLab wiki*. GitLab. https://gitlab.com/manchester_qbi/manchester_qbi_public/madym_cxx/-/wikis

- 81 Berks, M. (2021b). Madym matlab: A matlab interface the madym c++ DCE-MRI toolkit. In
 82 *GitLab repository*. GitLab. [https://gitlab.com/manchester_qbi/manchester_qbi_public/](https://gitlab.com/manchester_qbi/manchester_qbi_public/madym_matlab)
 83 [madym_matlab](https://gitlab.com/manchester_qbi/manchester_qbi_public/madym_matlab)
- 84 Berks, M. (2021c). Madym python: A python interface the madym c++ DCE-MRI toolkit. In
 85 *GitLab repository*. GitLab. [https://gitlab.com/manchester_qbi/manchester_qbi_public/](https://gitlab.com/manchester_qbi/manchester_qbi_public/madym_python)
 86 [madym_python](https://gitlab.com/manchester_qbi/manchester_qbi_public/madym_python)
- 87 Berks, Michael, Little, R. A., Watson, Y., Cheung, S., Datta, A., O'Connor, J. P., Scaramuzza,
 88 D., & Parker, G. J. (2021). A model selection framework to quantify microvascular liver
 89 function in gadoxetate-enhanced MRI: Application to healthy liver, diseased tissue, and
 90 hepatocellular carcinoma. *Magnetic Resonance in Medicine*.
- 91 Bochkhanov, S. (2019). *ALGLIB: C++ optimisation library*. <http://www.alglib.net>
- 92 ISMRM. (2021). MR-hub: Open-access software tools for the ISMRM community. In *GitHub*
 93 *pages*. GitHub. <https://ismrm.github.io/mrhub/>
- 94 Jayson, G. C., Zhou, C., Backen, A., Horsley, L., Marti-Marti, K., Shaw, D., Mescallado, N.,
 95 Clamp, A., Saunders, M. P., Valle, J. W., & others. (2018). Plasma Tie2 is a tumor
 96 vascular response biomarker for VEGF inhibitors in metastatic colorectal cancer. *Nature*
 97 *Communications*, 9(1), 4672.
- 98 O'Connor, J. P., Jackson, A., Parker, G. J., Roberts, C., & Jayson, G. C. (2012). Dynamic
 99 contrast-enhanced MRI in clinical trials of antivascular therapies. *Nature Reviews Clinical*
 100 *Oncology*, 9(3), 167.
- 101 Patlak, C. S., Blasberg, R. G., & Fenstermacher, J. D. (1983). Graphical evaluation of blood-
 102 to-brain transfer constants from multiple-time uptake data. *Journal of Cerebral Blood*
 103 *Flow and Metabolism*, 3(1), 1–7.
- 104 Sourbron, S., Ingris, M., Siefert, A., Reiser, M., & Herrmann, K. (2009). Quantification
 105 of cerebral blood flow, cerebral blood volume, and blood–brain-barrier leakage with DCE-
 106 MRI. *Magnetic Resonance in Medicine*, 62(1), 205–217. [https://doi.org/10.1002/mrm.](https://doi.org/10.1002/mrm.22005)
 107 [22005](https://doi.org/10.1002/mrm.22005)
- 108 Stikov, N., Trzasko, J. D., & Bernstein, M. A. (2019). Reproducibility and the future of MRI
 109 research. *Magnetic Resonance in Medicine*, 82(6), 1981–1983. <https://doi.org/https://doi.org/10.1002/mrm.27939>
 110 <https://doi.org/https://doi.org/10.1002/mrm.27939>
- 111 Tofts, P. S. (1997). Modeling tracer kinetics in dynamic gd-DTPA MR imaging. *Journal of*
 112 *Magnetic Resonance Imaging*, 7(1), 91–101. <https://doi.org/10.1002/jmri.1880070113>