

RHRT: An R package to assess Heart Rate Turbulence

- ² Valeria Blesius*¹ and Andreas Dominik^{†1}
- 1 THM University of Applied Sciences, Giessen, Germany

DOI: 10.21105/joss.03540

Software

- Review 🗗
- Repository 🗗
- Archive ♂

Editor: Brian McFee ♂

Reviewers:

- @brunaw
- @rrrlw

Submitted: 23 June 2021 **Published:** 26 July 2021

License

Authors of papers retain 15 copyright and release the work 16 under a Creative Commons 17 Attribution 4.0 International 18 License (CC BY 4.0). 19

Summary

Heart Rate Turbulence (HRT) is a naturally occurring phenomenon of the heart that was first described by Schmidt et al. (1999). After a premature ventricular contraction (VPC) – an atypical heart beat that originates in the ventricles and occurs with a lower latency than a regular beat – the heart rate fluctuates. This fluctuation consists of a fast increase and later decrease in heart rate. This variability of interval lengths is an indicator for the condition of the autonomic nervous system: While no reaction to the premature ventricular beat and therefore no variation suggests an underlying pathology, a distinctive turbulence is considered healthy. Therefore, HRT can be used in medicine to determine the risk status of a person especially with certain diseases or conditions, e.g. after a myocardial infarction. (reviewed in Bauer et al., 2008)

RHRT finds occurrences of HRT in heart beat interval data, calculates the most used parameters (Turbulence Onset (TO), Turbulence Slope (TS), Turbulence Timing (TT) and normalised Turbulence Slope (nTS)) and plots the results. The package works best and fastest when given annotation data, but can also find HRT based on commonly accepted filtering rules that were first published in Grimm et al. (2003). Most filtering parameters and calculation methods can be freely adjusted to enable research on the methodology itself. In addition to parameter calculation, RHRT can classify the data into common risk categories (HRT0-2 and HRTA-C) and estimate the reliability of the results based on the number and parameter values of the HRTs.

The package can be found on [CRAN] and GitHub.

Statement of need

- Since it reflects the status of the autonomic nervous system, HRT is a feasible method to estimate the health risk of a person (Lombardi, 2011). HRT analysis is already used for risk stratification in the clinical practice, although the optimal methodology for HRT assessment has still not been systematically set (Blesius et al., 2020). RHRT is the first R package for HRT analysis and aims to enable further research about the usefulness of HRT and its methodology.
- Minimal example
- 32 To install RHRT use
- install.packages("RHRT")

^{*}corresponding author

[†]co-author



- ₃₄ for the version on CRAN. To install the continuosly developed version on GitHub you can use
- 35 the devtools package:
- 36 devtools::install_github("VBlesius/RHRT/RHRT")

have a look at the data and the parameters

- 37 The most straightforward use of RHRT is to scan your interval data for valid HRTs and analyse
- 38 the results:

plot(hrtl)

```
library(RHRT)
## scan your interval data and save the results as an HRTList
### the input should be a numeric vector consisting of RR interval data
### testdataLong is dummy data included in the package
hrtl <- vectorToHRT(testdataLong)

## get the HRT class of your data
getResults(hrtl, type = "class")
#> [1] "HRTO"
```




Figure 1: Plot of an HRTList Object: the plot resembles the standard visualisation of HRT – a tachogram – in which the indices of the intervals are plotted against their lengths. The tachogram and HRT parameters TO and TS are drawn in black, red and blue, respectively, while the tachograms of all underlying HRTs are drawn in grey in the background. The plot is zoomed in by default to show the HRT parameter values more precisely, so the intervals before and after the VPC are outside the plot range. RR interval: interval between two heartbeats measured between the R-peaks in the ECG, couplRR: coupling interval (interval between last sinus induced contraction and VPC), compRR: compensatory interval (interval between VPC and following sinus induced contraction), TO: Turbulence Onset, TS: Turbulence Slope.

- 39 More examples and a detailed description of the objects and functions can be found in the
- vignette of the package.



41 Acknowledgements

- 42 We are sincerely grateful to Christopher Schölzel for testing the package and providing excel-
- lent input through the course of its development.

44 References

- Bauer, A., Malik, M., Schmidt, G., Barthel, P., Bonnemeier, H., Cygankiewicz, I., Guzik, P., Lombardi, F., Müller, A., Oto, A., Schneider, R., Watanabe, M., Wichterle, D., & Zareba, W. (2008). Heart rate turbulence: Standards of measurement, physiological interpretation, and clinical use: International Society for Holter and Noninvasive Electrophysiology Consensus. *Journal of the American College of Cardiology*, *52*(17), 1353–1365. https://doi.org/10.1016/j.jacc.2008.07.041
- Blesius, V., Schölzel, C., Ernst, G., & Dominik, A. (2020). HRT assessment reviewed: A systematic review of heart rate turbulence methodology. *Physiological Measurement*, 41(8), 08TR01. https://doi.org/10.1088/1361-6579/ab98b3
- Grimm, W., Sharkova, J., Christ, M., Schneider, R., Schmidt, G., & Maisch, B. (2003).
 Heart rate turbulence following ventricular premature beats in healthy controls. *Annals of Noninvasive Electrocardiology*, 8(2), 127–131.
- Lombardi, F. (2011). Origin of heart rate variability and turbulence: An appraisal of autonomic modulation of cardiovascular function. *Clinical and Translational Physiology*, *2*, 95. https://doi.org/10.3389/fphys.2011.00095
- Schmidt, G., Malik, M., Barthel, P., Schneider, R., Ulm, K., Rolnitzky, L., Camm, A. J.,
 Bigger Jr, J. T., & Schömig, A. (1999). Heart-rate turbulence after ventricular premature
 beats as a predictor of mortality after acute myocardial infarction. *The Lancet*, 353(9162),
 1390–1396. https://doi.org/10.1016/S0140-6736(98)08428-1