

CS 228 : Logic in Computer Science

S. Krishna

Recap

- ▶ Signature τ : set containing relations and constants

Recap

- ▶ Signature τ : set containing relations and constants
- ▶ Each relation in τ has an arity

Recap

- ▶ Signature τ : set containing relations and constants
- ▶ Each relation in τ has an arity
- ▶ A FO formula is written over some signature τ ; that is, it uses the relations and constants from τ . It also uses variables denoted x_i , boolean connectives and quantifiers \forall, \exists .

Recap

- ▶ Signature τ : set containing relations and constants
- ▶ Each relation in τ has an arity
- ▶ A FO formula is written over some signature τ ; that is, it uses the relations and constants from τ . It also uses variables denoted x_i , boolean connectives and quantifiers \forall, \exists .
- ▶ A relation R of arity k is used in the formula as $R(t_1, \dots, t_k)$ where t_i 's are variables or constants

Recap

- ▶ Signature τ : set containing relations and constants
- ▶ Each relation in τ has an arity
- ▶ A FO formula is written over some signature τ ; that is, it uses the relations and constants from τ . It also uses variables denoted x_i , boolean connectives and quantifiers \forall, \exists .
- ▶ A relation R of arity k is used in the formula as $R(t_1, \dots, t_k)$ where t_i 's are variables or constants
- ▶ Equality $t_1 = t_2$ is available irrespective of τ

Recap

- ▶ Signature τ : set containing relations and constants
- ▶ Each relation in τ has an arity
- ▶ A FO formula is written over some signature τ ; that is, it uses the relations and constants from τ . It also uses variables denoted x_i , boolean connectives and quantifiers \forall, \exists .
- ▶ A relation R of arity k is used in the formula as $R(t_1, \dots, t_k)$ where t_i 's are variables or constants
- ▶ Equality $t_1 = t_2$ is available irrespective of τ
- ▶ To make sense out of a formula, we need structures

Recap

- ▶ A structure of signature τ consists of a universe, and assigns meanings to all the entities of τ

Recap

- ▶ A structure of signature τ consists of a universe, and assigns meanings to all the entities of τ
- ▶ So, if R is a k -ary relational symbol of τ , the structure specifies which k -tuples of elements from the universe are legitimate relations for R
- ▶ If c is a constant in τ , the structure also maps c to some fixed element of the universe

Recap

- ▶ A structure of signature τ consists of a universe, and assigns meanings to all the entities of τ
- ▶ So, if R is a k -ary relational symbol of τ , the structure specifies which k -tuples of elements from the universe are legitimate relations for R
- ▶ If c is a constant in τ , the structure also maps c to some fixed element of the universe
- ▶ So, structures of τ **give life** to τ

Recap

- ▶ A structure of signature τ consists of a universe, and assigns meanings to all the entities of τ
- ▶ So, if R is a k -ary relational symbol of τ , the structure specifies which k -tuples of elements from the universe are legitimate relations for R
- ▶ If c is a constant in τ , the structure also maps c to some fixed element of the universe
- ▶ So, structures of τ **give life** to τ
- ▶ Structures also tell you the set of values your variables x_i can assume : these are the elements from the universe

Recap

- ▶ A structure of signature τ consists of a universe, and assigns meanings to all the entities of τ
- ▶ So, if R is a k -ary relational symbol of τ , the structure specifies which k -tuples of elements from the universe are legitimate relations for R
- ▶ If c is a constant in τ , the structure also maps c to some fixed element of the universe
- ▶ So, structures of τ **give life** to τ
- ▶ Structures also tell you the set of values your variables x_i can assume : these are the elements from the universe
- ▶ A structure in PL will just consist of the universe $\{0, 1\}$, since there is no signature. All variables assume values from this boolean universe.

Satisfiability in PL and FO

- ▶ The satisfiability of a PL formula depends on the existence of an assignment satisfying it; likewise, the satisfiability of a FO formula φ over signature τ depends on the existence of a structure \mathcal{A} of τ such that φ is true on \mathcal{A} .

Free and Bound Variables

- ▶ For a wff $\varphi = \forall x\psi$, ψ is said to be the **scope** of the quantifier x

Free and Bound Variables

- ▶ For a wff $\varphi = \forall x\psi$, ψ is said to be the **scope** of the quantifier x
- ▶ Every occurrence of x in $\forall x\psi$ is **bound**
- ▶ Any occurrence of x which is not bound is called **free**


Free and Bound Variables

- ▶ For a wff $\varphi = \forall x\psi$, ψ is said to be the **scope** of the quantifier x
- ▶ Every occurrence of x in $\forall x\psi$ is **bound**
- ▶ Any occurrence of x which is not bound is called **free**
- ▶ $\varphi = P(x, y) \rightarrow \forall x((\forall yR(x, y)) \rightarrow Q(x, y))$
 - ▶ y is free in $Q(x, y)$ and bound in $R(x, y)$,
 - ▶ x is free in $P(x, y)$, and bound in $Q(x, y), R(x, y)$

Free and Bound Variables

- ▶ For a wff $\varphi = \forall x\psi$, ψ is said to be the **scope** of the quantifier x
- ▶ Every occurrence of x in $\forall x\psi$ is **bound**
- ▶ Any occurrence of x which is not bound is called **free**
- ▶ $\varphi = P(x, y) \rightarrow \forall x((\forall yR(x, y)) \rightarrow Q(x, y))$
 - ▶ y is free in $Q(x, y)$ and bound in $R(x, y)$,
 - ▶ x is free in $P(x, y)$, and bound in $Q(x, y), R(x, y)$
- ▶ Given φ , denote by $\varphi(x_1, \dots, x_n)$, that x_1, \dots, x_n are the free variables of φ , also $free(\varphi)$
- ▶ A **sentence** is a formula φ **none** of whose variables are **free**


$$P(x, y) \rightarrow \forall x((\forall y R(x, y)) \rightarrow Q(x, y))$$




$$P(x, y) \rightarrow \forall x((\forall y R(x, y)) \rightarrow Q(x, y))$$




$$P(x, y) \rightarrow \forall x((\forall y R(x, y)) \rightarrow Q(x, y))$$




$$P(x, y) \rightarrow \forall x((\forall yR(x, y)) \rightarrow Q(x, y))$$




$$P(x, y) \rightarrow \forall x((\forall y R(x, y)) \rightarrow Q(x, y))$$




$$P(x, y) \rightarrow \forall x((\forall y R(x, y)) \rightarrow Q(x, y))$$




$$P(x, y) \rightarrow \forall x((\forall y R(x, y)) \rightarrow Q(x, y))$$




$$P(x, y) \rightarrow \forall x((\forall y R(x, y)) \rightarrow Q(x, y))$$



$$P(x, y) \rightarrow \forall x((\forall y R(x, y)) \rightarrow Q(x, y))$$




$$P(x, y) \rightarrow \forall x((\forall y R(x, y)) \rightarrow Q(x, y))$$




$$P(x, y) \rightarrow \forall x((\forall y R(x, y)) \rightarrow Q(x, y))$$


$$\varphi(s, t) = P(s, t) \rightarrow \forall x((\forall y R(x, y)) \rightarrow Q(x, t))$$




$$\forall x(R(x, y) \rightarrow \forall yP(y))$$




$$\forall x(R(x, y) \rightarrow \forall yP(y))$$




$$\forall x(R(x, y) \rightarrow \forall yP(y))$$




$$\forall x(R(x, y) \rightarrow \forall yP(y))$$




$$\forall x(R(x, y) \rightarrow \forall yP(y))$$



$$\forall x(R(x, y) \rightarrow \forall yP(y))$$



$$\forall x(R(x, y) \rightarrow \forall yP(y))$$



$$\varphi(t) = \forall x(R(x, t) \rightarrow \forall yP(y))$$

Assignments on τ -structures

Assignments

For a τ -structure \mathcal{A} , an assignment over \mathcal{A} is a function $\alpha : \mathcal{V} \rightarrow u(\mathcal{A})$ that assigns every variable $x \in \mathcal{V}$ a value $\alpha(x) \in u(\mathcal{A})$. If t is a constant symbol c , then $\alpha(t)$ is $c^{\mathcal{A}}$

Assignments

Binding on a Variable

For an assignment α over \mathcal{A} , $\alpha[x \mapsto a]$ is the assignment

$$\alpha[x \mapsto a](y) = \begin{cases} \alpha(y), & y \neq x, \\ a, & y = x \end{cases}$$

Assignments

Binding on a Variable

For an assignment α over \mathcal{A} , $\alpha[x \mapsto a]$ is the assignment

$$\alpha[x \mapsto a](y) = \begin{cases} \alpha(y), & y \neq x, \\ a, & y = x \end{cases}$$

Let $u(\mathcal{A}) = \{a, b, c, d\}$, and consider assignment $\alpha : \{x, y, z\} \rightarrow u(\mathcal{A})$ defined by $\alpha(x) = d, \alpha(y) = b, \alpha(z) = c$. Then,

- ▶ $\alpha[x \mapsto a]$ is the assignment α' where $\alpha'(x) = a, \alpha'(y) = \alpha(y), \alpha'(z) = \alpha(z)$.
- ▶ $\alpha[x \mapsto c]$ is the assignment α'' where $\alpha''(x) = c, \alpha''(y) = \alpha(y), \alpha''(z) = \alpha(z)$.

Satisfaction

We define the relation $\mathcal{A} \models_{\alpha} \varphi$ (read as φ is true in \mathcal{A} under the assignment α) inductively:

Satisfaction

We define the relation $\mathcal{A} \models_{\alpha} \varphi$ (read as φ is true in \mathcal{A} under the assignment α) inductively:

► $\mathcal{A} \not\models_{\alpha} \perp$

Satisfaction

We define the relation $\mathcal{A} \models_{\alpha} \varphi$ (read as φ is true in \mathcal{A} under the assignment α) inductively:

- ▶ $\mathcal{A} \not\models_{\alpha} \perp$
- ▶ $\mathcal{A} \models_{\alpha} t_1 = t_2$ iff $\alpha(t_1) = \alpha(t_2)$

Satisfaction

We define the relation $\mathcal{A} \models_{\alpha} \varphi$ (read as φ is true in \mathcal{A} under the assignment α) inductively:

- ▶ $\mathcal{A} \not\models_{\alpha} \perp$
- ▶ $\mathcal{A} \models_{\alpha} t_1 = t_2$ iff $\alpha(t_1) = \alpha(t_2)$
- ▶ $\mathcal{A} \models_{\alpha} R(t_1, \dots, t_k)$ iff $(\alpha(t_1), \dots, \alpha(t_k)) \in R^{\mathcal{A}}$

Satisfaction

We define the relation $\mathcal{A} \models_{\alpha} \varphi$ (read as φ is true in \mathcal{A} under the assignment α) inductively:

- ▶ $\mathcal{A} \not\models_{\alpha} \perp$
- ▶ $\mathcal{A} \models_{\alpha} t_1 = t_2$ iff $\alpha(t_1) = \alpha(t_2)$
- ▶ $\mathcal{A} \models_{\alpha} R(t_1, \dots, t_k)$ iff $(\alpha(t_1), \dots, \alpha(t_k)) \in R^{\mathcal{A}}$
- ▶ $\mathcal{A} \models_{\alpha} (\varphi \rightarrow \psi)$ iff $\mathcal{A} \not\models_{\alpha} \varphi$ or $\mathcal{A} \models_{\alpha} \psi$

Satisfaction

We define the relation $\mathcal{A} \models_{\alpha} \varphi$ (read as φ is true in \mathcal{A} under the assignment α) inductively:

- ▶ $\mathcal{A} \not\models_{\alpha} \perp$
- ▶ $\mathcal{A} \models_{\alpha} t_1 = t_2$ iff $\alpha(t_1) = \alpha(t_2)$
- ▶ $\mathcal{A} \models_{\alpha} R(t_1, \dots, t_k)$ iff $(\alpha(t_1), \dots, \alpha(t_k)) \in R^{\mathcal{A}}$
- ▶ $\mathcal{A} \models_{\alpha} (\varphi \rightarrow \psi)$ iff $\mathcal{A} \not\models_{\alpha} \varphi$ or $\mathcal{A} \models_{\alpha} \psi$
- ▶ $\mathcal{A} \models_{\alpha} (\forall x)\varphi$ iff for every $a \in u(\mathcal{A})$, $\mathcal{A} \models_{\alpha[x \mapsto a]} \varphi$

Satisfaction

We define the relation $\mathcal{A} \models_{\alpha} \varphi$ (read as φ is true in \mathcal{A} under the assignment α) inductively:

- ▶ $\mathcal{A} \not\models_{\alpha} \perp$
- ▶ $\mathcal{A} \models_{\alpha} t_1 = t_2$ iff $\alpha(t_1) = \alpha(t_2)$
- ▶ $\mathcal{A} \models_{\alpha} R(t_1, \dots, t_k)$ iff $(\alpha(t_1), \dots, \alpha(t_k)) \in R^{\mathcal{A}}$
- ▶ $\mathcal{A} \models_{\alpha} (\varphi \rightarrow \psi)$ iff $\mathcal{A} \not\models_{\alpha} \varphi$ or $\mathcal{A} \models_{\alpha} \psi$
- ▶ $\mathcal{A} \models_{\alpha} (\forall x)\varphi$ iff for every $a \in u(\mathcal{A})$, $\mathcal{A} \models_{\alpha[x \mapsto a]} \varphi$
- ▶ $\mathcal{A} \models_{\alpha} (\exists x)\varphi$ iff there is some $a \in u(\mathcal{A})$, $\mathcal{A} \models_{\alpha[x \mapsto a]} \varphi$

Last two cases, α has no effect on the value of x . Thus, assignments matter **only** to free variables.

Example of Satisfaction

$$\mathcal{G} = (\{1, 2, 3\}, E^{\mathcal{G}} = \{(1, 2), (2, 1), (2, 3), (3, 2)\})$$

- For any assignment α , $\mathcal{G} \models_{\alpha} \forall x \forall y (E(x, y) \rightarrow E(y, x))$ iff

Example of Satisfaction

$$\mathcal{G} = (\{1, 2, 3\}, E^{\mathcal{G}} = \{(1, 2), (2, 1), (2, 3), (3, 2)\})$$

- For any assignment α , $\mathcal{G} \models_{\alpha} \forall x \forall y (E(x, y) \rightarrow E(y, x))$ iff
for all $a \in \{1, 2, 3\}$, $\mathcal{G} \models_{\alpha[x \mapsto a]} \forall y (E(x, y) \rightarrow E(y, x))$ iff

Example of Satisfaction

$$\mathcal{G} = (\{1, 2, 3\}, E^{\mathcal{G}} = \{(1, 2), (2, 1), (2, 3), (3, 2)\})$$

- For any assignment α , $\mathcal{G} \models_{\alpha} \forall x \forall y (E(x, y) \rightarrow E(y, x))$ iff
for all $a \in \{1, 2, 3\}$, $\mathcal{G} \models_{\alpha[x \mapsto a]} \forall y (E(x, y) \rightarrow E(y, x))$ iff
for every $a, b \in \{1, 2, 3\}$, $\mathcal{G} \models_{\alpha[x \mapsto a, y \mapsto b]} (E(x, y) \rightarrow E(y, x))$

Example of Satisfaction

$$\mathcal{G} = (\{1, 2, 3\}, E^{\mathcal{G}} = \{(1, 2), (2, 1), (2, 3), (3, 2)\})$$

- ▶ For any assignment α , $\mathcal{G} \models_{\alpha} \forall x \forall y (E(x, y) \rightarrow E(y, x))$ iff
for all $a \in \{1, 2, 3\}$, $\mathcal{G} \models_{\alpha[x \mapsto a]} \forall y (E(x, y) \rightarrow E(y, x))$ iff
for every $a, b \in \{1, 2, 3\}$, $\mathcal{G} \models_{\alpha[x \mapsto a, y \mapsto b]} (E(x, y) \rightarrow E(y, x))$
 - ▶ for $\alpha_1 : \alpha_1(x) = 1, \alpha_1(y) = 1, \mathcal{G} \models_{\alpha_1} (E(x, y) \rightarrow E(y, x))$,
 - ▶ for $\alpha_2 : \alpha_2(x) = 1, \alpha_2(y) = 2, \mathcal{G} \models_{\alpha_2} (E(x, y) \rightarrow E(y, x))$,
 - ▶ for $\alpha_3 : \alpha_3(x) = 1, \alpha_3(y) = 3, \mathcal{G} \models_{\alpha_3} (E(x, y) \rightarrow E(y, x))$,
 - ▶ for $\alpha_4 : \alpha_4(x) = 2, \alpha_4(y) = 1, \mathcal{G} \models_{\alpha_4} (E(x, y) \rightarrow E(y, x))$,
 - ▶ \vdots
 - ▶ for $\alpha_9 : \alpha_9(x) = 3, \alpha_9(y) = 3, \mathcal{G} \models_{\alpha_9} (E(x, y) \rightarrow E(y, x))$

Example of Satisfaction

$$\mathcal{G} = (\{1, 2, 3\}, E^{\mathcal{G}} = \{(1, 2), (2, 1), (2, 3), (3, 2)\})$$

- ▶ For any assignment α , $\mathcal{G} \models_{\alpha} \forall x \forall y (E(x, y) \rightarrow E(y, x))$ iff
for all $a \in \{1, 2, 3\}$, $\mathcal{G} \models_{\alpha[x \mapsto a]} \forall y (E(x, y) \rightarrow E(y, x))$ iff
for every $a, b \in \{1, 2, 3\}$, $\mathcal{G} \models_{\alpha[x \mapsto a, y \mapsto b]} (E(x, y) \rightarrow E(y, x))$
 - ▶ for $\alpha_1 : \alpha_1(x) = 1, \alpha_1(y) = 1, \mathcal{G} \models_{\alpha_1} (E(x, y) \rightarrow E(y, x))$,
 - ▶ for $\alpha_2 : \alpha_2(x) = 1, \alpha_2(y) = 2, \mathcal{G} \models_{\alpha_2} (E(x, y) \rightarrow E(y, x))$,
 - ▶ for $\alpha_3 : \alpha_3(x) = 1, \alpha_3(y) = 3, \mathcal{G} \models_{\alpha_3} (E(x, y) \rightarrow E(y, x))$,
 - ▶ for $\alpha_4 : \alpha_4(x) = 2, \alpha_4(y) = 1, \mathcal{G} \models_{\alpha_4} (E(x, y) \rightarrow E(y, x))$,
 - ▶ \vdots
 - ▶ for $\alpha_9 : \alpha_9(x) = 3, \alpha_9(y) = 3, \mathcal{G} \models_{\alpha_9} (E(x, y) \rightarrow E(y, x))$
- ▶ There is an assignment α which satisfies
 $\mathcal{G} \models_{\alpha} \exists x (E(x, y) \wedge E(x, z) \wedge y \neq z)$

Example of Satisfaction

$$\mathcal{G} = (\{1, 2, 3\}, E^{\mathcal{G}} = \{(1, 2), (2, 1), (2, 3), (3, 2)\})$$

- ▶ For any assignment α , $\mathcal{G} \models_{\alpha} \forall x \forall y (E(x, y) \rightarrow E(y, x))$ iff
for all $a \in \{1, 2, 3\}$, $\mathcal{G} \models_{\alpha[x \mapsto a]} \forall y (E(x, y) \rightarrow E(y, x))$ iff
for every $a, b \in \{1, 2, 3\}$, $\mathcal{G} \models_{\alpha[x \mapsto a, y \mapsto b]} (E(x, y) \rightarrow E(y, x))$
 - ▶ for $\alpha_1 : \alpha_1(x) = 1, \alpha_1(y) = 1, \mathcal{G} \models_{\alpha_1} (E(x, y) \rightarrow E(y, x))$,
 - ▶ for $\alpha_2 : \alpha_2(x) = 1, \alpha_2(y) = 2, \mathcal{G} \models_{\alpha_2} (E(x, y) \rightarrow E(y, x))$,
 - ▶ for $\alpha_3 : \alpha_3(x) = 1, \alpha_3(y) = 3, \mathcal{G} \models_{\alpha_3} (E(x, y) \rightarrow E(y, x))$,
 - ▶ for $\alpha_4 : \alpha_4(x) = 2, \alpha_4(y) = 1, \mathcal{G} \models_{\alpha_4} (E(x, y) \rightarrow E(y, x))$,
 - ▶ \vdots
 - ▶ for $\alpha_9 : \alpha_9(x) = 3, \alpha_9(y) = 3, \mathcal{G} \models_{\alpha_9} (E(x, y) \rightarrow E(y, x))$
- ▶ There is an assignment α which satisfies
 $\mathcal{G} \models_{\alpha} \exists x (E(x, y) \wedge E(x, z) \wedge y \neq z)$
 $\alpha(y) = 1, \alpha(z) = 3$, and consider $\alpha[x \mapsto 2]$.

Example of Satisfaction

$$\mathcal{G} = (\{1, 2, 3\}, E^{\mathcal{G}} = \{(1, 2), (2, 1), (2, 3), (3, 2)\})$$

- ▶ For any assignment α , $\mathcal{G} \models_{\alpha} \forall x \forall y (E(x, y) \rightarrow E(y, x))$ iff
for all $a \in \{1, 2, 3\}$, $\mathcal{G} \models_{\alpha[x \mapsto a]} \forall y (E(x, y) \rightarrow E(y, x))$ iff
for every $a, b \in \{1, 2, 3\}$, $\mathcal{G} \models_{\alpha[x \mapsto a, y \mapsto b]} (E(x, y) \rightarrow E(y, x))$
 - ▶ for $\alpha_1 : \alpha_1(x) = 1, \alpha_1(y) = 1, \mathcal{G} \models_{\alpha_1} (E(x, y) \rightarrow E(y, x))$,
 - ▶ for $\alpha_2 : \alpha_2(x) = 1, \alpha_2(y) = 2, \mathcal{G} \models_{\alpha_2} (E(x, y) \rightarrow E(y, x))$,
 - ▶ for $\alpha_3 : \alpha_3(x) = 1, \alpha_3(y) = 3, \mathcal{G} \models_{\alpha_3} (E(x, y) \rightarrow E(y, x))$,
 - ▶ for $\alpha_4 : \alpha_4(x) = 2, \alpha_4(y) = 1, \mathcal{G} \models_{\alpha_4} (E(x, y) \rightarrow E(y, x))$,
 - ▶ \vdots
 - ▶ for $\alpha_9 : \alpha_9(x) = 3, \alpha_9(y) = 3, \mathcal{G} \models_{\alpha_9} (E(x, y) \rightarrow E(y, x))$
- ▶ There is an assignment α which satisfies
 $\mathcal{G} \models_{\alpha} \exists x (E(x, y) \wedge E(x, z) \wedge y \neq z)$
 $\alpha(y) = 1, \alpha(z) = 3$, and consider $\alpha[x \mapsto 2]$.
- ▶ Check this: $\mathcal{G} \not\models \exists x \forall y E(x, y)$,

Example of Satisfaction

$$\mathcal{G} = (\{1, 2, 3\}, E^{\mathcal{G}} = \{(1, 2), (2, 1), (2, 3), (3, 2)\})$$

- ▶ For any assignment α , $\mathcal{G} \models_{\alpha} \forall x \forall y (E(x, y) \rightarrow E(y, x))$ iff
for all $a \in \{1, 2, 3\}$, $\mathcal{G} \models_{\alpha[x \mapsto a]} \forall y (E(x, y) \rightarrow E(y, x))$ iff
for every $a, b \in \{1, 2, 3\}$, $\mathcal{G} \models_{\alpha[x \mapsto a, y \mapsto b]} (E(x, y) \rightarrow E(y, x))$
 - ▶ for $\alpha_1 : \alpha_1(x) = 1, \alpha_1(y) = 1, \mathcal{G} \models_{\alpha_1} (E(x, y) \rightarrow E(y, x))$,
 - ▶ for $\alpha_2 : \alpha_2(x) = 1, \alpha_2(y) = 2, \mathcal{G} \models_{\alpha_2} (E(x, y) \rightarrow E(y, x))$,
 - ▶ for $\alpha_3 : \alpha_3(x) = 1, \alpha_3(y) = 3, \mathcal{G} \models_{\alpha_3} (E(x, y) \rightarrow E(y, x))$,
 - ▶ for $\alpha_4 : \alpha_4(x) = 2, \alpha_4(y) = 1, \mathcal{G} \models_{\alpha_4} (E(x, y) \rightarrow E(y, x))$,
 - ▶ \vdots
 - ▶ for $\alpha_9 : \alpha_9(x) = 3, \alpha_9(y) = 3, \mathcal{G} \models_{\alpha_9} (E(x, y) \rightarrow E(y, x))$
- ▶ There is an assignment α which satisfies
 $\mathcal{G} \models_{\alpha} \exists x (E(x, y) \wedge E(x, z) \wedge y \neq z)$
 $\alpha(y) = 1, \alpha(z) = 3$, and consider $\alpha[x \mapsto 2]$.
- ▶ Check this: $\mathcal{G} \not\models \exists x \forall y E(x, y)$, $\mathcal{G} \models \forall x \exists y E(x, y)$

Example of Satisfaction

- ▶ $\mathcal{W} = abaaa$ or,
 $\mathcal{W} = (\{0, \dots, 4\}, <^{\mathcal{W}}, S^{\mathcal{W}}, Q_a^{\mathcal{W}} = \{0, 2, 3, 4\}, Q_b^{\mathcal{W}} = \{1\})$

Example of Satisfaction

- ▶ $\mathcal{W} = abaaa$ or,
 $\mathcal{W} = (\{0, \dots, 4\}, <^{\mathcal{W}}, S^{\mathcal{W}}, Q_a^{\mathcal{W}} = \{0, 2, 3, 4\}, Q_b^{\mathcal{W}} = \{1\})$
 - ▶ There is an assignment α for which
 $\mathcal{W} \models_{\alpha} (Q_a(x) \wedge Q_a(y) \wedge S(x, y))$

Example of Satisfaction

- ▶ $\mathcal{W} = abaaa$ or,
 $\mathcal{W} = (\{0, \dots, 4\}, <^{\mathcal{W}}, S^{\mathcal{W}}, Q_a^{\mathcal{W}} = \{0, 2, 3, 4\}, Q_b^{\mathcal{W}} = \{1\})$
 - ▶ There is an assignment α for which
 $\mathcal{W} \models_{\alpha} (Q_a(x) \wedge Q_a(y) \wedge S(x, y))$
One possibility : $\alpha(x) = 2, \alpha(y) = 3$
 - ▶ There is no assignment α which satisfies
 $\exists x \exists y (Q_b(x) \wedge Q_b(y) \wedge x \neq y)$

Example of Satisfaction

- ▶ $\mathcal{W} = abaaa$ or,
 $\mathcal{W} = (\{0, \dots, 4\}, <^{\mathcal{W}}, S^{\mathcal{W}}, Q_a^{\mathcal{W}} = \{0, 2, 3, 4\}, Q_b^{\mathcal{W}} = \{1\})$
 - ▶ There is an assignment α for which
 $\mathcal{W} \models_{\alpha} (Q_a(x) \wedge Q_a(y) \wedge S(x, y))$
One possibility : $\alpha(x) = 2, \alpha(y) = 3$
 - ▶ There is no assignment α which satisfies
 $\exists x \exists y (Q_b(x) \wedge Q_b(y) \wedge x \neq y)$
 - ▶ Prove or disprove : $\mathcal{W} \models \exists x \forall y [Q_b(x) \wedge x < y \wedge Q_a(y)]$
 - ▶ Prove or disprove : $\mathcal{W} \models \exists x \forall y [Q_b(x) \wedge x < y \Rightarrow Q_a(y)]$

Satisfiability, Validity, Equivalence and Equisatisfiability

- ▶ A formula φ over a signature τ is said to be **satisfiable** iff for some τ -structure \mathcal{A} and assignment α , $\mathcal{A} \models_{\alpha} \varphi$

Satisfiability, Validity, Equivalence and Equisatisfiability

- ▶ A formula φ over a signature τ is said to be **satisfiable** iff for some τ -structure \mathcal{A} and assignment α , $\mathcal{A} \models_{\alpha} \varphi$
- ▶ A formula φ over a signature τ is said to be **valid** iff for every τ -structure \mathcal{A} and assignment α , $\mathcal{A} \models_{\alpha} \varphi$

Satisfiability, Validity, Equivalence and Equisatisfiability

- ▶ A formula φ over a signature τ is said to be **satisfiable** iff for some τ -structure \mathcal{A} and assignment α , $\mathcal{A} \models_{\alpha} \varphi$
- ▶ A formula φ over a signature τ is said to be **valid** iff for every τ -structure \mathcal{A} and assignment α , $\mathcal{A} \models_{\alpha} \varphi$
- ▶ Formulae $\varphi(x_1, \dots, x_n)$ and $\psi(x_1, \dots, x_n)$ are **equivalent** denoted $\varphi \equiv \psi$ iff for every \mathcal{A} and α , $\mathcal{A} \models_{\alpha} \varphi$ iff $\mathcal{A} \models_{\alpha} \psi$

Satisfiability, Validity, Equivalence and Equisatisfiability

- ▶ A formula φ over a signature τ is said to be **satisfiable** iff for some τ -structure \mathcal{A} and assignment α , $\mathcal{A} \models_{\alpha} \varphi$
- ▶ A formula φ over a signature τ is said to be **valid** iff for every τ -structure \mathcal{A} and assignment α , $\mathcal{A} \models_{\alpha} \varphi$
- ▶ Formulae $\varphi(x_1, \dots, x_n)$ and $\psi(x_1, \dots, x_n)$ are **equivalent** denoted $\varphi \equiv \psi$ iff for every \mathcal{A} and α , $\mathcal{A} \models_{\alpha} \varphi$ iff $\mathcal{A} \models_{\alpha} \psi$
- ▶ Consider $\varphi_1(x) = \forall y R(x, y)$ and $\varphi_2 = \exists x \forall y R(x, y)$.
- ▶ It is clear that whenever φ_2 is satisfiable on \mathcal{A} , $\mathcal{A} \models_{\alpha[x \mapsto a]} \forall y R(x, y)$, for some $a \in u(\mathcal{A})$. Then one can find the assignment α such that $\mathcal{A} \models_{\alpha} \varphi_1(x)$, $\alpha(x) = a$.
- ▶ Likewise, if $\mathcal{A} \models_{\alpha} \varphi_1(x)$, then

Satisfiability, Validity, Equivalence and Equisatisfiability

- ▶ A formula φ over a signature τ is said to be **satisfiable** iff for some τ -structure \mathcal{A} and assignment α , $\mathcal{A} \models_{\alpha} \varphi$
- ▶ A formula φ over a signature τ is said to be **valid** iff for every τ -structure \mathcal{A} and assignment α , $\mathcal{A} \models_{\alpha} \varphi$
- ▶ Formulae $\varphi(x_1, \dots, x_n)$ and $\psi(x_1, \dots, x_n)$ are **equivalent** denoted $\varphi \equiv \psi$ iff for every \mathcal{A} and α , $\mathcal{A} \models_{\alpha} \varphi$ iff $\mathcal{A} \models_{\alpha} \psi$
- ▶ Consider $\varphi_1(x) = \forall y R(x, y)$ and $\varphi_2 = \exists x \forall y R(x, y)$.
- ▶ It is clear that whenever φ_2 is satisfiable on \mathcal{A} , $\mathcal{A} \models_{\alpha[x \mapsto a]} \forall y R(x, y)$, for some $a \in u(\mathcal{A})$. Then one can find the assignment α such that $\mathcal{A} \models_{\alpha} \varphi_1(x)$, $\alpha(x) = a$.
- ▶ Likewise, if $\mathcal{A} \models_{\alpha} \varphi_1(x)$, then $\mathcal{A} \models_{\alpha'[x \mapsto \alpha(x)]} \varphi_2$, and $\alpha'(y)$ can be defined as $\alpha(y)$.
- ▶ Thus, $\varphi_1(x)$, φ_2 agree on satisfiability : equisatisfiable.

True or False?

For a formula φ and assignments α_1 and α_2 such that for every $x \in \text{free}(\varphi)$, $\alpha_1(x) = \alpha_2(x)$, $\mathcal{A} \models_{\alpha_1} \varphi$ iff $\mathcal{A} \models_{\alpha_2} \varphi$

True or False?

For a formula φ and assignments α_1 and α_2 such that for every $x \in \text{free}(\varphi)$, $\alpha_1(x) = \alpha_2(x)$, $\mathcal{A} \models_{\alpha_1} \varphi$ iff $\mathcal{A} \models_{\alpha_2} \varphi$

- For example, $\varphi(y) = \forall x(R(x, y) \rightarrow \forall zP(z))$

True or False?

For a formula φ and assignments α_1 and α_2 such that for every $x \in \text{free}(\varphi)$, $\alpha_1(x) = \alpha_2(x)$, $\mathcal{A} \models_{\alpha_1} \varphi$ iff $\mathcal{A} \models_{\alpha_2} \varphi$

- ▶ For example, $\varphi(y) = \forall x(R(x, y) \rightarrow \forall zP(z))$
- ▶ Consider two assignments α_1, α_2 such that $\alpha_1(y) = \alpha_2(y) = \alpha(\text{say})$

True or False?

For a formula φ and assignments α_1 and α_2 such that for every $x \in \text{free}(\varphi)$, $\alpha_1(x) = \alpha_2(x)$, $\mathcal{A} \models_{\alpha_1} \varphi$ iff $\mathcal{A} \models_{\alpha_2} \varphi$

- ▶ For example, $\varphi(y) = \forall x(R(x, y) \rightarrow \forall zP(z))$
- ▶ Consider two assignments α_1, α_2 such that $\alpha_1(y) = \alpha_2(y) = \alpha(\text{say})$
- ▶ Evaluate for all $a, b \in u(\mathcal{A})$, $R(a, \alpha) \rightarrow P(b)$

True or False?

For a formula φ and assignments α_1 and α_2 such that for every $x \in \text{free}(\varphi)$, $\alpha_1(x) = \alpha_2(x)$, $\mathcal{A} \models_{\alpha_1} \varphi$ iff $\mathcal{A} \models_{\alpha_2} \varphi$

- ▶ For example, $\varphi(y) = \forall x(R(x, y) \rightarrow \forall zP(z))$
- ▶ Consider two assignments α_1, α_2 such that $\alpha_1(y) = \alpha_2(y) = \alpha(\text{say})$
- ▶ Evaluate for all $a, b \in u(\mathcal{A})$, $R(a, \alpha) \rightarrow P(b)$
- ▶ $\mathcal{A} \models_{\alpha_1} \varphi$ iff $\mathcal{A} \models_{\alpha_2} \varphi$

True or False?

For a sentence φ , and any two assignments α_1 and α_2 , $\mathcal{A} \models_{\alpha_1} \varphi$ iff $\mathcal{A} \models_{\alpha_2} \varphi$

True or False?

For a sentence φ , and any two assignments α_1 and α_2 , $\mathcal{A} \models_{\alpha_1} \varphi$ iff $\mathcal{A} \models_{\alpha_2} \varphi$

No free variables!

Check SAT

- ▶ $\varphi = \exists x[(\forall y E(x, y)) \wedge \forall z[(\forall y E(z, y)) \rightarrow z = x]]$. Does φ evaluate to true under some graph structure?

Check SAT

- ▶ $\varphi = \exists x[(\forall y E(x, y)) \wedge \forall z[(\forall y E(z, y)) \rightarrow z = x]]$. Does φ evaluate to true under some graph structure?
- ▶ $\psi = \exists x[Q_a(x) \wedge \forall y[(y < x \wedge Q_b(y)) \rightarrow (z < x \wedge y < z \wedge Q_a(z))]]$. Does ψ evaluate to true under some word structure?