

Supplementary Materials for:

Abiraterone acetate plus prednisolone for metastatic patients starting hormone therapy: 5-year follow-up results from the STAMPEDE randomised trial (NCT00268476)

Authors: Nicholas D James, Noel W Clarke, Adrian Cook, Adnan Ali, Alex P Hoyle, Gert Attard, Chris D Brawley, Simon Chowdhury, William R Cross, David P Dearnaley, Johann S de Bono, Carlos Diaz Montana, Duncan Gilbert, Silke Gillesen, Clare Gilson, Rob J Jones, Ruth E Langley, Zafar I Malik, David J Matheson, Robin Millman, Chris C Parker, Cheryl Pugh, Hannah Rush, J Martin Russell, Dominic R Berthold, Michelle L Buckner, Malcolm D Mason, Alastair WS Ritchie, Alison J Birtle, Susannah J Brock, Prantik Das, Dan Ford, Joanna Gale, Warren Grant, Emma K Gray, Peter Hoskin, Mohammad M Khan, Caroline Manetta, Neil J McPhail, Joe M O'Sullivan, Omi Parikh, Carla Perna, Carmel J Pezaro, Andrew S Protheroe, Angus J Robinson, Sarah M Rudman, Denise J Sheehan, Narayanan N Srihari, Isabel Syndikus, Jacob Tanguay, Carys W Thomas, Salil Vengalil, John Wagstaff, James P Wylie, Mahesh KB Parmar, Matthew R Sydes

CONTENTS

CONTENTS	1
SUPPLEMENTARY TABLES.....	2
SUPPLEMENTARY FIGURES.....	5
STAMPEDE OVERSIGHT COMMITTEES, STAFF AND COLLABORATORS.....	8
PROTOCOL	9

SUPPLEMENTARY TABLES

Table S1: Post-progression treatment

Patients with reported progression*	SOC-alone		SOC+AAP		p
	n=437	(100%)	n=282	(100%)	
Any second-line treatment	405	(93%)	217	(77%)	<0.0001
Anti-androgen	316	(72%)	84	(30%)	<0.0001
Abiraterone	131	(30%)	10	(4%)	<0.0001
Enzalutamide	157	(36%)	46	(16%)	<0.0001
Docetaxel	199	(46%)	137	(49%)	0.424
Zoledronic acid	81	(19%)	51	(18%)	0.88
Dexamethasone	116	(27%)	66	(23%)	0.344
Prednisolone	93	(21%)	43	(15%)	0.044

* Biochemical progression, local progression, metastatic progression or skeletal-related event

Note: SOC = Standard-of-care, AAP = Abiraterone acetate + prednisolone

Table S2: Worst grade toxicity reported two years after randomisation*, and four years after randomisation*

Grade	<u>Two years</u>					<u>Four years</u>				
	<u>SOC-alone</u>		<u>SOC+AAP</u>		p	<u>SOC-alone</u>		<u>SOC+AAP</u>		p
	N=136	(100%)	N=291	(100%)		N=72	(100%)	N=197	(100%)	
0	36	(27%)	59	(21%)	0.29	5	(10%)	18	(12%)	0.56
1	63	(47%)	143	(50%)		25	(50%)	57	(38%)	
2	22	(17%)	64	(22%)		12	(24%)	51	(34%)	
3	12	(9%)	20	(7%)		8	(16%)	23	(15%)	
4	0	(0%)	0	(0%)		0	(0%)	1	(1%)	
5	0	(0%)	0	(0%)		0	(0%)	0	(0%)	
Missing	3	n/a	5	n/a		22	n/a	47	n/a	

* Toxicity data is collected until first progression in the SOC-alone group and until the end of abiraterone treatment for SOC+AAP patients. This determines the number of patients included in the denominator at two and four years, with patients counted as missing if data is expected but not reported within a 12-week window.

Note: SOC = Standard-of-care, AAP = Abiraterone acetate + prednisolone

Table S3: Overall survival and Failure-free survival, at time of primary analysis and at long-term follow-up

	<u>Primary analysis (2017)</u>		<u>Long-term follow-up analysis (2020)</u>	
	<u>SOC-alone</u> n=502	<u>SOC+AAP</u> n=501	<u>SOC-alone</u> n=502	<u>SOC+AAP</u> n=501
Overall survival				
Events	218	150	329	244
Proportion event-free at 5 yrs (95%CI)	0.43 (0.36,0.50)	0.58 (0.47,0.67)	0.41 (0.37,0.45)	0.60 (0.55,0.64)
SOC+AAP vs. SOC-alone, HR=(95%CI) p	0.61 (0.49,0.75)	<0.0001	0.60 (0.50,0.71)	<0.0001
p (proportional hazards)		0.33		0.78
RMST (months)	42 (40,44)	48 (46,50)	54 (51,57)	66 (63,69)
Failure-free survival				
Events	393	210	437	282
Proportion event-free at 5 yrs (95%CI)	0.18 (0.15,0.22)	0.51 (0.44,0.57)	0.13 (0.11,0.17)	0.45 (0.41,0.50)
SOC+AAP vs. SOC-alone, HR=(95%CI) p	0.31 (0.26,0.37)	<0.0001	0.34 (0.29,0.40)	<0.0001
p (proportional hazards)		0.0199		0.0001
RMST (months)	20 (18,23)	40 (38,43)	24 (21,27)	55 (51,59)

Note: SOC = Standard-of-care, AAP = Abiraterone acetate + prednisolone,


RMST = restricted mean “survival” (event-free) time, CI = confidence interval,

yr = years, HR = hazard ratio

SUPPLEMENTARY FIGURES

Figure S1: Time from randomisation to event for two secondary outcome measures

(A) Progression-free survival



(B) Metastatic progression-free survival





Figure S2: Time from randomisation to event for two secondary outcome measures

(A) Skeletal-related events



(B) Disease-specific survival






Figure S3: Time to from first progression event to starting of reported further treatment**(A) Abiraterone**

SOC					
Progressed	437	405	356	309	274
Censored	0	16	39	63	82
Started	0	16	42	65	81
SOC+AAP					
Progressed	282	259	230	212	185
Censored	0	22	49	67	92
Started	0	1	3	3	5

(B) Docetaxel (outside of treatment allocation)

SOC					
Progressed	437	380	310	261	230
Censored	0	14	37	53	65
Started	0	43	90	123	142
SOC+AAP					
Progressed	282	208	153	127	112
Censored	0	19	41	47	58
Started	0	55	88	108	112

STAMPEDE OVERSIGHT COMMITTEES, STAFF AND COLLABORATORS

See separate **supplementary document** for pdf list of STAMPEDE oversight committees, staff and collaborators.

Any subsequent updates may also be found through:

<http://www.stampede-trial.org/media-section/presentation-repository/trial-recognition/>

PROTOCOL

Pertinent versions of the protocol are provided as separate **supplementary documents**.

A full list of previous protocol versions can be found via the trial website:

<http://www.stampededtrial.org/>