

VIJAYA DIAGNOSTIC CENTRE®

Opp.Metro Pillar No. A1516, Indira Nagar, Dilsukhnagar, Hyderabad - 500089

TEST REPORT

Name : Mrs. SWETHA M Age/Gender : 38 Years / Female

Registration ID : 220600002580

Ref. By : Self
Sample Type : Serum

Registered on: 04-Dec-2022 09:24 Collected on: 04-Dec-2022 09:29

Released on : 04-Dec-2022 14:30

Printed on : 11-May-2024 18:11

Regn Centre : Bandlaguda-60

3rd Trimester : 0.8 - 5.2

THYROID PROFILE

TEST NAME RESULT UNIT BIOLOGICAL REFERENCE INTERVAL T3 Total :1.11 Non pregnant : 0.60 - 1.81 ng/mL Pregnant Method: Chemiluminiscence Immunoassay 1st Trimester : 0.81 - 1.90 2nd & 3rd Trimester: 1.0 - 2.60 **T4 Total** : 8.60 : 3.2 - 12.6 μg/dL Pregnant : 6.4 - 10.7 Method: Chemiluminiscence Immunoassay **TSH** - Ultrasensitive : 8.542 Non pregnant : 0.55 - 4.78 μIU/mL Pregnant Method: Chemiluminiscence Immunoassay 1st Trimester : 0.3 - 4.5 2nd Trimester : 0.5 - 4.6

Interpretation / Comments:

....

- Patient preparation is particularly important for hormone studies, results of which may be markedly
 affectedby many factors such as stress, position, fasting state, time of the day, preceding diet and drug
 therapy.
- The levels of T3 helps in the diagnosis of T3 thyrotoxicosis and monitoring the course of hyperthyroidism.T3 is not recommended for diagnosis of hypothyroidism as decreased values have minimal clinical significance.
- Values below the lower limits can be caused by a number of conditions including non-thyroidal illness, acute and chronic stress and hypothyroidism.
- Elevated level of T4 are seen in hyperthyroidism, pregnancy, euthyroid patients with increased serum thyroxine binding globulin.
- Decreased levels are noted in hypothyroidism, hypoproteinemia, euthyroid sick syndrome, decrease in thyroxine binding globulin.
- TSH levels are increased in primary hypothyroidism, insufficient thyroid hormone replacement therapy, Hashimotos thyroiditis, use of amphetamines, dopamine antagonists, iodine containing agents, lithium and iodine induced or deficiency goiter.
- Decreased levels of TSH may be seen in graves disease, toxic multinodular goitre, thyroiditis, excessivetreatment with thyroid hormone replacement and central hypothyroidism.





Anitha

DR. R ANITHA.
MD,BIOCHEMISTRY
CONSULTANT BIOCHEMISTRY

Registration No: 70111

----- End of Report -----