

VIJAYA DIAGNOSTIC CENTRE®

Opp.Metro Pillar No. A1516, Indira Nagar, Dilsukhnagar, Hyderabad - 500089

TEST REPORT

Name : Mrs. SWETHA M
Age/Gender : 40 Years / Female
Registration ID : 240600008455

Ref. By : Self Sample Type : Serum Registered on: 25-Apr-2024 07:47 Collected on: 25-Apr-2024 07:52 Released on: 25-Apr-2024 16:43

Printed on : 08-May-2024 12:51

Regn Centre : Bandlaguda-60

THYROID PROFILE

<u>TEST NAME</u> <u>RESULT</u> <u>UNIT</u> <u>BIOLOGICAL REFERENCE INTERVAL</u>

 $\textbf{T3 Total} \hspace{1.5cm} : 1.04 \hspace{1.5cm} \text{ng/mL} \hspace{1.5cm} \text{Non pregnant} \hspace{1.5cm} : 0.60 - 1.81$

Method: Chemiluminiscence Immunoassay

Pregnant

1st Trimester : 0.81 - 1.90 2nd & 3rd Trimester : 1.0 - 2.60

 $\textbf{T4 Total} \hspace{1.5cm} : 8.70 \hspace{1.5cm} \mu g/dL \hspace{1.5cm} Adult \hspace{1.5cm} : 3.2 - 12.6$

Pregnant : 6.4 - 10.7

TSH - Ultrasensitive : 10.658 μIU/mL Non pregnant : 0.55 - 4.78

Pregnant

1st Trimester : 0.3 - 4.5 2nd Trimester : 0.5 - 4.6 3rd Trimester : 0.8 - 5.2

Interpretation / Comments:

Method: Chemiluminiscence Immunoassay

Method: Chemiluminiscence Immunoassay

• Patient preparation is particularly important for hormone studies, results of which may be markedly affected by many factors such as stress, position, fasting state, time of the day, preceding diet and drug therapy.

- The levels of T3 helps in the diagnosis of T3 thyrotoxicosis and monitoring the course of hyperthyroidism.T3 is not recommended for diagnosis of hypothyroidism as decreased values have minimal clinical significance.
- Values below the lower limits can be caused by a number of conditions including non-thyroidal illness, acute and chronic stress and hypothyroidism.
- Elevated level of T4 are seen in hyperthyroidism, pregnancy, euthyroid patients with increased serum thyroxine binding globulin.
- Decreased levels are noted in hypothyroidism, hypoproteinemia, euthyroid sick syndrome, decrease in thyroxine binding globulin.
- TSH levels are increased in primary hypothyroidism, insufficient thyroid hormone replacement therapy, Hashimotos thyroiditis, use of amphetamines, dopamine antagonists, iodine containing agents, lithium and iodine induced or deficiency goiter.
- Decreased levels of TSH may be seen in graves disease, toxic multinodular goitre, thyroiditis, excessivetreatment with thyroid hormone replacement and central hypothyroidism.





Vohahan

DR. SYED SHABBAR MASIH MD BIOCHEMISTRY

Registration No: TSMC/FMR/44589

----- End of Report -----

^{*} Suggested Clinical Correlation, If Necessary Kindly Discuss with signatory