- Create a table in your database with the following columns: `sale_date`, `product_id`, and `total_revenue`.

```
CREATE TABLE sales(
sale_date DATE,
 product_id INT,
total_revenue DECIMAL(10,2)
);
```

- Insert at least 10 rows of sample data into the table.

```
INSERT INTO sales VALUES
('2026-04-01', 101, 500.00),
('2026-04-01', 102, 750.00),
('2026-04-02', 101, 300.00),
('2026-04-02', 103, 900.00),
('2026-04-03', 102, 650.00),
('2026-04-03', 104, 1200.00),
('2026-04-04', 101, 400.00),
('2026-04-04', 103, 800.00),
('2026-04-05', 104, 1500.00),
('2026-04-05', 102, 700.00);
```

- Write SQL queries using aggregate functions (e.g., `SUM`, `AVG`, `MAX`, `MIN`, `COUNT`) to answer the following questions:
  - What is the total revenue for all sales?

```
select SUM(total_revenue) AS total_revenue_all_sales from sales;
```

  - What is the average revenue per sale?

```
select AVG(total_revenue) AS average_revenue from sales;
```

  - What is the maximum and minimum revenue for a single sale?

```
Select max(total_revenue) AS max_revenue,
```

```
min(total_revenue) AS min_revenue from sales;
```

- How many sales were made for each product ID?

```
select product_id ,count(*) AS total_sales
from sales
group by product_id ;
```

- Save your SQL queries and results in a file named `aggregate_functions_practice.sql`.

Results:

- **Total Revenue** → 7700
- **Average Revenue** → 770
- **Max Revenue** → 1500
- **Min Revenue** → 300