

## PH423 Assignment 1

Parth Sastry, Sahas Kamat, Sankalp Gambhir (180260026, 30, 32)

August 25, 2020

### Question 1.

---

[Sankalp: I got this one.] We consider the action of  $[\hat{\mathbf{X}}, \hat{\mathbf{P}}]$  on an arbitrary state  $\phi$ ,

$$[\hat{\mathbf{X}}, \hat{\mathbf{P}}] \phi(x) . \quad (1)$$

Starting with equation 1, we first write it in bra-ket notation as

$$[\hat{\mathbf{X}}, \hat{\mathbf{P}}] \phi(x) = \langle x | [\hat{\mathbf{X}}, \hat{\mathbf{P}}] | \phi \rangle . \quad (2)$$

Since  $[\hat{\mathbf{X}}, \hat{\mathbf{P}}] = i\hbar$ , we get

$$\langle x | [\hat{\mathbf{X}}, \hat{\mathbf{P}}] | \phi \rangle = \langle x | i\hbar | \phi \rangle . \quad (3)$$

On either side, we can introduce the identity operator as  $[\hat{\mathbf{X}}, \hat{\mathbf{P}}] \cdot \hat{\mathbf{1}}$  and  $i\hbar \cdot \hat{\mathbf{1}}$ , and use the position space completeness relation  $\int dx' |x'\rangle\langle x'| = \hat{\mathbf{1}}$  to obtain

$$\int dx' \langle x | [\hat{\mathbf{X}}, \hat{\mathbf{P}}] | x'\rangle\langle x'| | \phi \rangle = \int dx' \langle x | i\hbar | x'\rangle\langle x'| | \phi \rangle . \quad (4)$$

Expanding the commutator operation  $\langle x | [\hat{\mathbf{X}}, \hat{\mathbf{P}}] | x'\rangle$  as

$$\langle x | [\hat{\mathbf{X}}, \hat{\mathbf{P}}] | x'\rangle = \langle x | \hat{\mathbf{X}} \hat{\mathbf{P}} | x'\rangle - \langle x | \hat{\mathbf{P}} \hat{\mathbf{X}} | x'\rangle$$

and applying the action of  $\hat{\mathbf{X}}$  on  $|x'\rangle$  and  $\langle x|$ , we obtain

$$\int dx' (x - x') \langle x | \hat{\mathbf{P}} | x'\rangle\langle x'| | \phi \rangle = \int dx' \langle x | i\hbar | x'\rangle\langle x'| | \phi \rangle . \quad (5)$$

Since the state  $\phi(x)$  chosen was arbitrary, we get the weak equivalence

$$(x - x') \langle x | \hat{\mathbf{P}} | x'\rangle = i\hbar \langle x | x'\rangle . \quad (6)$$

We know  $\langle x | x'\rangle = \delta(x - x')$ , and using

$$\delta(x) = -x\delta'(x)$$

we get

$$(x - x') \langle x | \hat{P} | x' \rangle = -i\hbar(x - x') \frac{d}{dx} \delta(x - x') . \quad (7)$$

And finally, dividing both sides by  $(x - x')$ , assuming  $x \neq x'$ , we get the required matrix element

$$\langle x | \hat{P} | x' \rangle = -i\hbar \frac{d}{dx} \delta(x - x') . \quad (8)$$

□

## Question 2.

[Parth: Question 2 is mine] Starting from the abstract ket form of the Schrodinger eqn. derive the Schrodinger eqn. in the co-ordinate basis

The Schrodinger eqn. in the abstract ket form is given as follows -

$$\hat{H} |\psi\rangle = E |\psi\rangle \quad (9)$$

In the position basis,  $\hat{H}$  can be expressed in the form of a matrix, while  $|\psi\rangle$  can be expressed in terms of components in the basis. In the position basis, the components of  $|\psi\rangle$  form the wavefunction.

$$\langle x | \psi \rangle = \psi(x). \quad (10)$$

The Hamiltonian becomes a matrix in the position basis, with components given by -

$$H_{xy} = \langle x | \hat{H} | y \rangle . \quad (11)$$

Applying this to equation 9. We multiply by  $\langle x |$  on the left side and insert an identity on the right hand side. The identity is given by -

$$1 = \int dy |y\rangle \langle y| \quad (12)$$

So now we obtain -

$$\int dy \langle x | \hat{H} | y \rangle \langle y | \psi \rangle = E \langle x | \psi \rangle . \quad (13)$$

simplifying to components, we get -

$$\int dy H_{xy} \psi(y) = E \psi(x). \quad (14)$$

now our components are -

$$H_{xy} = -\frac{\hbar^2}{2m} \delta''(x - y) + V \delta(x - y) \quad (15)$$

we plug this in to equation 14. We get -

$$\int dy \left[ -\frac{\hbar^2}{2m} \delta''(x - y) \psi(y) \right] + \int dy V \psi(y) \delta(x - y) = E \psi(x) \quad (16)$$

$$-\frac{\hbar^2}{2m} \int dy \delta''(x-y) \psi(y) + V \psi(x) = E \psi(x) \quad (17)$$

Using Integration by Parts on the integral twice, we obtain -

$$-\frac{\hbar^2}{2m} \int dy \delta(x-y) \psi''(y) + V \psi(x) = E \psi(x) \quad (18)$$

$$-\frac{\hbar^2}{2m} \psi''(x) + V \psi(x) = E \psi(x) \quad (19)$$

Generalising this a bit -

$$-\frac{\hbar^2}{2m} \nabla^2 \psi + V \psi = E \psi \quad (20)$$

This is quite obviously, our Schrodinger's Equation in co-ordinate form.

### Question 3.

---

[Sahas: This one's mine.]