

# Using Digital Biomarkers for Objective Assessment of Perfusionists' Workload and Acute Stress during Cardiac Surgery

Roger D. Dias<sup>1,2</sup>[0000-0003-4959-5052], Lauren R. Kennedy-Metz<sup>3</sup>[0000-0002-2696-3943], Rithy Srey<sup>4</sup>, Geoffrey Rance<sup>5</sup>[0000-0002-4270-1142], Mahdi Ebnali<sup>1,2</sup>, David Arney<sup>1,6</sup>[0000-0002-5536-777X], Matthew Gombolay<sup>7</sup>[0000-0002-5321-6038] and Marco A. Zenati<sup>1,8</sup>[0000-0001-7139-0323]

<sup>1</sup> Harvard Medical School, Boston, MA

<sup>2</sup> Department of Emergency Medicine, Mass General Brigham, Boston, MA

<sup>3</sup> Department of Psychology, Roanoke College, Salem, VA

<sup>4</sup> Division of Cardiac Surgery, Veterans Affairs Boston Healthcare System, Boston, MA

<sup>5</sup> Department of Cardiac Surgery, Cape Cod Healthcare, Hyannis, MA

<sup>6</sup> Department of Anesthesia, Critical Care and Pain Medicine, Massachusetts General Hospital, Boston, MA

<sup>7</sup> Georgia Institute of Technology, Atlanta, GA

<sup>8</sup> Division of Cardiac Surgery, Veterans Affairs Boston Healthcare System and Medical Robotics and Computer Assisted Surgery Lab, Boston, MA  
rdias@bwh.harvard.edu

**Abstract.** The cardiac operating room (OR) is a high-risk, high-stakes environment inserted into a complex socio-technical healthcare system. During cardiopulmonary bypass (CPB), the most critical phase of cardiac surgery, the perfusionist has a crucial role within the interprofessional OR team, being responsible for optimizing patient perfusion while coordinating other tasks with the surgeon, anesthesiologist, and nurses. The aim of this study was to investigate objective digital biomarkers of perfusionists' workload and stress derived from heart rate variability (HRV) metrics captured via a wearable physiological sensor in a real cardiac OR. We explored the relationships between several HRV parameters and validated self-report measures of surgical task workload (SURG-TLX) and acute stress (STAI-SF), as well as surgical processes and outcome measures. We found that the frequency-domain HRV parameter HF relative power - FFT (%) presented the strongest association with task workload (correlation coefficient:-0.491, p-value: 0.003). We also found that the time-domain HRV parameter RMSSD (ms) presented the strongest correlation with perfusionists' acute stress (correlation coefficient:-0.489, p-value: 0.005). A few workload and stress biomarkers were also associated with bypass time and patient length of stay in the hospital. The findings from this study will inform future research regarding which HRV-based biomarkers are best suited for the development of cognitive support systems capable of monitoring surgical workload and stress in real time.

**Keywords:** Cognitive Workload, Acute Stress, Perfusionists, Cardiac Surgery, Digital Biomarkers.

## 1 Introduction

The cardiac operating room (OR) is a high-risk, high-stakes environment inserted into a complex socio-technical healthcare system [1,2]. Despite the tremendous improvements in surgical patients' mortality and morbidity achieved in the past two decades, patient safety and preventable adverse event rates are still suboptimal among cardiac surgery patients [3–5]. During the cardiopulmonary bypass (CPB), the most critical phase of cardiac surgery, the perfusionist has a crucial role within the interprofessional OR team, being responsible for optimizing patient perfusion while coordinating other tasks with the surgeon, anesthesiologist, and nurses [6]. The inherently complex nature of the OR setting, alongside the requirements for effective and sustained situational awareness, decision-making, and teamwork, make perfusionists particularly vulnerable to the deleterious effects that high cognitive workload and stress may have on clinical performance [7,8].

Extensive previous literature has shown that cognitive overload and high acute stress (distress) impair surgical performance in both simulated and real-life settings [9,10]. Nonetheless, most of the existing evidence is surgeon-centered, and only a few studies have included other OR team members, particularly the perfusionists. Furthermore, the majority of this previous research has used self-report instruments (e.g., questionnaires) that provide a validated overall measure of the entire surgical procedure but do not support objective monitoring of clinicians' workload and stress in real-time [9,11].

Among several real-time objective measures, heart rate variability (HRV) is a physiological biomarker that can be unobtrusively captured via digital wearable sensors and has been used as a proxy for cognitive workload and acute stress in various settings, including the OR [9,12,13]. Through HRV analysis, many parameters are generated in the time, frequency, and non-linear domains, with each of these parameters reflecting autonomic nervous system activity (sympathetic and/or parasympathetic) at varying levels [14]. HRV has been shown to index high-level cognitive functions (e.g., cognitive workload) and emotional regulation (e.g., stress) [15].

The aim of this study was to investigate objective digital biomarkers of perfusionists' workload and stress derived from HRV metrics captured via a wearable physiological sensor in a real cardiac OR. We explored the relationships between several HRV parameters and validated self-report measures of surgical task workload and acute stress, as well as surgical processes and outcomes measures.

## 2 Methods

### 2.1 Study Setting and Design

This was an observational study conducted in the cardiac OR of a tertiary hospital in the United States between January 2021 and May 2022. The research protocol was approved by the Institutional Review Boards (IRB) at the Veterans Affairs (VA) Bos-

ton Healthcare System and Harvard Medical School. Informed consent was obtained from all participants, including patients, perfusionists, and OR staff.

## 2.2 Population

Participants were perfusionists who worked in the cardiac OR and patients who were undergoing open cardiac surgery procedures: coronary artery bypass grafting (CABG) and/or aortic valve replacement (AVR). No exclusion criteria were applied.

## 2.3 Procedures

At the beginning of each operation, perfusionists were attached to a wireless 3-lead electrocardiogram (ECG) device (*MindWare Mobile Impedance Cardiograph*) that records ECG signals at 500Hz. After the operation, the ECG files were exported and analyzed in the *Kubios HRV* software [16]. In *Kubios*, an automatic artifact correction was selected.

At the end of the procedure, perfusionists completed two previously validated self-report questionnaires: the Surgery Task Load Index (SURG-TLX)[17], assessing their perceived surgical task workload, and the Spielberger Short-Form State-Trait Anxiety Inventory (STAI-SF)[17,18], assessing their perceived acute stress.

## 2.4 Measurements

Heart Rate Variability (HRV). Perfusionists' ECG signals were processed in *Kubios* [16] to extract a series of 44 different HRV parameters for each 5-minute time window, calculated based on established HRV analysis guidelines [16,19]. The analysis included HRV metrics from time, frequency, and non-linear domains. To gather a measure that reflects the overall surgical procedure, the 5-minute windows were averaged across the entire procedure for each HRV parameter.

**Perceived Surgical Task Workload.** To measure the perceived (self-reported) workload related to the entire surgical procedure, the SURG-TLX was completed by perfusionists [17,20]. This is a 20-point scale for which the score is calculated by multiplying the number of points selected by 5 (score range: 5-100). The tool covers six different workload domains: mental demands, physical demands, temporal demands, task complexity, distractions, and situational stress. Each domain is rated individually, then averaged to generate the SURG-TLX total score.

**Perceived Acute Stress.** To measure the perceived (self-reported) acute stress related to the entire surgical procedure, the STAI-SF was completed by perfusionists [18]. This is a 4-point scale for which half of the items have a reverse score. The tool has six different items, asking participants to report how much they feel: calm, tense, upset, relaxed, content, and worried. Each item is rated individually, then the sum of the scores (reversed for items 1, 3, and 5) generates the STAI-SF total score, ranging from 6 to 24 points.

**Surgical processes and outcome measures.** Patient data was collected from the electronic health record. Surgical processes measures included: total procedure duration,

cross-clamp time, and bypass time, all in minutes. Surgical outcome measures included: intrahospital mortality (%), and length of hospital stay after surgery (in days).

### 3 Results

A total of 37 cardiac surgery patients and 4 perfusionists were included in this study. Twenty-nine (78.4%) patients underwent an isolated CABG procedure, 5 (13.5%) patients received an isolated AVR procedure, and 3 (8.1%) patients had a combined CABG/AVR procedure. Table 1 displays the patient's demographic and surgical characteristics.

**Table 1.** Patient Demographic and Surgical Characteristics.

<b>Variables</b>	<b>Measures</b>
Sex male - <i>N</i> (%)	37 (100.0)
Age - <i>years</i>	71.0 (63.0 - 74.0)
BMI - <i>Kg/m<sup>2</sup></i>	29.7 (27.3 - 32.5)
BSA - <i>m<sup>2</sup></i>	2.1 (1.9 - 2.2)
<b><i>Pre-Operative Risk (VASQIP risk assessment core)</i></b>	
30-day Mortality Risk - %	0.8 (0.4 - 1.3)
30-day Morbidity Risk - %	6.9 (6.7 - 9.6)
30-day SSI Risk - %	1.4 (0.9 - 2.0)
<b><i>Surgical Process and Outcome Measures</i></b>	
Bypass Duration - <i>minutes</i>	112.0 (94.0 - 134.0)
Cross Clamp Time - <i>minutes</i>	71.0 (56.0 - 90.0)
Total Procedure Duration - <i>minutes</i>	330.0 (258.0 - 371.0)
Hospital Length of Stay - <i>days</i>	7.0 (6.0 - 9.0)
Mortality - <i>N</i> (%)	0 (0.0)

*\*BMI: body mass index; BSA: body surface area; VASQIP: Veteran Affairs Surgical Quality Improvement Program; SSI: surgical site infection*

A correlation analysis was performed between all objective (HRV parameters) and subjective (self-report instruments) measures of task workload and acute stress. Table 2 (Task Workload) and Table 3 (Acute Stress) show the correlation coefficients and corresponding p-values for all statistically significant associations.

**Table 2.** Significant Correlations between HRV parameters and SURG-TLX total score.

HRV Parameter	HRV Domain	Correlation Coefficient	p-value
HF relative power - FFT (%)	Frequency-domain	-0.491	0.003
HF absolute power - FFT (n.u.)	Frequency-domain	-0.485	0.004
LF absolute power - FFT (n.u.)	Frequency-domain	0.485	0.004
LF relative power - FFT (%)	Frequency-domain	0.476	0.004
LF/HF ratio - FFT	Frequency-domain	0.472	0.005
VLF peak frequency - FFT (Hz)	Frequency-domain	0.454	0.007
DFA alpha1 – short-term fluctuation slope	Non-linear	0.382	0.026

*\*HF: high-frequency band; LF: low-frequency band; VLF: very-low-frequency band; DFA: detrended fluctuation analysis; FFT: Fast Fourier Transformation*


**Table 3.** Significant Correlations between HRV parameters and STAI-SF total score.

HRV Parameter	HRV Domain	Correlation Coefficient	p-value
RMSSD (ms)	Time-Domain	-0.489	0.005
SD1 in Poincaré plot (ms)	Non-linear	-0.488	0.005
HF absolute power - FFT (ms2)	Frequency-Domain	-0.485	0.006
LF peak frequency - FFT (Hz)	Frequency-Domain	-0.469	0.008
HF absolute power - FFT (log)	Frequency-Domain	-0.465	0.008
PNS index	<i>Kubios</i> Metric [16]	-0.461	0.009
RPA Lmax - maximum line length (beats)	Non-linear	0.459	0.009
VLF absolute power - FFT (ms2)	Frequency-Domain	-0.459	0.009
pNN50 (%)	Time-Domain	-0.454	0.010
TINN (ms)	Time-Domain	-0.453	0.011
SD2/SD1 ratio	Non-linear	0.436	0.014
SDNN (ms)	Time-Domain	-0.436	0.014
Total power - FFT (ms2)	Frequency-Domain	-0.434	0.015
NN50 (beats)	Time-Domain	-0.431	0.015
DFA alpha1 - short-term fluctuation slope	Non-linear	0.425	0.017
HF relative power - FFT (%)	Frequency-Domain	-0.421	0.018
HF absolute power - FFT (n.u.)	Frequency-Domain	-0.418	0.019
LF absolute power - FFT (n.u.)	Frequency-Domain	0.418	0.019
Stress index	<i>Kubios</i> Metric [16]	0.409	0.022
LF relative power - FFT (ms2)	Frequency-Domain	-0.407	0.023
RPA Shannon Entropy	Non-linear	0.401	0.026
D2 - correlation dimension	Non-linear	-0.393	0.029


LF/HF ratio - FFT	Frequency-Domain	0.391	0.030
RPA Determinism (%)	Non-linear	0.379	0.036
RPA Recurrence Rate (%)	Non-linear	0.377	0.037
HRV Triangular Index	Time-Domain	-0.369	0.041
VLF absolute power - FFT (log)	Frequency-Domain	-0.369	0.041
ApEn - Approximate Entropy	Non-linear	-0.358	0.048
SampEn - Sample Entropy	Non-linear	-0.355	0.050

*\*RMSSD: root mean square of the successive differences between RR intervals; SD1: standard deviation perpendicular to the line-of-identity; SD2: standard deviation along the line-of-identity; HF: high-frequency band; LF: low-frequency band; VLF: very-low-frequency band; PNS: parasympathetic nervous system activity; RPA: recurrent plot analysis; NN50: Number of successive RR interval pairs that differ more than 50 ms; pNN50: NNxx divided by the total number of RR intervals; DFA: detrended fluctuation analysis; FFT: Fast Fourier Transformation; TINN: Baseline width of the RR interval histogram; SDNN: Standard deviation of RR intervals; Stress index: Square root of Baevsky's [21] stress index.*

The scatter plots in Fig. 1 and Fig. 2 display the relationship between the HRV parameters presenting the strongest correlations with the perfusionist's workload and acute stress. Both parameters - the high-frequency (HF) band and the root mean square of the successive differences between heartbeat intervals (RMSSD) - reflect parasympathetic autonomic nervous activity (vagal modulation). However, HF is influenced by respiratory rate while RMSSD is not [22]. The inverse relationship reflects the fact that high workload and stress situations lead to a decrease in parasympathetic activity while increasing sympathetic tone.



**Fig. 1.** Relationship between a parasympathetic-mediated HRV parameter (frequency-domain) and perfusionists' workload (SURG-TLX).



**Fig. 2.** Relationship between a parasympathetic-mediated HRV parameter (time-domain) and perfusionists' acute stress (STAI-SF).

The only HRV parameter associated with patient pre-operative risk scores was the very-low-frequency (VLF) relative power, which presented a correlation coefficient of -0.346, and a p-value of 0.045 with the risk for surgical site infection (SSI). No other relationships with pre-operative risk scores yielded a statistically significant correlation.

Additional correlational analysis between perfusionists' workload and acute stress metrics and surgical processes and outcome measures revealed that the bypass duration was associated with the following metrics: low-frequency (LF) absolute power (log) (correlation coefficient: 0.366, p-value: 0.033); Standard deviation along the line-of-identity (SD2) (correlation coefficient: -0.370, p-value: 0.031); Stress Index (correlation coefficient: -0.346, p-value: 0.045); and VLF absolute power (log) (correlation coefficient: -0.412, p-value: 0.016). Only the HRV parameter Mean RR-intervals (ms), which represent the mean interval between consecutive heartbeats in milliseconds, presented a statistically significant correlation with patient length of stay in the hospital after surgery (correlation coefficient: -0.346, p-value: 0.0045). No other relationships with surgical processes and outcome measures yielded a statistically significant correlation.

## 4 Discussion

In the present study, we collected data from patients and perfusionists in a real cardiac OR, leveraging an integrative approach based on human factors and cognitive engineering sciences [23]. To investigate objective digital biomarkers of perfusionists' workload and acute stress, we performed HRV analysis from ECG signals and ex-

plored the relationships between several HRV parameters and previously validated self-report tools. The findings from this study will inform future research regarding which HRV-based biomarkers are best suited for the development of cognitive support systems capable of monitoring surgical workload and stress in real time.

Several previous studies have assessed the cognitive workload of surgeons and, less often, other OR team members during cardiac surgery [9]. Most of these studies have used self-report instruments such as the NASA Task Load Index (NASA-TLX) [24] and, more recently, the SURG-TLX, which was validated specifically in the surgical setting [9,11]. Acute stress has also been investigated in many surgical studies, and the Spielberg STAI questionnaire is one of the most used instruments to assess acute stress in various settings [25]. Among HRV metrics, the vast majority of the studies have selected only a few parameters to investigate based on previous research [26,27]. To date, no surgical study has investigated and compared the strength of correlation among all the HRV parameters extracted in the time, frequency, and non-linear domains. Furthermore, most previous studies in the field have suggested the LF/HF ratio is the best marker of cognitive workload [9]. Nonetheless, our findings show that there are other HRV parameters with equal or even stronger correlation with SURG-TLX total score. Interestingly, our results showed that no time-domain parameter was associated with perfusionists' workload. Regarding acute stress, the strongest correlation was found with RMSSD, which corroborates previous literature that has suggested this time-domain parameter as a good indicator of acute stress [9,12]. It is also important to notice that many non-linear parameters were significantly associated with perfusionists' workload and stress, and studies exploring these parameters to infer cognitive and emotional functions in the OR are scarce. Future research should include non-linear HRV parameters in their analysis.

Surprisingly, the patient pre-operative risk, which is an indicator of expected task complexity and difficulty [28], was only associated with a few self-report items and HRV parameters. A possible explanation is that the population of patients included in this study did not substantially vary in their pre-operative risk (Table 1), therefore, not allowing the discovery of linear correlations with workload and stress. Another explanation is that high-risk patients may pose an additional workload to the surgeon, particularly in the psychomotor domain, but not necessarily to the perfusionist. The bypass duration, the time in which a heart-lung machine provides the patient perfusion, is a reliable surgical process measure that serves as a surrogate for various post-operative outcomes [29]. Several self-report and HRV parameters were associated with bypass duration, showing that a longer bypass phase is associated with a higher perfusionist's workload and stress. Length of hospital stay, an important patient outcome measure, was also moderately associated with an HRV parameter (mean R-R interval).

The findings from this study are promising, particularly as it relates to validating objective digital biomarkers of perfusionists' workload and stress that can be monitored in near or real-time during cardiac surgery. Although we have averaged the HRV parameters over the entire procedure, these metrics are extracted for each 5-minute window or even shorter, such as 1-minute, using ultra-short-term HRV analysis [30]. In the same way we currently monitor patient physiology in real-time, in-



forming clinical decision-making in the OR, the continuous monitoring of perfusionists' cognitive and emotional states can be used by intelligent ambient systems capable of detecting cognitive overload situations and re-allocating tasks across the OR team, distributing cognitive demands [31]. Such intelligent systems [32] can also manage unavoidable interruptions to the surgical workflow, providing the surgical team with a data-driven assessment of clinicians' workload and stress that enable timely interruptions at specific moments when the workload is not highly critical [33].

Although this study reveals important relationships between perfusionists' workload, stress, and surgical processes and outcome measures, it is important to highlight that these findings cannot infer a cause-effect relationship. In fact, in the same way that task difficulty and complexity can increase cognitive load and stress, these two states can also deteriorate performance, which per se, can increase stress [34]. These relationships between task workload, acute stress, and performance are intrinsically complex and should be understood and investigated through a feedback loop perspective. Future research should attempt to model these relationships using non-linear models and network analysis in order to capture further the complexity and nuances of such a dynamic system.

## 5 Conclusion

In conclusion, the findings from this study provide a comprehensive exploratory analysis of the several HRV parameters that have been reported in the literature to index cognitive and emotional states across various industries in which human performance plays a critical role in safety. By investigating the comparative association between these different HRV parameters and established measures of cognitive workload and acute stress in the cardiac OR, our results advance the current scientific knowledge toward the development and validation of objective digital biomarkers of human performance in the surgical setting. More importantly, because these metrics are unobtrusively captured in real-time, our findings provide the foundational evidence for future cognitive augmentation capabilities [35] that can augment and improve performance in the OR, reducing medical errors and enhancing patient safety.

## 6 Acknowledgment

This work was supported by the National Heart, Lung, and Blood Institute (NHLBI) of the National Institutes of Health (NIH) [Grant Numbers: R01HL126896, R01HL157457]. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

## References

1. Frank P, Nurok M, Sundt T. Human Factors Considerations in Cardiac Surgery. *Human Factors in Surgery*. 2020. pp. 131–140. doi:10.1007/978-3-030-53127-0\_13

2. Cohen TN, Ley EJ, Gewertz BL. *Human Factors in Surgery: Enhancing Safety and Flow in Patient Care*. Springer Nature; 2020. Available: [https://play.google.com/store/books/details?id=cuv\\_DwAAQBAJ](https://play.google.com/store/books/details?id=cuv_DwAAQBAJ)
3. Christian CK, Gustafson ML, Roth EM, Sheridan TB, Gandhi TK, Dwyer K, et al. A prospective study of patient safety in the operating room. *Surgery*. 2006;139: 159–173. doi:10.1016/j.surg.2005.07.037
4. Martinez EA, Shore A, Colantuoni E, Herzer K, Thompson DA, Gurses AP, et al. Cardiac surgery errors: results from the UK National Reporting and Learning System. *Int J Qual Health Care*. 2011;23: 151–158. doi:10.1093/intqhc/mzq084
5. Gurses AP, Kim G, Martinez EA, Marsteller J, Bauer L, Lubomski LH, et al. Identifying and categorising patient safety hazards in cardiovascular operating rooms using an interdisciplinary approach: a multisite study. *BMJ Qual Saf*. 2012;21: 810–818. doi:10.1136/bmjqs-2011-000625
6. Wiegmann D, Suther T, Neal J, Parker SH, Sundt TM. A human factors analysis of cardiopulmonary bypass machines. *J Extra Corpor Technol*. 2009;41: 57–63. Available: <https://www.ncbi.nlm.nih.gov/pubmed/19681301>
7. Flin R, Youngson GG, Yule S. *Enhancing Surgical Performance: A Primer in Non-technical Skills*. CRC Press; 2015. Available: <https://play.google.com/store/books/details?id=cn-JDwAAQBAJ>
8. Kennedy-Metz LR, Dias RD, Srey R, Rance GC, Conboy HM, Haime ME, et al. Analysis of Dynamic Changes in Cognitive Workload During Cardiac Surgery Perfusionists' Interactions With the Cardiopulmonary Bypass Pump. *Hum Factors*. 2021;63: 757–771. doi:10.1177/0018720820976297
9. Dias RD, Ngo-Howard MC, Boskovski MT, Zenati MA, Yule SJ. Systematic review of measurement tools to assess surgeons' intraoperative cognitive workload. *Br J Surg*. 2018;105: 491–501. doi:10.1002/bjs.10795
10. Arora S, Sevdalis N, Nestel D, Woloshynowych M, Darzi A, Kneebone R. The impact of stress on surgical performance: a systematic review of the literature. *Surgery*. 2010;147: 318–30, 330.e1–6. doi:10.1016/j.surg.2009.10.007
11. Kennedy-Metz LR, Wolfe HL, Dias RD, Yule SJ, Zenati MA. Surgery Task Load Index in Cardiac Surgery: Measuring Cognitive Load Among Teams. *Surg Innov*. 2020;27: 602–607. doi:10.1177/1553350620934931
12. Peabody JE, Ryznar R, Ziesmann MT, Gillman L. A Systematic Review of Heart Rate Variability as a Measure of Stress in Medical Professionals. *Cureus*. 2023;15: e34345. doi:10.7759/cureus.34345
13. Kennedy-Metz LR, Dias RD, Srey R, Rance GC, Furlanello C, Zenati MA. Sensors for Continuous Monitoring of Surgeon's Cognitive Workload in the Cardiac Operating Room. *Sensors*. 2020;20. doi:10.3390/s20226616
14. Zammuto M, Ottaviani C, Laghi F, Lonigro A. The Heart in the Mind: A Systematic Review and Meta-Analysis of the Association Between Theory of Mind and Cardiac Vagal Tone. *Front Physiol*. 2021;12: 611609. doi:10.3389/fphys.2021.611609
15. Forte G, Favieri F, Casagrande M. Heart Rate Variability and Cognitive Function: A Systematic Review. *Front Neurosci*. 2019;13: 710. doi:10.3389/fnins.2019.00710
16. Tarvainen MP, Niskanen J-P, Lipponen JA, Ranta-aho PO, Karjalainen PA. Kubios HRV – Heart rate variability analysis software. *Computer Methods and Programs in Biomedicine*. 2014. pp. 210–220. doi:10.1016/j.cmpb.2013.07.024
17. Wilson MR, Poolton JM, Malhotra N, Ngo K, Bright E, Masters RSW. Development and validation of a surgical workload measure: the surgery task load index (SURG-TLX). *World J Surg*. 2011;35: 1961–1969. doi:10.1007/s00268-011-1141-4

18. Marteau TM, Bekker H. The development of a six-item short-form of the state scale of the Spielberger State-Trait Anxiety Inventory (STAI). *Br J Clin Psychol.* 1992;31: 301–306. doi:10.1111/j.2044-8260.1992.tb00997.x
19. Heart rate variability. Standards of measurement, physiological interpretation, and clinical use. Task Force of the European Society of Cardiology and the North American Society of Pacing and Electrophysiology. *Eur Heart J.* 1996;17: 354–381. Available: <https://www.ncbi.nlm.nih.gov/pubmed/8737210>
20. Kennedy-Metz L, Dias R, Srey R, Rance G, Leissner K, Zorca S, et al. Human factors analysis of Goal-directed perfusion in cardiac surgery. *Healthcare and Medical Devices. AHFE International*; 2022. doi:10.54941/ahfe1002120
21. Baevsky RM, Institute of Biomedical Problems of the Russian Academy of Sciences, Chernikova AG. Heart rate variability analysis: physiological foundations and main methods. *Cardiometry.* 2017; 66–76. doi:10.12710/cardiometry.2017.10.6676
22. Shaffer F, Ginsberg JP. An Overview of Heart Rate Variability Metrics and Norms. *Front Public Health.* 2017;5: 258. doi:10.3389/fpubh.2017.00258
23. Zenati MA, Kennedy-Metz L, Dias RD. Cognitive Engineering to Improve Patient Safety and Outcomes in Cardiothoracic Surgery. *Semin Thorac Cardiovasc Surg.* 2020;32: 1–7. doi:10.1053/j.semtcvs.2019.10.011
24. Wadhwa RK, Parker SH, Burkhart HM, Greason KL, Neal JR, Levenick KM, et al. Is the “sterile cockpit” concept applicable to cardiovascular surgery critical intervals or critical events? The impact of protocol-driven communication during cardiopulmonary bypass. *J Thorac Cardiovasc Surg.* 2010;139: 312–319. doi:10.1016/j.jtcvs.2009.10.048
25. Dias RD, Scalabrini Neto A. Acute stress in residents during emergency care: a study of personal and situational factors. *Stress.* 2017;20: 241–248. doi:10.1080/10253890.2017.1325866
26. Rieger A, Stoll R, Kreuzfeld S, Behrens K, Weippert M. Heart rate and heart rate variability as indirect markers of surgeons’ intraoperative stress. *Int Arch Occup Environ Health.* 2014;87: 165–174. doi:10.1007/s00420-013-0847-z
27. Böhm B, Rötting N, Schwenk W, Grebe S, Mansmann U. A prospective randomized trial on heart rate variability of the surgical team during laparoscopic and conventional sigmoid resection. *Arch Surg.* 2001;136: 305–310. doi:10.1001/archsurg.136.3.305
28. Lowndes BR, Forsyth KL, Blocker RC, Dean PG, Truty MJ, Heller SF, et al. NASA-TLX Assessment of Surgeon Workload Variation Across Specialties. *Ann Surg.* 2020;271: 686–692. doi:10.1097/SLA.0000000000003058
29. Nadeem R, Agarwal S, Jawed S, Yasser A, Altahmody K. Impact of Cardiopulmonary Bypass Time on Postoperative Duration of Mechanical Ventilation in Patients Undergoing Cardiovascular Surgeries: A Systemic Review and Regression of Metadata. *Cureus.* 2019;11: e6088. doi:10.7759/cureus.6088
30. Shaffer F, Meehan ZM, Zerr CL. A Critical Review of Ultra-Short-Term Heart Rate Variability Norms Research. *Front Neurosci.* 2020;14: 594880. doi:10.3389/fnins.2020.594880
31. Hazlehurst B, McMullen CK, Gorman PN. Distributed cognition in the heart room: how situation awareness arises from coordinated communications during cardiac surgery. *J Biomed Inform.* 2007;40: 539–551. doi:10.1016/j.jbi.2007.02.001
32. Dias RD, Conboy HM, Gabany JM, Clarke LA, Osterweil LJ, Arney D, et al. Intelligent Interruption Management System to Enhance Safety and Performance in Complex Surgical and Robotic Procedures. *OR 20 Context Aware Oper Theaters Comput Assist Robot Endosc Clin Image Based Proc Skin Image Anal* (2018). 2018;11041: 62–68. doi:10.1007/978-3-030-01201-4\_8

33. Dias RD, Conboy HM, Gabany JM, Clarke LA, Osterweil LJ, Avrunin GS, et al. Development of an Interactive Dashboard to Analyze Cognitive Workload of Surgical Teams During Complex Procedural Care. *IEEE Int Interdiscip Conf Cogn Methods Situat Aware Decis Support*. 2018;2018: 77–82. doi:10.1109/COGSIMA.2018.8423995
34. Teo G, Reinerman-Jones L, Matthews G, Szalma J, Jentsch F, Hudson I, et al. Selecting workload and stress measures for performance prediction. *Proc Hum Factors Ergon Soc Annu Meet*. 2017;61: 2042–2046. doi:10.1177/1541931213601989
35. Dias RD, Yule SJ, Zenati MA. Augmented cognition in the operating room. *Digital Surgery*. Cham: Springer International Publishing; 2021. pp. 261–268. doi:10.1007/978-3-030-49100-0\_19