

Packing of \mathbb{R}^n with n -dimensional symmetric convex bodies

Sanket Biswas

Faculty of Mathematics,
University of Waterloo, Waterloo, ON, Canada
and
Department of Chemical Engineering,
SLIET Longowal, Punjab, India

GCT/1940108
1940108@sliet.ac.in

August 22, 2022



UNIVERSITY OF
WATERLOO

FACULTY OF MATHEMATICS
Department of
Pure Mathematics



Contents

- 1 About the Internship Faculty
 - Overview
 - Research Profile
- 2 Introduction and Motivation
 - Basic Definitions and Terminologies
 - Types of OPPs
- 3 Overview of Our Research
 - Background
 - Results and Conjectures



Overview



- It is the only dedicated Faculty of Mathematics in North America.
- It connects three areas central to our millennium: Mathematics, Statistics, and Computer Science.
- It possesses celebrated faculty from around the world.
- Has access to institutes such as the Perimeter Institute, the Institute for Quantum Computing, and the Cybersecurity and Privacy Institute.
- Has strong connections with industry leaders like Google Canada, OpenText, Maplesoft, and multiple insurance companies.



Research Profile

- ➊ **Applied Mathematics:** This department is at the forefront of a number of exciting research areas including Control and Dynamical System, Fluid Mechanics, Mathematical Medicine and Biology, Mathematical Physics and Scientific Computing.
- ➋ **Combinatorics & Optimisation:** This department is doing cutting-edge research in algebraic combinatorics, continuous optimization, cryptography, discrete optimization, graph theory and quantum computing.
- ➌ **Computer Science:** The David R. Cheriton School of Computer Science is engaged in research on an incredible range of computer science and information systems topics. About 24% of all university spin-off companies in Canada are from the University of Waterloo.
- ➍ **Pure Mathematics:** This department is home to an exciting, active research environment focused on core disciplines within pure mathematics, including algebra, analysis, geometry and topology, mathematical logic and number theory.
- ➎ **Statistics & Actuarial Science:** This department is working in a variety of fields in statistics, probability, actuarial science and quantitative finance.

Contents

- 1 About the Internship Faculty
 - Overview
 - Research Profile
- 2 Introduction and Motivation
 - Basic Definitions and Terminologies
 - Types of OPPs
- 3 Overview of Our Research
 - Background
 - Results and Conjectures



Basic Definitions and Terminologies

- A *convex body* is a compact convex subset of the n -dimensional Euclidean space \mathbb{R}^n with nonempty interior.
- An *arrangement* \mathcal{A} in \mathbb{R}^n is a collection of convex bodies in \mathbb{R}^n .
- A *packing* \mathcal{P} is an arrangement with objects (elements) having mutually disjoint interiors.
- The *density* of a packing \mathcal{P} in \mathbb{R}^n is given by the limit

$$d(\mathcal{P}) := \lim_{r \rightarrow \infty} \frac{\text{vol}(\mathcal{P} \cap B^n(r))}{\text{vol}(B^n(r))},$$

where $B^n(r)$ is the n -dimensional open ball centred at the origin having radius r , and for a measurable subset $C \in \mathbb{R}^n$, $\text{vol}(C)$ denotes the *volume* of C .



Basic Definitions and Terminologies Contd...

- The *bounded density* of a packing \mathcal{P} in a *bounded domain* D of \mathbb{R}^n is given by

$$d_D(\mathcal{P}) := \frac{\text{vol}(\mathcal{P})}{\text{vol}(D)}.$$

- For a convex body C in \mathbb{R}^n , the *packing density* of C is given by

$$\delta(C) := \max\{d(\mathcal{P}) : \mathcal{P} \text{ is a packing in } \mathbb{R}^n \text{ with congruent copies of } C\},$$

and for a bounded domain D of \mathbb{R}^n , the *bounded packing density* of C is given by


$$\delta_D(C) := \max\{d_D(\mathcal{P}) : \mathcal{P} \text{ is a packing in } D \text{ with congruent copies of } C\}.$$

Objective: To find the optimal packing of \mathbb{R}^n and its bounded domains using different convex bodies. Such problems are known as *object packing problems (OPPs)*.



ICP, GCP, and EPP

- An *identical circle packing (ICP)* is an optimized non-overlapping arrangement of n identical circles inside a container set such as a circle, square, or a general rectangle.
- A *general circle packing (GCP)* is an optimized non-overlapping arrangement of n arbitrary circles inside a container set.
- An *ellipse packing problem (EPP)* is an optimized non-overlapping arrangement of n ellipses with arbitrary size and orientation inside a container set.



Contents

1 About the Internship Faculty

- Overview
- Research Profile

2 Introduction and Motivation

- Basic Definitions and Terminologies
- Types of OPPs



3 Overview of Our Research

- Background
- Results and Conjectures

Background of Our Research

- The problem of finding the optimal packing of the n -dimensional Euclidean space \mathbb{R}^n using congruent copies of an n -dimensional sphere has received considerable attention in the literature.
- The following table shows the values of n for which $\delta(B^n)$ is known, along with the values of $\delta(B^n)$.

n	2	3	4	5	6	7	8	24
$\delta(B^n)$	$\frac{\pi}{2\sqrt{3}}$	$\frac{\pi}{\sqrt{18}}$	$\frac{\pi^2}{16}$	$\frac{\pi^2}{15\sqrt{2}}$	$\frac{\pi^3}{48\sqrt{3}}$	$\frac{\pi^3}{105}$	$\frac{\pi^4}{384}$	$\frac{\pi^{12}}{12!}$



UNIVERSITY OF
WATERLOO

FACULTY OF MATHEMATICS
Department of
Pure Mathematics



Results and Conjectures

- We are interested in finding the optimal packing of \mathbb{R}^n using congruent copies of an n -dimensional ellipsoid—a convex body of the form

$$E^n[\mathbf{p}, \mathbf{a}] := \{\mathbf{x} \in \mathbb{R}^n : |\mathbf{a} \cdot (\mathbf{x} - \mathbf{p})| \leq 1\},$$

where $\mathbf{p} \in \mathbb{R}^n$ is the centre of the ellipsoid and $\mathbf{a} = (a_1, a_2, \dots, a_n) \in \mathbb{R}^n$ is the coefficient vector such that, $1/a_i$ is the length of the semi-axis along the i direction.

- **Result:** The packing density is independent of the centre of the ellipsoid, and hence we can denote it by just $\delta(E^n[\mathbf{a}])$.
- But, is this density also independent of the coefficient vector \mathbf{a} ? We suspect that the answer is YES for any dimension n !!!
- **Conjecture:** For any dimension n and for any coefficient vector \mathbf{a} ,

$$\delta(E^n[\mathbf{a}]) = \delta(E^n[\mathbf{1}]) = \delta(B^n),$$

where $\mathbf{1} = (1, 1, \dots, 1)$.



UNIVERSITY OF
WATERLOO

FACULTY OF MATHEMATICS
Department of
Pure Mathematics




More on the Conjecture

If our conjecture holds true, then we would have the following:


n	2	3	4	5	6	7	8	24
$\delta(E^n)$	$\frac{\pi}{2\sqrt{3}}$	$\frac{\pi}{\sqrt{18}}$	$\frac{\pi^2}{16}$	$\frac{\pi^2}{15\sqrt{2}}$	$\frac{\pi^3}{48\sqrt{3}}$	$\frac{\pi^3}{105}$	$\frac{\pi^4}{384}$	$\frac{\pi^{12}}{12!}$





References


 Berni Julian Alder and Thomas Everett Wainwright.
Phase transition for a hard sphere system.

The Journal of chemical physics, 27(5):1208–1209, 1957.

 John D Bernal.
A geometrical approach to the structure of liquids.
Nature, 183(4655):141–147, 1959.

 E. G. Birgin, R. D. Lobato, and J. M. Martínez.
Packing ellipsoids by nonlinear optimization.
J. Global Optim., 65(4):709–743, 2016.

 E. G. Birgin, R. D. Lobato, and J. M. Martínez.
A nonlinear programming model with implicit variables for packing ellipsoids.
J. Global Optim., 68(3):467–499, 2017.

 Ignacio Castillo, Frank J. Kampas, and János D. Pintér.
Solving circle packing problems by global optimization: numerical results and industrial applications.
European J. Oper. Res., 191(3):786–802, 2008.



UNIVERSITY OF
WATERLOO

FACULTY OF MATHEMATICS
Department of
Pure Mathematics

